

Original Research Article

“Response of Phosphorus and Molybdenum on growth and yield of summer groundnut (*Arachis hypogea* L.)”

ABSTRACT

The experiment was carried out during *Zaid* season, 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh. The experiment was laid out in Randomised Block Design with ten treatments which are replicated thrice with three levels of application of Phosphorus 20, 40, 60 kg ha⁻¹ and three levels of foliar application of Molybdenum 0.05, 0.15, 0.25% and control. Result revealed that higher plant height (67.83 cm), plant dry weight (35.27g), number of pods plant⁻¹ (25.37), kernel yield (2.14 t ha⁻¹), halum yield (3.23t ha⁻¹), gross return (INR 137458.00 ha⁻¹), net return (INR 90113.60 ha⁻¹) and B:C ratio (1.90) was found in treatment (T8)with the application of Phosphorus at 60 kg ha⁻¹ along with Molybdenum 0.15% whereas number of nodules plant⁻¹ (103.67) was obtained by the application of Phosphorus at the rate 60 kg ha⁻¹ along with Molybdenum 0.25% in treatment (T9).

Keywords: Groundnut, phosphorus, molybdenum, growth, yield.

INTRODUCTION

Groundnut is a leguminous plant, valued for its high-oil edible seeds and as such, it is the fourth most important source of edible oil and a third most important source of vegetable protein in the world. Groundnut is an excellent source of plant nutrients contains 45-50% oil, 27-33% protein as well as essential minerals and vitamins. They play an important role in the dietary requirements of resource poor women and children and haulms are used as livestock feed. Groundnut oil is composed of mixed glycerides and contains a high proportion of unsaturated fatty acids, in particular, oleic (50-65%) and linoleic acid (18-30%).

Globally, Groundnut covers 295 lakh hectares with the production of 487 lakh tones with the productivity of 1647 kg ha⁻¹. With annual all-season coverage of 55.6 lakh hectares, globally, India ranks first in Groundnut acreage and is the second largest producer of Groundnut in the world with 101 lakh tones with a productivity of 1816 kg ha⁻¹ in 2020-21. Groundnut is cultivated in one or more (kharif, rabi and summer) seasons, but nearly 80% of acreage and production comes from kharif crop (**FAOSTAT 2021**). India is one among the top three producing countries of Groundnut in the world. It ranks second next to China (37% contribution to the world Groundnut production). Nearly 15% of Groundnut production is contributed by India to the world production during 2018. Nigeria ranks third in world Groundnut production with 6% contribution.

Phosphorus is the second major essential nutrient element for crop growth and good quality yield. The most obvious effect of P is on the plant root system. The requirement of P in nodulating legumes is higher compared to non-nodulating crops as it plays a significant role in nodule formation and fixation of atmospheric nitrogen. Due to the important role played by P in the physiological processes of plants, application of P to soil deficient in this nutrient leads to increase groundnut yield (**Islam et al. 2013**). It is necessary for the synthesis of protoplasm and chlorophyll, for cell division, and for the growth of meristematic tissues. It aids in the maturation and growth of the seed plant. The production of this vitamin is necessary for is also engaged in oil, protein, acid, and development of glucosinolates that have the amount of oil is increased through hydrolysis. Its deficit result in significant restriction in the plant top and root growth (**Mirvat et al. 2006**).

Molybdenum is important for many enzymatic and physiological processes in plants, with a clear contribution to nitrogen fixation in leguminous crops via root nodules. The nitrate reductase enzyme needs molybdenum to produce, and in legumes, it also participates in symbiotic nitrogen fixation. Without sufficient amounts of molybdenum and iron, the nitrogen-fixing enzyme nitrogenase cannot function and nitrogen fixation cannot take place. Molybdenum, improved groundnut production by increasing the vegetative development and weight of nodular materials. (**Togay et al. 2008**)

MATERIAL AND METHODS

The field experiment was conducted during summer season of 2022 at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom

University of Agriculture, Technology & Sciences, Prayagraj, Uttar Pradesh. The soil in the experimental area was sandy loam with pH (7.7), organic carbon (0.57%), available N (171.48 kg ha⁻¹), available P (27.0 kg ha⁻¹), and available K (291.2 kg ha⁻¹). The experiment was laid out in Randomised Block Design with ten treatments which are replicated thrice with three levels of application of Phosphorus 20, 40, 60 kg ha⁻¹ and three levels of foliar application of Molybdenum 0.05, 0.15, 0.25% and control. Treatment combinations are T1: Phosphorus 20 kg ha⁻¹ + Molybdenum 0.05%, T2: Phosphorus 20 kg ha⁻¹ + Molybdenum 0.15%, T3: Phosphorus 20 kg ha⁻¹ + Molybdenum 0.25%, T4: Phosphorus 40 kg ha⁻¹ + Molybdenum 0.05%, T5: Phosphorus 40 kg ha⁻¹ + Molybdenum 0.15%, T6: Phosphorus 40 kg ha⁻¹ + Molybdenum 0.25%, T7: Phosphorus 60 kg ha⁻¹ + Molybdenum 0.05%, T8: Phosphorus 60 kg ha⁻¹ + Molybdenum 0.15%, T9: Phosphorus 60 kg ha⁻¹ + Molybdenum 0.25%, T10: RDF: 20-40-60 NPK kg ha⁻¹ (Control) are used. Seeds are sown at a spacing of 30cm×10cm according to a seed rate of 90 kg/ha. The recommended dose of nitrogen (20 kg/ha) and potassium (60 kg/ha) was applied as basal dose just before sowing and phosphorus, molybdenum (Foliar spray at 25 DAS) were applied as per the treatments. Urea, SSP and MOP were taken as fertilizer sources of N, P and K respectively.

RESULT AND DISCUSSION

GROWTH ATTRIBUTES

Plant height (cm) The data revealed that significant and maximum plant height (67.83 cm) was recorded in treatment 8 [Phosphorus (60 kg ha⁻¹) + Molybdenum (0.15%)]. However, treatment 7 and treatment 9 were found to be statistically at par with treatment 8 in table no.1. Significant and maximum plant height was recorded by the application of Phosphorus might be due to P application may be attributable to the fact that P is known to help in the development of more extensive root system and thus enables plants to absorb more water and nutrients from depth of the soil. Similar result was reported by **Kabir *et al.* (2013)**. Further, foliar application of molybdenum has significant effect on the traits such as plant height and the biological performance. The result were in conformity with those of **Manjili *et al.* (2014)**.

Number of nodules plant⁻¹ The data revealed that significant and maximum number of nodules plant⁻¹ (107) was recorded in treatment 9 [Phosphorus (60 kg ha⁻¹) + Molybdenum (0.25%)]. treatment 8 and treatment 7 were found to be statistically at par with treatment 8. Phosphorus fertilizer is required for plant growth and development as it helps in root development and also serves as energy source for the rhizobium which in turn may lead to increased nodule formation thereby enhancing N₂ fixation. Similar result was reported by **Badar et al. (2015)** Molybdenum can play a vital role in increasing the nitrogen fixation process by Rhizobium and is responsible for the formation of nodule tissue and increase in N₂ fixation. Molybdenum is a constituent of the nitrogenase enzyme and every bacterium, which fixes nitrogen, need molybdenum during the fixation process reported by **Hirpara et al. (2019)**

Plant dry weight (g) The data revealed that significant and maximum plant dry weight (35.27 g) was recorded in treatment 8 [Phosphorus (60 kg ha⁻¹) + Molybdenum (0.15%)]. treatment 7 and treatment 9 were found to be statistically at par with treatment 8 in table 1. Significant and maximum dry weight by the application of Phosphorus might be due to phosphorus is known to help developing a more extensive root system and thus enabling plants to extract water and nutrients from more depth and this could enhance the plants to produce more assimilates which was reflected in high biomass. Similar result was reported by **Gobarah et al.(2006)**. The improvement in nodulation might have resulted in higher amount of nitrogen fixation and there by better vegetative growth and dry matter production. The results were similar with findings of **Balla et al. (2020)**.

YIELD ATTRIBUTES

Number of pods plant⁻¹ The data revealed that significant maximum number of pods plant⁻¹ (25.37) was recorded in treatment 8 [Phosphorus (60 kg ha⁻¹) + Molybdenum (0.15%)]. However, treatment 9 and treatment 7 were found to be statistically at par with treatment 8 in table 2. The increased number of Pods/Plant might be due to effect of phosphatic fertilizer that increased nutrient availability, resulting in better nutrient absorption and crop growth. Similar result was reported by **Sagar et al. (2020)**.

Kernel yield (t ha⁻¹) The data revealed that significant higher kernel yield (2.14 t ha⁻¹) was recorded in treatment 8 [Phosphorus (60 kg ha⁻¹) + Molybdenum (0.15%)].

treatment 9 and treatment 7 were found to be statistically at par with treatment 8 in table 2. Increasing phosphorus fertilizers rates increased all yield components. Such favourable effects on yield and yield attributes could be due to the stimulation effects of P on number of nodules and nitrogen activity which in turn reflected positively on groundnut yield attributes. Similar result was reported by **Vali et al. (2020)**.

Halum yield (t ha⁻¹) The data revealed that significant higher haulm yield (3.23 t ha⁻¹) was recorded in treatment 8 [Phosphorus (60 kg ha⁻¹) + Molybdenum (0.15%)]. However, treatment 9 and treatment 7 were found to be statistically at par with treatment 8 in table 2. The improvement of dry matter was mainly responsible for the increased stover yield. The effect of phosphorus will induce the uptake ability of roots to nutrients and positive increase in the yield parameters because of improving the root system as a source-sink relationship to the reproductive part (shoot). Similar result was reported by **Mouri et al. (2018)**. Molybdenum increased nitrogen fixation which effects plant growth rate and metabolism which results in higher halum yields. Similar result was reported by **Bhagiya et al. (2005)**.

ECONOMICS Highest gross return (INR 137458.00 ha⁻¹), net return (INR 86126.60 ha⁻¹) and Benefit cost ratio (1.90) was obtained in the treatment 8 (Phosphorus 60 kg ha⁻¹ + Molybdenum 0.15%) in table 3.

REFERENCES

1. **Bhagiya, S. R., Polara, K. B., & Polara, J. V. (2005)**. Effect of B and Mo on yield, quality and nutrient absorption by groundnut. *Advances in Plant Sciences*, 18(2), 803.
2. **Badar, R., Nisa, Z., & Ibrahim, S. (2015)**. Supplementation of P with rhizobial inoculants to improve growth of Peanut plants. *International Journal of Applied Research*, 1(4), 19-23.
3. **Balla, H., Singh, V., Tiwari, D., & Shaik, M. A. (2020)**. Effect of boron and molybdenum on growth rate and yield of groundnut (*Arachis hypogea L.*). *Journal of Pharmacognosy and Phytochemistry*, 9(6), 1416-1419.
4. **FAOSTAT, F. (2021)**. New food balances. *FAOSTAT*. Available via *FAO*. Accessed, 25.

- 5. Gobarah, M. E., Mohamed, M. H., & Tawfik, M. M. (2006).** Effect of phosphorus fertilizer and foliar spraying with zinc on growth, yield and quality of groundnut under reclaimed sandy soils. *J. Appl. Sci. Res*, 2(8), 491-496.
- 6. Hirpara, D. V., Sakarvadia, H. L., Jadeja, A. S., Vekaria, L. C., & Ponkia, H. P. (2019).** Response of boron and molybdenum on groundnut (*Arachis hypogea L.*) under medium black calcareous soil. *Journal of pharmacognosy and phytochemistry*, 8(5), 671-677.
- 7. Islam, A. M. (2013).** Effect of phosphorus, calcium and boron on the growth and yield of groundnut (*Arachis hypogea L.*). *International Journal of Bio-science and Bio-technology*, 5(3).
- 8. Kabir, R., Yeasmin, S., Islam, A. K. M. M., & Sarkar, M. R. (2013).** Effect of phosphorus, calcium and boron on the growth and yield of groundnut (*Arachis hypogea L.*). *International Journal of Bio-science and bio-Technology*, 5(3), 51-60
- 9. Mirvat, E.G., Magda, H.M., and Tawfik, M.M. (2006)** Effect of phosphorus fertilizer and foliar spraying with zinc on growth, yield and quality of groundnut under reclaimed sandy soils. *Journal of Applied Science and Research*. 2(8): 491-496.
- 10. Manjili, M. J., Bidarigh, S., & Amiri, E. (2014).** Study the effect of foliar application of nano chelate molybdenum fertilizer on the yield and yield components of peanut. *Egyptian Academic Journal of Biological Sciences, H. Botany*, 5(1), 67-71.
- 11. Mouri, S. J., Sarkar, M. A. R., Uddin, M. R., Sarker, U. K., & Hoque, M. M. I. (2018).** Effect of variety and phosphorus on the yield components and yield of groundnut. *Progressive Agriculture*, 29(2), 117-126.
- 12. Togay, Y., Togay, N., & Dogan, Y. (2008).** Research on the effect of phosphorus and molybdenum applications on the yield and yield parameters in lentil (*Lens culinaris Medic.*). *African Journal of Biotechnology*, 7(9).
- 13. Vidya-Sagar, D. R. M. S., Dawson, J., & Reddy, R. U. K. (2020).** Effect of phosphorus and gypsum on growth, yield and economics of groundnut (*Arachis hypogea L.*). *Int. J. Curr. Microbiol. App. Sci*, 9, 1635-1638.

14. Vali, G. M., Singh, S., Sruthi, D. S. V., Hinduja, N., Talasila, V., & Tiwari, D. (2020). Effect of phosphorus and zinc on growth and yield of Summer groundnut (OF SUMMER GROUNDNUT (*Arachis hypogea* *Arachis hypogea* *Arachis hypogea* L.)).

UNDER PEER REVIEW

Table 1 Effect of phosphorus and molybdenum on growth attributes of groundnut.

S.no.	Treatment combinations	Plant height (cm)	No. of nodules plant ⁻¹	Plant dry weight (g)
1.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.05%	60.67	91.33	26.83
2.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.15%	59.00	91.67	27.53
3.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.25%	62.77	92.33	30.17
4.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.05%	63.90	98.00	28.67
5.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.15%	61.20	100.67	31.20
6.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.25%	63.73	98.00	31.10
7.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.05%	66.50	101.33	34.23
8.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.15%	67.83	100.67	35.27
9.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.25%	65.53	103.67	32.73
10.	Control RDF (20:40:60 kg ha ⁻¹)	54.43	71.33	26.33
	F test	S	S	S
	SEm.(±)	2.38	4.07	1.64
	CD (p=0.05)	6.65	12.08	4.88

Table 2 Effect of phosphorus and molybdenum on yield attributes of groundnut.

Treatment Combinations		No. of pods plant ⁻¹ (No.)	Halum Yield t/ha	Kernel Yield t/ha
1.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.05%	18.07	2.30	1.51
2.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.15%	18.27	2.17	1.50
3.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.25%	19.43	2.50	1.59
4.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.05%	20.50	2.53	1.65
5.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.15%	20.83	2.67	1.67
6.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.25%	22.80	2.83	1.69
7.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.05%	23.20	3.10	2.05
8.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.15%	25.37	3.23	2.14
9.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.25%	24.47	3.17	2.10
10.	Control RDF (20:40:60 kg ha ⁻¹)	21.40	2.13	1.42
F test		S	S	S
SEm(±)		0.76	0.14	0.13
CD (p=0.05)		2.26	0.42	0.39

Table 3 Effect of phosphorus and molybdenum on Economics of groundnut

	Treatment Combination	Cost of cultivation (INR/ha)	Gross returns (INR/ha)	Net returns (INR/ha)	B:C ratio
1.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.05%	45854.80	97047.00	51192.00	1.12
2.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.15%	47054.80	96249.00	49194.00	1.05
3.	Phosphorus 20kg ha ⁻¹ + Molybdenum 0.25%	48254.80	102622.00	54367.20	1.13
4.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.05%	45999.60	105906.00	59906.40	1.30
5.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.15%	47199.60	107898.00	60698.40	1.29
6.	Phosphorus 40kg ha ⁻¹ + Molybdenum 0.25%	48399.60	109393.00	60993.40	1.26
7.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.05%	46144.40	131685.00	85540.60	1.85
8.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.15%	47344.40	137458.00	90113.60	1.90
9.	Phosphorus 60kg ha ⁻¹ + Molybdenum 0.25%	48544.40	134671.00	86126.60	1.77
10.	Control RDF (20:40:60 kg ha ⁻¹)	44510.00	90975.00	46175.40	1.03

UNDER PEER REVIEW