

Estimation of genetic variability parameters and parent-offspring correlation, regression in groundnut (*Arachis hypogaea* L.)

Abstract:

Genetic variability is major component which helps in selecting better genotypes under different environmental conditions, with this aim an experiment was conducted to understand the genetic potential, heritability, genetic advance and traits association of yield contributing characters for F₆ and F₇ families derived from the cross GKVK-6 × KCG-2 at the University of Agricultural Sciences, Bangalore in an augmented block design along with three checks *viz.*, TMV-2, KCG-6 and KCG-2 during summer and *Kharif* 2017. Results from analysis of variance (ANOVA) revealed that highly significant differences were observed for all the characters studied. High genetic variability was observed for major yield contributing characters like the number of pods per plant (g), pod yield (g), kernel yield per plant (g), SMK% (sound mature kernel *per cent*), SCMR (SPAD chlorophyll meter reading) and SLA (specific leaf area) (cm²/g). Narrow difference between GCV (genotypic coefficient of variation) and PCV (phenotypic coefficient of variation) was observed for pods per plant, pod yield, SCMR and SLA and high heritability coupled with moderate genetic advance *per cent* mean was recorded for pod yield, SCMR and SLA indicating the involvement of additive gene action in controlling these traits. Three superior families were noticed with more pods plant⁻¹, high pod yield plant⁻¹, high kernel yield⁻¹, high shelling (%), high SMK (%), high SCMR value, and low SLA value. Further, these superior families also revealed the presence of high parent offspring regression and intergeneration correlation, implying increased efficiency of selection for most of the traits considered and these were identified to be the important characters that could be used in selection for yield and the selected families will be tested further for yield stabilization.

Keywords: GCV, PCV, Heritability, GAM, water use efficiency, Groundnut.

Introduction

Groundnut (*Arachis hypogaea* L.) is a self-pollinated, cleistogamous annual herb belonging to the family Leguminaceae with a chromosome number of $2n=40$. It is the most important oilseed crop of India and the world in terms of area and production. It is vernacularly known by different names as peanut, monkey nut, earthnut and pigmy nut. Cultivated groundnut is classified into two subspecies, subs. *fastigiata* and subs. *hypogaea*. The subs. *fastigiata* contains four botanical varieties, var. *vulgaris*, var. *fastigiata*, var. *peruviana*, and var. *aequatoriana*. The subs. *hypogaea* contain two varieties, var. *hypogaea* and var. *hirsuta*. Each of those botanical types has the contrasting plant, pod and seed characteristics (Krapovickas and Gregory, 1994). Yield is a complex trait, governed by many traits and there is ample evidence to show that selection directly for grain yield in plants is not easy. Since the economic a part of groundnut may be a pod that's developed under the soil, prediction of its performance supported aerial morphological characters is nearly difficult (Weiss, 2000). Gain under direct selection for pod yield in groundnut is low and slow as pod yield isn't only polygenically controlled but also influenced by its component characters (Alam *et al.*, 1985). The knowledge of existing variability and degree of association between pod yield and its contributing characters and their relative contribution to pod yield is important for developing high yielding genotypes. Heritability and genetic advance is a useful tool for breeders in determining the amount of genetic variation present in crops. Correlation analysis is useful to work out the magnitude of association among the characters and their relative contribution to pod yield. More than 70% of the groundnut growing area falls under arid and semi-arid regions, where groundnut frequently experiences drought stresses for various durations and intensities (Reddy *et al.*, 2003). Yield losses attributable to drought are highly variable depending on timing, intensity and duration in addition to other location-specific environmental stress factors like high irradiation and temperature (Kambiranda *et al.*, 2011). Losses in yield that are due to water stress can be tackled partially by developing varieties that are better adapted to water scares conditions (Ravi *et al.*, 2011).

Water use efficiency is defined as the amount of water utilized in transpiration to produce dry matter during a specific growth period. Water Use Efficiency (WUE) is one such trait which will contribute to productivity under drought. There are numerous easily measurable traits having a high correlation with WUE that are identified as surrogate traits. Traits that have practical advantages for WUE are specific leaf area (SLA) and soil-plant analysis development and SPAD chlorophyll meter reading (SCMR). Nageswara Rao *et al.* (2001) and Sheshashayee *et al.* (2006) reported a direct correlation between SLA and carbon isotope discrimination, and an indirect correlation with WUE in groundnut, suggesting that SLA can be used as a surrogate trait to measure WUE in groundnut. SCMR has been used effectively to determine leaf nitrogen content non-destructively in several crops including groundnut (Nageswara Rao *et al.*, 2001) reported significant and high negative inter-relationship among SLA and SCMR. Zhang *et al.* (2022), identified the genotypes PI 502120 and AU-NPL 17 as water spender genotypes as they showed high yield, $\Delta^{13}\text{C}$, photosynthesis, and stomatal conductance under drought. Upadhyaya (2002) used SCMR and SLA as surrogate traits for assessing WUE in groundnut mini core germplasm collection and reported a negative correlation between SCMR and SLA. There are many reasons which are attributed to low yield levels *viz.*, lack or non-availability of improved high yielding cultivars, cultivation under shallow soils of low fertility, uneven rainfall distribution, continuous cropping without rotation of crop, low plant population and incidence of foliar diseases and pests are cited as the major limiting factors in most of the groundnut growing regions. Keeping all the above points in view, an effort has been made to study the genetic variability, an association of traits related to WUE with pod yield and its component traits, and to identify superior families performing better under drought stress conditions in F₆ and F₇ generation.

Materials and Methods

Plant material and experimental site

The present study was conducted during summer and *Khariif* 2017 at the experimental field, GKVK, University of Agricultural Sciences, Bangalore. The experimental material for the present study comprised of 12 F₆ and F₇ families along with three checks KCG-6, KCG-2 and TMV-2. The populations were developed by crossing the parents which were contrasting for the traits, SLA, SCMR, pod yield per plant, kernel yield per plant and pods per plant. The selected families from the F₅ generation of the cross were forwarded to the F₆ generation and the same procedure was repeated to rise F₇ generation. Statistical analysis of the mean data was analyzed in WINDOSTAT *version* 8.5 for augmented design and SPSS for descriptive statistics.

Evaluation of plant material

The F₆ plants of the cross, their respective parents and checks (KCG-2, KCG-6 and TMV-2) were raised in a plant to progeny row method in an augmented design during summer 2017 with 5 m row length and 30 cm and 10 cm inter and intra row spacing respectively. Individual F₆ progenies which had mean values higher than the families grand mean and mean more than the checks and the parents for the traits *viz.*, SPAD chlorophyll meter reading (SCMR), pod yield, kernel yield, shelling *per cent* and sound mature kernel (SMK) *per cent* and the low mean values for specific leaf area (SLA) were selected and sown in a plant to progeny row for F₇ evaluation of yield and water use efficiency-related traits as the salient features of parents and checks were given in (Table 1).

Data Recording

The data on ten morphological characters namely days to 50% flowering (days to first flowering), plant height (cm), primary branches per plant, pods per plant, pod yield per plant (g), kernel yield per plant (g), shelling percentage, sound mature kernel (SKM) *per cent*, specific leaf area (SLA) and SPAD chlorophyll meter reading (SCMR) were recorded. Shelling percentage: was calculated by using the following formulae. Kernel weight (g)/Pod weight (g) ×100, Sound mature kernel (SMK) *per cent*: was calculated by using the following formulae. Number of well-developed kernels/Total number of kernels ×100.

Water use efficient traits

Specific leaf area (SLA): The second or third fully expanded leaf of the main axis was collected in butter paper covers at 65 days after sowing. The leaf area was measured using a leaf area meter. Then the leaves were kept in an oven at 70 °C for 3 days. The dry weight of the leaf was accurately measured using a sensitive balance. SLA was computed using the formula given below and expressed as cm²/g.

SPAD Chlorophyll Meter Reading (SCMR): Leaf nitrogen content normally influences the leaf chlorophyll content. A device has been developed by Minolta company, New Jersey USA (SPAD-502) which measures the light attenuation at 430 nm (the peak wavelength for chlorophyll a and chlorophyll b) and 750 nm (near-infrared) with no transmittance. The unit less value measured by the chlorophyll meter is termed SCMR (SPAD chlorophyll meter reading), which provides information on the relative amount of leaf chlorophyll. The SPAD meter (soil-plant analytical development) is a simple handheld instrument, which operates with a DC power of three Volts.

The second or third leaf from the apex was selected to record the SCMR. The selected leaf of groundnut was clamped avoiding the midrib region into the sensor head of the SPAD meter. A gentle stroke was given to record the SPAD reading and the average of such four strokes per leaflet was considered. Since groundnut is a tetra-foliolate leaf, SCMR was recorded in all four leaflets and the average value was recorded. The SCMR was recorded under normal sunlight between 9.00 am to 4.00 pm.

Narrow sense heritability estimates (based on regression co-efficient)

Narrow sense heritability estimates were performed based on the regression of F₇ on F₆, using the following formula (Cahaner and Hiller, 1980).

$$b (F_7, F_6) = \frac{\text{Covariance of } F_7 F_6}{\text{Variance of } F_6}$$

$$h^2 (ns) = b (F_7.F_6) \times 0.85$$

Regression co-efficient between F_6 and F_7 , generations was estimated as heritability value using multiplicative factor of 6/7. Heritability estimates were computed for the selected families for all the characters.

Results and Discussion

Analysis of variance (ANOVA)

Analysis of variance was carried out for growth, traits related to water use efficiency, yield and its component characters and is presented in Table 1 Figure 1. Mean sum of squares of families exhibited highly significant difference for all the traits. Further, mean sum of squares of checks *versus* families exhibited greater significant difference for all the characters like days to first flowering, plant height, branches per plant, SCMR, SLA, pod yield per plant, kernel yield per plant and sound mature kernel *per cent* except SMK *per cent*. Hence the choice of material for the study is suitable and further selections could be practiced.

Estimates of genetic variability parameters

The genetic variability parameters were analyzed and presented in the Table 2. Wide range of variation was observed for the traits SLA (132 to 180.20, 98 to 156 cm^2/g with the mean of 153.47, 122.50), pods per plant (17.50 to 44.16, 18 to 49 with the mean of 28.30, 32.58), pod yield per plant (15.33 to 31.71, 13.74 to 39.25 with the mean of 22.13, 25.81), kernel yield per plant (8 to 19.46, 7.42 to 35 with the mean of 13.41 19.18), shelling% (50 to 65.29, 40.98 to 60.22 with the mean of 58.82, 52.46), SMK% (8.55 to 82, 31.77 to 60.11 with the mean of 58.49, 48.09) in F_6 and F_7 generation respectively. High GCV and PCV with narrow difference between GCV and PCV was observed for the traits Primary branches plant^{-1} (12.77-15.11, 13.46-15.26), Pods plant^{-1} (19.29-24.52, 17.81-25.20), Pod yield $\text{plant}^{-1}(\text{g})$ (19.72-24.14, 17.91-19.56), Kernel yield $\text{plant}^{-1}(\text{g})$ (23.08-29.09, 26.35-31.89). High heritability coupled with high GAM was observed for the traits DFF, Plant height, Primary branches plant^{-1} , SCMR, SLA(cm^2/g), Pods plant^{-1} ,

Pod yield plant⁻¹(g), Kernel yield plant⁻¹(g), Shelling (%) and SMK (%). Chauhan *et al.* (2022) reported high GCV, PCV for shelling %, kernel yield per plant and pod yield per plant in BC₁F₁, BC₁F₂ and BC₁F₃ populations. Similar results were reported by Meta and Monpara (2010), Makhan *et al.* (2003); Golakia *et al.* (2005); John *et al.* (2005); Rao *et al.* (2012) in groundnut. Sridevi *et al.* (2022). Similar results of High heritability coupled with high genetic advance over mean were reported by Nath *et al.* (2002) and Golakia *et al.* (2005) for plant height, pods per plant and pod yield per plant which indicates that these characters are under the influence of additive genetic control and selection will be effective and may rapidly contribute to yield. Rao *et al.* 2012 for pods per plant, Padmaja *et al.* 2013a for pod yield per plant, Zaman *et al.* 2011 and Rao *et al.* 2012 for kernel yield per plant, Reddy *et al.* 1991; Venkataravana *et al.* 2001 for SMK *per cent*.

Estimation of correlation coefficients

Correlation coefficient is an essential tool. Correlation studies between yield and its component traits would help plant breeders to enhance crop growth and yield of crop. In the present study phenotypic correlation between pod yield per plant with component characters and also with physiological traits were studied in both F₆ and F₇ generations.

Phenotypic correlation coefficients for traits related to water use efficiency, pod yield and yield and its component traits are presented in Table 3. Phenotypic correlation coefficient revealed that pod yield per plant had significant positive correlation with SCMR (0.65, 0.52), pods per plant (0.94, 0.93), kernel yield per plant (0.76, 0.92), DFF (0.20, 0.39) and plant height (0.46, 0.42), however, Pod yield per plant had significant negative correlation with SLA (-0.43, -0.36), and shelling per cent (-0.42, -0.55) in F₆ and F₇ generation. This indicated that improvement in SCMR, pods per plant, kernel yield per plant and plant height will lead to improvement in yield. These results are in accordance with the reports of Sridevi *et al.* (2022), Mukhtar *et al.* (2011), Shoba *et al.* (2012). Koolachart *et al.* (2013) and Thakur *et al.* (2013). This indicates that selection of traits for low SLA leads to improvement in yield.

Shelling percentage showed significant positive correlation with kernel yield per plant (0.41, 0.35) in F₆ and F₇ generation indicated the shelling *per cent* could be improved by selecting more number of pod per plant with bold kernels. Similar result was noticed by Nandini *et al.* (2012). SLA exhibited significant negative correlation with SCMR (-0.67, -0.50), pods per plant (-0.39, -0.48), pod yield per plant (-0.43, -0.36) and kernel yield per plant (-0.43, -0.55) in F₆ and F₇ generation, suggesting the improvement of yield and water use efficiency could be done by selecting families that show low SLA. Rekha (2005) and Reddy *et al.* (2003) also reported similar kind of outcomes. SCMR exhibited highly significant positive association with pods per plant (0.69, 0.58), pod yield per plant (0.65, 0.52), kernel yield per plant (0.50, 0.48) and SMK *per cent* (0.48, 0.29) in F₆ and F₇ generation. Therefore, selection of genotypes with high SCMR offers the scope for simultaneous improvement of yield and water use efficiency in groundnut as higher SCMR indicate high photosynthetic efficient genotypes. The results are in agreement with the reports of Songsri *et al.* (2009) and Rekha (2005). John *et al.*, 2005 for SCMR. This shows that selection of families whose SCMR value were higher than checks, which indirectly lead to improvement of yield in groundnut as high SCMR indicates high photosynthetic efficient genotypes with high water use efficiency. The reports of Nageshwar Rao *et al.* (2001), Talwar *et al.* (2004), Rekha (2005), John *et al.* (2005) and Krishnamurthy *et al.* (2007) also confirmed the same association in groundnut.

Estimation of parent-offspring correlation, regression and heritability in F₆ and F₇ generations.

Data presented in the Table 4 indicates the presence of significant positive correlation coupled with higher magnitude of parent-offspring regression observed for the traits like days to first flowering ($r=0.61$, $b=0.44$), plant height ($r=0.71$, $b=0.56$), SCMR ($r=0.38$, $b=0.70$), pods per plant ($r=0.60$, $b=0.56$), pod yield per plant ($r=0.53$, $b=0.67$) and kernel yield per plant ($r=0.27$, $b=0.21$) respectively. Further high narrow sense heritability was observed for traits like days to first flowering (75.23), plant height (48.31), SCMR (59.79), pods per plant (47.61), and pod yield per plant (57.34) respectively.

Lush (1945) defined heritability in broad and narrow sense and emphasized that characters are subjected to different amount of non-heritable variation. The broad sense heritability includes genotypic variance and phenotypic variance, but genotype variance includes both dominance and additive variance and hence not a reliable index for practicing selections. While narrow sense heritability includes

additive variance and phenotypic variance, hence, additive variance is a reliable index of the total genotypic variance, and selections will be effective for forwarding the lines to next generation. Hence, a comparison was made between narrow sense and broad sense heritability.

Estimation of parent-offspring correlation, regression and heritability exhibited significant positive intergeneration correlation coupled with higher magnitude of parent-offspring regression between F_6 and F_7 generation for the traits like days to first flowering, plant height, SCMR, pods per plant, pod yield per plant and kernel yield per plant, this suggests that these characters could be used as selection criteria for selection of superior families from F_7 generation. Further high broad sense heritability coupled with high narrow sense heritability has been observed for traits like DFF, plant height, SCMR, pods per plant, pod yield per plant, indicated the characters are governed by additive gene action. The higher value of narrow sense heritability suggests that there is predominance of additive gene action in the inheritance of the trait and hence effective selection can be made in the early generation. It helps in estimating the performance of the future generations depending on present generation. Hence, performance of future generations can be estimated depending on the traits having higher value of Narrow sense heritability further improvement of pod yield upon selection could be possible. These findings were supported by Kulkarni *et al.* (1976) and Reddy *et al.* (1993). Overall three superior families were noticed with more pods plant⁻¹, high pod yield plant⁻¹, high kernel yield⁻¹, high shelling (%), high SMK (%), high SCMR value, and low SLA value (Table 5 Figure 2).

Conclusions

Analysis of variance for all the characters studied in both F_6 and F_7 generations revealed highly significant differences among the families suggesting the presence of sufficient amount of variability. Further, genetic variability estimates such as PCV and GCV coupled with high heritability and genetic advance as *per cent* mean for kernel yield per plant, pod yield per plant and sound mature kernel percentage, SLA indicating the presence of sufficient variability and involvement of additive gene action in both F_6 and F_7 generations for these traits. Days to first flowering SCMR exhibited lower PCV and GCV estimates and thus indicating less variation

for this trait. Phenotypic correlation coefficient depicted significant positive association of SCMR, pods per plant and kernel yield per plant with pod yield per plant. SLA exhibited significant negative association with pod yield per plant and SCMR. Therefore, superior genotypes for high yield and water use efficiency were selected depending on the relationship between SLA, SCMR and pod yield. Three superior families were noticed with more pods plant⁻¹, high pod yield plant⁻¹, high kernel yield⁻¹, high shelling (%), high SMK (%), high SCMR value, and low SLA value. Further, these superior families also revealed the presence of high parent offspring regression and intergeneration correlation, implying increased efficiency of selection for most of the traits considered and these were identified to be the important characters that could be used in selection for yield and the selected families will be tested further for yield stabilization.

References

- Alam, M. S., Rahman, A. R. M. S. and Khair, A. B. M. A. (1985). Genetic variability and character association in groundnut (*Arachis hypogaea* L. of Bangladesh]. *Bangladesh Journal of Agriculture (Bangladesh)*.
- Cahaner, A. and Hillel, J. (1980). Estimating heritability and genetic correlation between traits from generations F₂ and F₃ of self-fertilizing species: a comparison of three methods. *TAG. Theoretical and applied genetics. Theoretische und angewandte Genetik*, 58(1), 33-38.
- Chauhan, S., Savithamma, D. L. and Kundu, S. (2022). Genetic variability, heritability and genetic advance for yield and its related traits in BC1F1, BC1F2 and BC1F3 back cross generation of two crosses of groundnut (*Arachis hypogaea* L.).
- Golakia, P. R., Makne, V. G. and Monpara, B. A. (2005). Heritable variation and association in Virginia runner and Spanish bunch group of groundnut. *National Journal of Plant Improvement*, 7(1), 50-53.
- John, K., Vasanthi, R. P., Venkateswarlu, O. and Sudhakar, P. (2005). Genetic variability and correlation studies among F1s and parents in groundnut (*Arachis hypogaea* L.). *Legume Research-An International Journal*, 28(4), 262-267.
- Kambiranda, D. M., Vasanthaiah, H. K., Katam, R., Ananga, A., Basha, S. M. and Naik, K. (2011). Impact of drought stress on peanut (*Arachis hypogaea* L.) productivity and food safety. *Plants and environment*, 1, 249-272.
- Koolachart, R., B. Suriharn, S. Jogloy, N. Vorasoot, S. Wongkaew, C. C. Holbrook, N. Jongrunklang, T. Kesmala. and A. Patanothai (2013). "Relationships between physiological traits and yield components of peanut genotypes with different levels of terminal drought resistance." *SABRAO Journal of Breeding and Genetics*, 45(3), 422-446.

- Krapovickas, A. and Gregory, W. C. (1994). Taxonomia Del Genero " Arachis (leguminosae)". *Bonplandia*, 1-186.
- Krishnamurthy, L., Vadez, V., Devi, M.J., Serraj, R., Nigam, S.N., Sheshshayee, M.S., Chandra, S. and Aruna, R. (2007). Variation in transpiration efficiency and its related traits in a groundnut (*Arachis hypogaea* L.) mapping population. *Field Crops Research*, 103(3), pp.189-197.
- Kulkarni, R. S., Swamy Rao, T., & Virupaksappa, K. (1976). Gene action in Bhindi.
- Lush, J. L. (1945). Heritability of quantitative characters in farm animals. *Proc. 8th Cong. Genet. Hereditas*, 35: 356-375.
- Meta, H. R. and Monpara, B. A. (2010). Genetic variation and trait relationships in summer groundnut, (*Arachis hypogaea* L.). *Journal of Oilseeds Research*, 27(1), 8-11.
- Mukhtar, A. A., Tanimu, B. S., Ibrahim, B. S., Abubakar, I. U. and Babaji, B. A. (2011). Correlations and path coefficients analysis between pod yield and some quantitative parameters in ground nuts (*Arachis hypogaea* L.). *International Journal of Science and Nature*, 2(4), 799-804.
- Nageswara Rao, R. C., Talwar, H. S. and Wright, G. C. (2001). Rapid assessment of specific leaf area and leaf nitrogen in peanut (*Arachis hypogaea* L.) using a chlorophyll meter. *Journal of Agronomy and Crop Science*, 186(3), 175-182.
- Nandini, C., Savithramma, D. L. and Babu, N. N. (2011). Genetic variability analysis for water use efficiency in F8 recombinant inbred lines of groundnut (*Arachis hypogaea* L.). *Current Biotica*, 5(3), 282-288.
- Nath, U. K. and Alam, M. S. (2002). Genetic variability, heritability and genetic advance of yield and related traits of groundnut (*Arachis hypogaea* L.). *Journal of Biological Sciences*, 2(11), 762-764.
- Padmaja, D., Eswari K. B., Rao, B. M. V. and Reddy, S. M. (2013a). Genetic variability, heritability for late leaf spot tolerance and productivity traits in a recombinant inbred line population of groundnut (*Arachis hypogaea* L.). *IOSR Journal of agriculture and veterinary science*, 5, 36-41.
- Rao, V. T., Bhadru, D., Murthy, K. G. K. and Bharathi, D. (2012). Genetic variability and association among the various characters in groundnut (*Arachis hypogaea* L.). *International journal of Applied Biology and Pharmaceutical*, 3, 337-341.

Ravi, K., Vadez, V., Isobe, S., Mir, R. R., Guo, Y., Nigam, S. N., Gowda, M. V. C., Radhakrishnan, T., Bertoli, D. J., Knapp, S. J. and Varshney, R. K. (2011). Identification of several small main-effect QTLs and a large number of epistatic QTLs for drought tolerance related traits in groundnut (*Arachis hypogaea* L.). *Theoretical and Applied Genetics*, 122, pp.1119-1132.

Reddy, G. L. K., Reddy, M. S. S., & Reddy, P. R. (1993). Intergeneration correlation between F3-F4 and F4-F5 generations in some crosses of groundnut. *Crop Improvement*.

Reddy, M. V., Reddy, K. R., Reddy, K. H. P., & Rao, J. A. (1991). Heritability, genetic advance and character association in groundnut over three environments. *Journal of Maharashtra Agricultural Universities (India)*.

Reddy, P. V., Sudhakar, P., Sujatha, D., Babitha, M. and Latha, P. (2003). Transgressive segregation in groundnut. In *Proceedings of National Seminar on Physiological Interventions for Improved Crop Productivity and Quality: Opportunities and Constraints*, Indian Society for Plant Physiology, New Delhi (Vol. 84).

Reddy, P. V., Sudhakar, P., Sujatha, D., Babitha, M. and Latha, P. (2003). Transgressive segregation in groundnut. In *Proceedings of National Seminar on Physiological Interventions for Improved Crop Productivity and Quality: Opportunities and Constraints*, Indian Society for Plant Physiology, New Delhi (Vol. 84).

Rekha, D. (2005). Genetic Analysis and Molecular Characterization for water use efficiency in groundnut. *M.Sc. (Agri.)* thesis submitted to the University of Agricultural Sciences, Bangalore.

Sheshshayee, M. S., Bindumadhava, H., Rachaputi, N. R., Prasad, T. G., Udayakumar, M., Wright, G. C. and Nigam, S. N. (2006). Leaf chlorophyll concentration relates to transpiration efficiency in peanut. *Annals of Applied Biology*, 148(1), 7-15.

Shoba, D., Manivannan, N. and Vindhiyavarman, P. (2012). Correlation and path coefficient analysis in groundnut (*Arachis hypogaea* L.), *Madras Agricultural Journal*, 99(1-3), 18-20.

Songsri, P., Jogloy, S., Holbrook, C. C., Kesmala, T., Vorasoot, N., Akkasaeng, C. and Patanothai, A. (2009). Association of root, specific leaf area and SPAD chlorophyll meter reading to water use efficiency of peanut under different available soil water. *Agricultural water management*, 96(5), 790-798.

Sridevi, S., Meenakumari, B., Manivannan, N. and Ravichandran, V. (2022). Assessment of genetic variability, character association and path analysis of kernel yield and yield components in groundnut (*Arachis hypogaea* L.). *Electronic Journal of Plant Breeding*, 13(2), 476-481.

Talwar, H. S., Nageshwar Rao, R. C., Nigam, S. N. and Wright, G. C. (2004). Leaf Anatomical characteristics associated with Water Use Efficiency in groundnut. In *proceedings of National symposium on enhancing productivity for sustaining food and nutritional security*, NRCG, Junagadh (p. 128).

Thakur, S. B., Ghimire, S. K., Chaudhary, N. K., Shrestha, S. M. and Mishra, B. (2013). Variability in groundnut (*Arachis hypogaea* L.) to *Cercospora* Leaf spot disease tolerance. *International Journal of Life Sciences Biotechnology and Pharm Research*, 2, 254-262.

Upadhyaya, H. D., Bramel, P. J., Ortiz, R. and Singh, S. (2002). Developing a mini core of peanut for utilization of genetic resources. *Crop Science*, 42(6), 2150-2156.

Venkataravana, P., Sheriff, R. A., Kulkarni, R. S., Shankaranarayana, V., & Fathima, P. S. (2000). Correlation and path analysis in groundnut (*Arachis hypogaea* L.). *Mysore Journal of Agricultural Sciences*, 34(4), 321-325.

Weiss, E. A. (2000). *Oilseed crops* (Second Ed.). Blackwell Science Ltd., London.

Zaman, M. A., Tuhina-Khatun, M., Ullah, M. Z., Moniruzzamn, M. and Alam, K. H. (2011). Genetic variability and path analysis of groundnut (*Arachis hypogaea* L.). *The Agriculturists*, 9(1-2), 29-36.

Zhang, Q., Dang, P., Chen, C., Feng, Y., Batchelor, W., Lamb, M. and Sanz-Saez, A. (2022). Tolerance to mid-season drought in peanut can be achieved by high water use efficiency or high efficient use of water. *Crop Science*, 62(5), 1948-1966.

Table 1: Estimation of Analysis of variance (ANOVA)

SV	Generations	df	DF	Plant height (cm)	Primary Branches plant ⁻¹	SCMR	SLA (cm ² /g)	Pods Plant ⁻¹	Pod yield plant ⁻¹ (g)	Kernel yield plant ⁻¹ (g)	Shelling (%)	SMK (%)
Blocks	F ₆	2.00	14.33	13.87	0.40	12.91	52.64	29.31	43.16	40.49	11.06	68.76*
	F ₇	1.00	12.00	28.20	0.53	20.01	54.50	31.52	33.14	39.36	17.69	85.80*
Checks	F ₆	2.00	12.33	15.04	0.64	54.74	71.40	11.47	21.51	34.65	40.66	55.81
	F ₇	2.00	10.66	23.74	1.38	32.90	94.00*	20.20	18.78	14.14	32.18	62.83
Families	F ₆	11.00	22.56**	94.82**	7.59**	180.62**	594.25**	66.16**	139.76**	117.10**	139.94**	91.40**
	F ₇	11.00	33.36**	104.74**	8.33**	229.74**	651.84**	76.44**	104.86**	127.15**	129.38**	98.51**
Checks vs. Families	F ₆	22.00	48.89**	231.42**	33.45**	528.18**	887.73**	238.88**	357.85**	552.73**	289.41**	331.53**
	F ₇	22.00	40.11**	387.69**	29.93**	562.87**	2450.25**	183.24**	389.40**	423.19**	335.08**	205.11**
Error	F ₆	18.00	0.66	2.71	0.08	3.11	6.82	6.90	5.93	3.28	2.85	20.35
	F ₇	18.00	1.20	1.82	0.05	2.41	4.66	7.88	7.45	2.04	4.55	11.49

Note: * Significant at 0.05 probability level. ** Significant at 0.01 probability level.

SV- Source of variation, df: Degrees of freedom, DFF: Days to first flowering

Table 2: Estimates of genetic variability parameters

Traits	Crosses	Mean	Range		Standardized Range	GCV (%)	PCV (%)	$h^2_{(bs)}$ %	GAM%
			Min.	Max.					
DFF	F ₆	30.25	29.00	32.00	0.10	02.31	02.57	89.93	10.53
	F ₇	30.44	26.00	35.00	0.30	04.23	05.63	75.23	21.76
Plant height (cm)	F ₆	37.83	34.28	41.33	0.19	04.65	05.03	92.50	05.44
	F ₇	36.08	12.40	49.36	01.02	19.09	21.83	87.49	25.86
Primary branches plant⁻¹	F ₆	04.73	03.20	05.55	0.50	12.77	15.11	84.56	07.65
	F ₇	04.86	03.12	07.56	0.91	13.46	15.26	88.23	17.75
SCMR	F ₆	38.02	32.00	43.66	0.31	08.42	10.69	78.83	08.55
	F ₇	42.98	35.00	55.36	0.47	06.12	06.99	87.63	19.74
SLA(cm²/g)	F ₆	153.47	132.00	180.20	0.31	06.65	9.26	71.86	13.70
	F ₇	122.50	98.00	156.00	0.47	08.18	09.08	90.10	16.86
Pods plant⁻¹	F ₆	28.30	17.50	44.16	0.94	19.29	24.52	78.68	29.64
	F ₇	32.58	18.00	49.00	0.95	17.81	25.20	70.69	35.14
Pod yield plant⁻¹(g)	F ₆	22.13	15.33	31.71	0.74	19.72	24.14	81.72	30.69
	F ₇	25.81	13.74	39.25	0.99	17.91	19.56	91.60	24.83
Kernel yield plant⁻¹(g)	F ₆	13.41	08.00	19.46	0.85	23.08	29.09	79.37	11.61
	F ₇	19.18	07.42	35.00	01.44	26.35	31.89	82.64	10.93
Shelling (%)	F ₆	58.82	50.00	65.29	0.26	7.01	08.97	78.21	12.61
	F ₇	52.46	40.98	60.22	0.37	06.46	09.16	70.55	13.31
SMK (%)	F ₆	58.49	8.55	82.00	01.26	24.65	26.87	91.76	50.79

	F ₇	48.09	31.77	60.11	0.59	17.66	18.06	97.82	36.40
--	----------------	-------	-------	-------	------	-------	-------	-------	-------

Note: GCV-Genotypic coefficient of variation, PCV-Phenotypic coefficient of variation, $h^2_{(bs)}\%$ –Heritability in broad sense, GAM%-Genetic advance as *per cent* of mean.

Table 3: Estimation of phenotypic correlation coefficients

Traits	Generations	Plant height (cm)	Primary branches plant ⁻¹	SCMR	SLA (cm ² /g)	Pods Plant ⁻¹	Pod yield plant ⁻¹ (g)	Kernel yield plant ⁻¹ (g)	Shelling (%)	SMK (%)
DFF	F ₆	0.48**	0.30**	0.32**	0.22*	0.32**	0.20*	0.41**	0.32**	0.30*
	F ₇	0.57**	0.36**	0.21*	0.17	0.26*	0.39**	0.39**	0.25*	0.43**
Plant height(cm)	F ₆	1.00	-0.20*	0.35*	0.32*	0.41*	0.46**	0.51**	-0.45**	0.48*
	F ₇	1.00	-0.32**	0.42	0.35*	0.32**	0.42**	0.45**	-0.37**	0.20*
Primary branches plant ⁻¹	F ₆		1.00	0.39*	-0.18	0.58**	0.32**	0.52**	0.32**	0.45**
	F ₇		1.00	0.25*	-0.20	0.46**	0.25*	0.65**	0.44**	0.36*
SCMR	F ₆			1.00	-0.67*	0.69**	0.65**	0.50*	-0.58**	0.48*
	F ₇			1.00	-0.5**	0.58**	0.52**	0.48*	-0.36**	0.29*
SLA(cm ² /g)	F ₆				1.00	-0.39*	-0.43**	-0.43**	-0.24*	0.18
	F ₇				1.00	-0.48**	-0.36**	-0.55**	-0.37**	0.07
Pods plant ⁻¹	F ₆					1.00	0.94**	0.62**	-0.45**	0.21
	F ₇					1.00	0.93**	0.87**	-0.35**	0.10
Pod yield plant ⁻¹ (g)	F ₆						1.00	0.76**	-0.42**	0.24*
	F ₇						1.00	0.92**	-0.55**	0.33**
Kernel yield plant ⁻¹ (g)	F ₆							1.00	0.41**	0.25*
	F ₇							1.00	0.35**	0.36**
Shelling (%)	F ₆								1.00	0.65**
	F ₇								1.00	0.49**

SMK (%)	F ₆									1.00
	F ₇									1.00

Note: *Significant at 0.05 probability level. ** Significant at 0.01 probability level

Table 4: Estimation of parent-offspring correlation, regression and heritability

Traits	Regression components			
	R	B	$h^2_{(bs)}$	$h^2_{(ns)}$
DFF	0.61*	0.44	75.23	38.08
Plant height (cm)	0.71**	0.56	87.49	48.31
Primary branches plant ⁻¹	0.25*	0.15	88.23	13.13
SCMR	0.38*	0.70	87.63	59.79
SLA(cm ² /g)	0.14	0.21	90.10	18.61
Pods plant ⁻¹	0.60**	0.56	70.69	47.61
Pod yield plant ⁻¹ (g)	0.53**	0.67	91.60	57.34
Kernel yield plant ⁻¹ (g)	0.27*	0.21	82.64	18.34
Shelling (%)	0.05	0.12	70.55	10.88
SMK (%)	0.28*	0.29	97.82	25.08

Note: r- Inter-generation correlation, b-regression, $h^2_{(bs)}$ - Heritability in broad sense, $h^2_{(ns)}$ - Heritability in narrow sense

Table 5: Performance of Selected superior families in F₇ generation for water use efficiency related traits, pod yield and its component traits

Sl. No	Name	SCMR	SLA(cm ² /g)	Pods plant ⁻¹	Pod yield plant ⁻¹ (g)	Kernel yield per plant (g)	Shelling%
1	C3-32-1-7-4-1	51.00	110.00	67.00	55.33	43.00	58.61
2	C3-130-5-6-2-4	48.00	114.00	54.00	49.00	31.40	58.97
3	C3-88-4-5-1-3	42.00	121.00	53.00	48.33	35.25	56.37
	GKVK-6	32.33	163.00	22.00	15.60	8.00	51.28
	KCG-2	38.10	150.00	25.00	19.10	15.33	49.60
	KCG-6	39.11	145.66	34.00	28.36	20.69	53.78

UNDER PEER REVIEW




Figure 1: Box plot showing variation for trait studied in the *Kharif* and summer seasons

UNDER PEER REVIEW



Figure 2 Selected superior plants for yield traits