

Original Research Article

Performance, haematology and serum biochemical status of two strains of broiler chicken fed *Ficus thonningii* leaf powder and vitamin C supplemented diets

ABSTRACT

Aims:To evaluate the effect of *Ficus thonningii* leaf powder (FTLP) and vitamin C supplemented diets on performance and haemato-biochemical status of two strains of broiler chickens.

Methodology:Four hundred and eighty-one-day-old of Arbor acre (AB) and Cobb 500 (CO) broiler chicks were randomized to eight dietary treatments; diet 1 and 2 (Control), diets 3 and 4 (200mg/kg vitamin C inclusion), diets 5 and 6 (1% inclusion of FTLP) and (diets 7 and 8 (200mg/kg + 1% FTLP inclusions) for AB and CO strains respectively.

Results:Result shows that feed intake, body weight gain and FCR were significant ($P<0.05$) at the starter phase. Interaction between strains and vitamin C were also significant ($P<0.05$) for body weight gain, FCR and feed intake. Vitamin C and FTLP supplementation significantly ($P<0.05$) improved the erythrocytes count while the MCV, MCH, WBC and lymphocytes of birds fed diets 2 and 6 were significantly ($P<0.05$) higher than the other diets. The dietary supplementation of vitamin C and FTLP shows that strain AB had a significantly ($P<0.05$) higher erythrocytes count than CO strain while the CO showed significant ($P<0.05$) improvement for the MCV and MCH. Vitamin C supplementation showed significant difference ($P<0.05$) for the RBC and the interaction of vitamin C and FTLP significantly ($P<0.05$) improved PCV, HB, MCV, MCH, WBC and lymphocytes. The interactive effect of vitamin C and FTLP significantly ($P<0.05$) improved the total protein and albumin. Cholesterol and creatinine levels were significantly ($P<0.05$) lowered by the inclusion of vitamin C and FTLP

Conclusion:As observed in this study, the inclusion of FTLP and vitamin C as a supplement in the diets of broiler chickens improved their performance, haematological indices and serum biochemical status of the tested birds.

Keywords: Ficus thonningii, supplementation, performance, haematological, serum biochemical

1. INTRODUCTION

The negative impacts of high ambient temperature on the performance and health status of broiler chicken raised outside their thermal neutral zone of $18^{\circ}\text{C} - 24^{\circ}\text{C}$, particularly in the tropics are on the increase [1]. This is due to climate problems, which have resulted in widespread heat stress among livestock species [2]. Studies had indicated that supplementation of antioxidants ameliorated the effects of heat stress on poultry; and there are calls for the adjustment of birds' diets to the climatic conditions of the regions where they are produced [3][4][2]. The inclusion of phyto-genic substances of plant origin in poultry diets has contributed to enhance the performance of animals, improving carcass traits and health status, ameliorating the negative impacts of oxidative stress and conferring positive effect on animal products [5], [6], [7]. The common wild fig, *Ficus thonningii* (odan) is one of the many fruit-bearing trees that have traditionally been used for treating diseases in Africa and

beyond Macerations of fresh *F. thonningii* leaves, taken orally, have been used traditionally for the treatment of diarrhea, gonorrhoea, and diabetes mellitus [8]. *F. thonningii* contains various biologically active compounds that exhibit physiological effects which could be responsible for its curative potentials for a wide range of diseases. These compounds known as phytochemicals are naturally produced by the tree as protection against biotic and abiotic stresses [9]. As a result of the antioxidant activities of *F. thonningii*, it has been reported that there is a protective effect on erythrocyte membrane from acetaminophen induced membrane peroxidation [10]. The antihaemolytic and haematinic potential is possibly due to its antagonistic action against the depletion of glutathione and hence prevention of the generation of free radicals which result in oxidative stress [10]. The inclusion of vitamin C in the diets of poultry birds has been reported to be a means of ameliorating the impact of heat stress due to its antioxidant properties, scavenging of free radicals [11], reduction of its biosynthesis during heat stress as well as poultry birds' ability to overcome infectious conditions [12]. Thus, two strains of broiler chickens were examined in this study for their performance, haematological and serum chemistry status by being fed diets supplemented with vitamin C and *Ficus thonningii* leaf powder.

2. MATERIAL AND METHODS

2.1 Ethical approval; collection, processing, and analysis of phytogens

The Research and Ethics Committee of the Department of Animal Production and Health, The Federal University of Technology, Akure, Nigeria, accepted the experiment's requirements and criteria for animal and animal protocol. [13]described the procedure for gathering and preparing *Ficus thonningii* leaf powder (FTLP) as well as investigated the proximate, phytochemical, antioxidant and mineral components of FTLP.

2.2 Experimental diets, experimental design and the birds' living environment

To meet the nutritional needs of the birds, a broiler chicken's basal diet was compounded for the starter phase (0 to 21 days) and finisher phase (21-42 days) using the requirement of [14]. For each phase, the basal diet was divided into eight equal portions and named diets 1 to 8. Diet 1 and 2 were not supplemented (negative control), while diets 3 and 4 had vitamin C (200mg/Kg of basal diet) supplementation. Then diets 5 and 6, received 1g FTLP/Kg of basal diet supplementation each while diets 7 and 8 had 1g FTLP/kg of basal diet + 200mg of vitamin C. The feeding trial was undertaken at the Federal University of Technology's Teaching and Research Farm in Akure, Nigeria.

In a completely randomized design, 240 one-day-old chicks each of Cobb 500 and Arbor Acre strains of broiler chicks weighing 37.40 ± 0.45 g were randomly assigned to all the eight experimental diets. The diets were replicated six times with ten birds (10 birds/replicate) and a total of 60 birds per treatment. Wood shavings were used as bedding for the floor of the experimental pen (2m x 1m) to a depth of 3cm. The temperature of the experimental house was kept at 31 ± 2 degrees Celsius for the first week and then reduced by 2 degrees Celsius each week after that until the temperature reached 26 ± 2 degrees Celsius. The lighting was turned on for 24 hours on the first day and 23 hours on consecutive days.

2.3 Performance characteristics

The body weight (BW) and feed intake (FI) of the tested birds were measured and recorded weekly. The average body weight gain (BWG) was calculated as the difference between the initial and final weights of the birds. The feed conversion ratio (FCR) was calculated by dividing the feed consumed by the weight gained.

2.4 Collection and analysis of Blood samples

On the last day of the feeding trial, 18 birds per replicate were randomly selected for blood collection. Syringe and needle were used to draw blood from the wing vein. For the determination of the serum biochemical indices (i. e. total protein, aspartate aminotransferase, creatinine, and cholesterol), 4ml of the blood sample was drawn into a plain bottle vial. The sample bottles were spun and the serum decanted into another plain bottle before freezing at -20°C . The remainder (2ml) was dispensed into Ethylenediaminetetraacetic acid bottle for haematological indices determination [15]. The serum biochemical concentrations were measured using a Reflectron @Plus 8C79 (Roche Diagnostic, GombH Mannheim, Germany) and kits.

2.5 Statistical data evaluation

The model: $T_{xy} = \mu + bx + \beta_{xy}$, was used in this experiment, where T_{xy} any of the factors of response; x = the overall average; bx = the xth treatment's effect (T = diets 1, 2, 3, 4, 5, 6, 7 and 8); and β_{xy} = random error due to the investigation. Using SPSS version 20, all of the data were subjected to one-way ANOVA. Duncan multiple range test of SPSS was used to detect the differences between the treatment means ($P < 0.05$).

3. RESULT

Table 2 shows the growth performance of different breeds of broiler chickens fed *Ficus thonningii* leaf powder (FTLP) and vitamin C supplemented diets. The body weight gain (BWG) at the starter phase (1 – 3 weeks) were not significantly different ($P > 0.05$) for birds across all the diets except diet 6 and the feed intake (FI) for birds fed diets 1, 2, 3 and 4 were significantly ($P < 0.05$) higher than the other diets. The FCR for birds fed diets 6 was significantly higher than the other tested diets. The breed and FTLP supplementation were significant ($P < 0.05$) for BWG, FI and FCR at the starter phase. The interaction between BRD x VC were significant ($P < 0.05$) for BWG and FCR; BRD x FTLP was significant ($P < 0.05$) for FCR while VC x FTLP showed significance ($P < 0.05$) for BWG and FI. The interactive effect of BRD x VC x FTLP were significant ($P < 0.05$) for BWG and FCR at the starter phase. At the finisher phase (4-6 weeks), the BWG were significantly higher ($P < 0.01$) in diets 2 and 4 though not significantly different ($P > 0.05$) from diet 1; FI was not significantly different ($P > 0.05$) except for diet 3 while the FCR was significantly higher ($P < 0.05$) for diet 5 though not significantly different ($P > 0.05$) from diet 6. The breed CO was significantly ($P < 0.05$) higher than the AB for the BWG while FTLP inclusion showed significance ($P < 0.05$) for BWG and FCR. The interaction between breed and vitamin C were significant ($P < 0.05$) for the FI and FCR; VC x FTLP showed significance ($P < 0.05$) for BWG while the interactive effect of **strain**, vitamin C and FCLP were significant ($P < 0.05$) for BWG and FI at the finisher phase. At the overall phase (1-6 weeks), the BWG was significantly ($P < 0.05$) higher for diets 2 and 4, though not significantly different from diet 1 while the FCR was significantly ($P < 0.05$) higher for diets 5 and 6 but not significantly different ($P > 0.05$) from diets 4, 7 and 8. The inclusion of FTLP at 1% showed significance ($P < 0.01$) for BWG and FCR. The interaction between breed and vitamin C was significant ($P < 0.05$) for FI; VC x FTLP was significant ($P < 0.05$) for FCR and the interaction between **strain**, vitamin C and FCLP was significant ($P < 0.01$) for FI.

Data on haematological indices of breeds of broiler chickens fed FTLP and vitamin C supplemented diets is presented in Table 3. The RBC of birds fed diet 7 was significantly ($P < 0.05$) higher, though not significantly different ($P > 0.05$) from birds fed diets 3 and 8. The MCV and MCH values of birds fed diets 2 and 6 were significantly ($P < 0.05$) higher than the

other diets while the WBC of birds fed diet 2 was significantly ($P<0.05$) higher, though not significantly ($P>0.05$) higher than birds fed diet 7. The lymphocytes of birds fed diets 1, 3, 4, 5 and 6 were lower significantly ($P>0.05$) than those fed diets 2 and 7. The AB strain showed higher significance ($P<0.05$) than the CO breed in the value of RBC while the MCV and MCH were significantly ($P<0.05$) higher in the CO breed. The inclusion of 200mg/kg of vitamin C showed significance ($P<0.05$) for the RBC while the inclusion was significantly ($P>0.05$) lower for the MCV and MCH. The interaction between BRD x VC were significant ($P<0.05$) for the PCV, HB, MCV, MCH, WBC and LYM. BRD x FTLP were significantly ($P<0.05$) higher for WBC and lymphocytes while there was no significant ($P>0.05$) interactive effect between BRD x VC x FTLP.

The result of serum chemistry indices of the two strain of broiler chickens fed diets supplemented with FTLP and Vitamin C is presented in Table 4. The different dietary inclusion had a positive significant ($P<0.05$) influence on the Aspartate aminotransferase (AST), Creatine (CREA), Cholesterol (CHOL), Globulin (GLO) and Total Protein (TP). The AST performance of diets 1 and 2 for the strain AB and CO were significantly ($P<0.05$) higher than those in diets 3 and 4 though, not significantly ($P>0.05$) different from diets 5 and 6. The serum creatinine level of the control diets (diets 1 and 2) for the strains AB and CO were significantly ($P<0.05$) different from diets 3, 4, 5, 6, 7 and 8. The serum cholesterol level of diets 1, 2 were significantly ($P<0.05$) higher than those of the other tested diets. The TP of diets 7 and 8 were significantly ($P<0.05$) higher than those of diets 1 and 2 but not significantly ($P>0.05$) different from diets 3, 4, 5 and 6. The GLB performance of diets 5 and 6 (FTLP) and diets 7 and 8 (vitamin C + FTLP) were not significantly ($P>0.05$) different but they were significantly ($P<0.05$) higher than those of diets 1, 2, 3 and 4, Vitamin C inclusion also had significance ($P<0.05$) on the AST, CREA, CHOL, TP and ALB. The inclusion of FTLP has significant effect ($P<0.06$) on the serum creatinine, serum cholesterol, total protein and serum globulin. The interactive effect of vitamin C and FTLP on AST, CHOL and CREA levels of the birds were significant ($P<0.05$).

Table 1: Composition of the experimental diets

Ingredients (g/kg)	Starter feed (%)	Finisher diet (%)
Maize	52.35	59.35
Rice bran	7	0
Maize bran	0	6
Soybean meal	30	24
Soy oil	3	3
Fish meal	3	3
Limestone	0.5	0.5
Bone meal	3	3
Salt	0.3	0.3
Premix	0.3	0.3
Methionine	0.3	0.3
Lysine	0.25	0.25
<u>Nutrient composition (g/kg)</u>	100	100
*Crude protein	22.18	20.03
Metabolizable energy (Kcal/kg)	3018	3108.1

Table 2 The growth performance response of different strains of broiler chickens to *Ficus thonningii*, leaf powder and vitamin C dietary supplementations

Diet	STRN	VC mg/kg	FTLP %	IW g/b	BWG 1-3 wks	FI 1-3 wks	FCR 1-3 wks	BWG 4-6 wks	FI 4-6 wks	FCR 4-6 wks	BWG 1-6 wks	FI 1-6 wks	FCR 1-6 wks
1	AB	0	0	37.66	845.47 ^a	1022.48 ^a	1.21 ^{bc}	1777.11 ^{ab}	3262.23 ^{bc}	1.8 ^{bc}	2622.58 ^{ab}	4284.71	1.62 ^{ab}
2	CO	0	0	37.57	792.45 ^a	982.48 ^a	1.24 ^b	1883.76 ^a	2752.24 ^{bc}	1.4 ^c	2676.21 ^a	3734.71	1.40 ^b
3	AB	200	0	37.34	812.71 ^a	986.61 ^a	1.21 ^{bc}	1600.96 ^{bc}	2564.39 ^c	1.6 ^{bc}	2413.67 ^{bc}	3551.00	1.47 ^b
4	CO	200	0	37.56	807.12 ^a	1002.41 ^a	1.24 ^b	1911.59 ^a	3529.52 ^a	1.8 ^{bc}	2718.71 ^a	4531.93	1.67 ^{ab}
5	AB	0	1	37.61	820.06 ^a	856.95 ^c	1.04 ^e	1328.55 ^d	3193.86 ^{abc}	2.4 ^a	2148.61 ^d	4050.81	1.90 ^a
6	CO	0	1	37.52	648.59 ^b	876.55 ^{bc}	1.35 ^a	1543.93 ^{bcd}	3138.92 ^{abc}	2.0 ^{ab}	2192.52 ^{cd}	4015.47	1.84 ^a
7	AB	200	1	37.65	787.91 ^a	899.22 ^{bc}	1.14 ^{cd}	1624.30 ^{bc}	3071.24 ^{abc}	1.8 ^{bc}	2412.21 ^{bc}	3970.47	1.64 ^{ab}
8	CO	200	1	37.42	808.01 ^a	904.24 ^b	1.11 ^{de}	1533.00 ^{cd}	2921.14 ^{abc}	1.9 ^{bc}	2341.01 ^{cd}	3825.38	1.63 ^{ab}
SEM				0.04	13.67	13.07	0.01	44.04	81.28	0.07	47.79	83.93	0.04
P-value				0.60	0.01	0.01	0.01	0.01	0.05	0.01	0.01	0.06	0.02
AB				37.56	816.54 ^a	941.32	1.15 ^d	1582.73 ^d	3022.93	1.93	2399.27	3964.25	1.66
CO				37.52	764.04 ^b	941.42	1.23 ^a	1718.07 ^a	3085.45	1.82	2482.11	4026.87	1.63
SEM				0.06	12.06	6.79	0.01	36.73	94.43	0.07	38.75	98.92	0.04
P-value				0.60	0.01	0.99	0.01	0.01	0.64	0.25	0.15	0.66	0.69
0				37.63	720.52	929.51	1.12	1552.83	3228.04	1.76	2385.60	4167.66	1.62
200				37.49	807.56	953.32	1.17	1612.63	2817.81	1.87	2412.94	3760.73	1.65
SEM				0.08	17.06	9.61	0.01	51.94	133.54	0.09	54.81	139.89	0.06
P value				0.28	0.12	0.17	0.07	0.52	0.63	0.17	0.27	0.71	0.19
0				37.50	829.09 ^a	1004.54 ^a	1.21 ^a	1689.03 ^a	2913.31	1.71 ^b	2518.13 ^a	3917.85	1.55 ^b
1				37.63	803.99 ^b	878.09 ^b	1.09 ^b	1476.42 ^b	3132.55	2.16 ^a	2280.41 ^b	4010.64	1.77 ^a
SEM				0.08	17.06	9.61	0.01	51.94	133.54	0.09	54.81	139.89	0.06
P value				0.82	0.01	0.01	0.01	0.01	0.69	0.01	0.01	0.67	0.01
Interactions P-value													
STRN x VC				0.63	0.01	0.29	0.01	0.62	0.02	0.02	0.54	0.02	0.08
STRN x FTLP				0.24	0.19	0.22	0.01	0.17	0.23	0.53	0.09	0.29	0.86
VC x FTLP				0.44	0.04	0.04	0.06	0.05	0.44	0.48	0.01	0.55	0.03
STRN x VC x FTLP				0.21	0.05	0.08	0.01	0.02	0.01	0.62	0.11	0.01	0.16

Means with a different superscript in the same column are significantly (P<0.05) different; STRN: Strains; AB: Arbor acre; CO: Cobb 500; VC: Vitamin C; FTLP: *Ficus Thonningii*; IW: Initial weight; BWG: Body weight gain; FI: Feed intake; FCR: Feed conversion ratio; SEM: Standard error of the means.

Table 3 The haematological indices response of different strains of broiler chickens to *Ficus thonningii* leaf powder and vitamin C dietary supplementations

Diet	STRN	VC mg/kg	FTLP %	PCV %	RBC x10 ⁶ /l	HB g/dl	MCHC g/dl	MCV fl	MCH Pg/cell	WBC x10 ⁹ /l	GRA x10 ⁹ /l	LYM x10 ⁹ /l	MON x10 ⁹ /l
1	AB	0	0	30.50	2.26 ^c	10.16	33.56	135.90 ^b	45.30 ^b	2.86 ^{bc}	0.80	1.96 ^{bc}	0.06
2	CO	0	0	32.00	1.87 ^c	10.66	33.30	173.26 ^a	57.76 ^a	4.06 ^a	1.00	3.00 ^a	0.07
3	AB	200	0	31.5	3.26 ^{ab}	10.50	33.31	97.46 ^c	32.46 ^c	2.86 ^{bc}	0.80	2.10 ^{bc}	0.00
4	CO	200	0	30.00	2.86 ^b	10.00	33.30	105.33 ^c	35.10 ^c	2.66 ^{bc}	0.90	1.70 ^{bc}	0.06
5	AB	0	1	30.66	2.17 ^c	10.23	33.00	142.50 ^b	47.50 ^b	3.10 ^{abc}	1.00	2.10 ^{bc}	0.00
6	CO	0	1	30.67	2.07 ^c	11.23	33.13	164.50 ^a	54.83 ^a	2.90 ^{bc}	0.90	1.93 ^{bc}	0.06
7	AB	200	1	34.00	3.73 ^a	11.32	33.33	92.63 ^c	30.86 ^c	3.53 ^{ab}	0.90	2.56 ^{ab}	0.07
8	CO	200	1	29.33	3.20 ^{ab}	9.77	33.06	91.80 ^c	30.60 ^c	2.23 ^c	0.80	1.43 ^c	0.00
SEM				0.51	0.14	0.17	0.06	6.78	2.62	0.14	0.04	0.12	0.01
P-value				0.196	0.01	0.17	0.43	0.01	0.01	0.03	0.92	0.02	0.17
A				31.67	2.85 ^a	10.55	33.30	117.13 ^b	39.03 ^b	3.09	0.87	2.18	0.33
R				31.25	2.50 ^b	10.41	33.20	133.72 ^a	44.57 ^a	2.96	0.90	2.01	0.05
SEM				0.65	0.09	0.22	0.08	3.65	1.22	0.16	0.07	0.14	0.01
P-value				0.66	0.02	0.64	0.43	0.01	0.01	0.59	0.81	0.40	0.38
0				31.71	2.09 ^b	10.57	33.25	154.04 ^a	51.35 ^a	3.23	0.92	2.25	0.05
200				31.21	3.26 ^a	10.40	33.25	96.81 ^b	32.25 ^b	2.82	0.85	1.95	0.03
SEM				0.65	0.09	0.22	0.08	3.65	1.22	0.16	0.07	0.14	0.01
P value				0.59	0.01	1.00	1.00	0.01	0.01	0.09	0.46	0.14	0.38
0				31.00	2.56	10.33	33.36	127.99	42.65	3.12	0.87	2.19	0.05
1				31.91	2.79	10.64	33.13	122.85	40.95	2.94	0.90	2.01	0.03
SEM				0.65	0.09	0.22	0.08	3.65	1.22	0.16	0.07	0.13	0.01
P value				0.33	0.11	0.32	0.07	0.33	0.34	0.45	0.81	0.36	0.38
Interactions P-value													
STRN x VC				0.01	0.43	0.01	0.79	0.02	0.02	0.01	0.81	0.01	0.38
STRN x FTLP				0.66	0.76	0.64	0.79	0.26	0.26	0.02	0.23	0.02	0.38
VC x FTLP				1.00	0.21	0.97	0.29	0.45	0.44	0.22	0.81	0.16	0.38
STRN x VC x FTLP				0.22	0.43	0.21	0.19	0.75	0.75	0.74	0.80	0.55	0.06

Means with a different superscript in the same column are significantly (P<0.05) different; STRN.: Strains; AB: Abore acre; CO: Cobb 500; VC: Vitamin C; FTLP: *Ficus Thonningii* leaf powder; PCV: Packed cell volume; RBC: Red blood cells; HB: Haemoglobin conc.; MCHC: Mean cell haemoglobin concentration; MCV: Mean cell volume; MCH: Mean cell haemoglobin; WBC: White blood cells; GRA: Granulocytes; LYM: Lymphocytes; MON: Monocytes; SEM: Standard error of the means.

Table 4: Serum biochemical profiles of different strains of broiler chicken fed *Ficus thonningii* leaf powder and vitamin c supplemented diets

Diet	STRN	VC mg/kg	FTLP %	AST (IU/L)	ALT (IU/L)	CREA (μ mol/L)	CHOL (mmol/L)	TP (g/L)	ALB (g/L)	GLB (g/L)	ALB/ GLO
1	AB	0	0	168.06 ^a	18.37	79.73 ^a	157.95 ^a	52.66 ^b	23.97	28.68 ^b	0.83 ^b
2	CO	0	0	171.28 ^a	17.54	81.20 ^a	155.53 ^a	53.65 ^b	23.88	29.77 ^b	0.80 ^b
3	AB	200	0	132.98 ^{bc}	16.55	65.18 ^b	134.96 ^b	63.47 ^{ab}	32.51	30.95 ^b	1.05 ^{ab}
4	CO	200	0	128.61 ^c	17.63	65.31 ^b	139.11 ^b	64.41 ^{ab}	35.17	29.24 ^b	1.19 ^a
5	AB	0	1	152.41 ^{ab}	16.28	67.75 ^b	129.78 ^b	61.93 ^{ab}	27.00	34.93 ^a	0.78 ^b
6	CO	0	1	152.39 ^{ab}	17.24	67.81 ^b	117.43 ^c	58.52 ^{ab}	25.30	33.22 ^a	0.75 ^b
7	AB	200	1	135.07 ^{bc}	16.91	63.93 ^b	119.09 ^c	66.79 ^a	32.35	34.44 ^a	0.94 ^{ab}
8	CO	200	1	132.22 ^{bc}	17.19	63.84 ^b	114.50 ^c	68.84 ^a	35.57	33.36 ^a	1.06 ^{ab}
SEM				3.77	0.57	1.62	3.42	1.56	1.39	0.52	0.04
<i>P</i> -value				0.01	0.99	0.01	0.01	0.05	0.10	0.01	0.05
	AB			147.13	17.02	69.15	135.44	61.21	28.96	32.25	0.90
	CO			146.12	17.40	69.54	131.64	61.35	29.95	31.40	0.95
	SEM			3.16	0.95	1.46	1.77	1.80	1.70	0.37	0.05
	<i>P</i> -value			0.82	0.78	0.85	0.14	0.95	0.68	0.13	0.46
	0			161.03	17.36	74.12 ^a	140.17 ^a	56.69 ^b	25.03 ^b	31.65	0.79 ^b
	200			132.22	17.07	64.57 ^b	126.91 ^b	65.87 ^a	33.88 ^a	31.99	1.06 ^a
	SEM			3.16	0.95	1.46	1.77	1.80	1.70	0.37	0.05
	<i>P</i> value			0.01	0.83	0.01	0.01	0.01	0.01	0.52	0.01
	0			150.23	17.52	72.86 ^a	146.89 ^a	58.55 ^b	28.88	29.66 ^b	0.97
	1			143.02	16.90	65.83 ^b	120.20 ^b	64.02 ^a	30.03	33.99 ^a	0.88
	SEM			3.16	0.95	1.46	1.77	1.80	1.70	0.37	0.05
	<i>P</i> value			0.12	0.65	0.01	0.01	0.04	0.64	0.01	0.24
	Interactions <i>P</i> -value										
	STRN x VC			0.56	0.82	0.85	0.17	0.60	0.44	0.32	0.25
	STRN x FTLF			0.92	0.85	0.84	0.08	0.75	0.90	0.33	0.94
	VC x FTLF			0.03	0.67	0.01	0.02	0.54	0.66	0.34	0.62
	STRN x VC x FTLF			0.79	0.63	0.88	0.90	0.59	0.83	0.13	0.94

Means with a different superscript in the same column are significantly ($P < 0.05$) different; STRN: Strains; AB: Arbor acre; CO: Cobb 500; VC: Vitamin C; FTLF: *Ficus thonningii* leaf powder; AST: Aspartate aminotransferase; ALT: Alanine aminotransferase; CREA: Creatine; CHOL: Cholesterol; TP: Total Protein; ALB: Albumin; GLB: Globulin; SEM: Standard error of the means.

4. Discussion

[16] reported that feed ingredients which are rich in antioxidant and polyphenols which are naturally abundant in them are getting attention, and [17] as well as [18] agreed that they could be used to boost performance in heat-stressed birds. The inclusion of phytogetic substances in feed have shown positive effects on the performance and biological health of broiler chickens [19]. Supplementation of broiler feed with plant extract containing secondary metabolites has been shown to promote growth performance and immune response in broiler chickens [20]. However, result of the present study does not indicate significant improvement in feed intake and body weight gain due to vitamin C and FTLP supplementation. The lower BWG in the experimental birds due to FTLP supplementation in this study suggests a possible compromise of nutritive quality and impairment of feed acceptability by the birds, because some phytochemicals may interfere with nutrient bioavailability for the birds, especially when present in excessive amounts in their diets and feed physical properties [21][5]. [22] reported that the usage of feed supplements can affect an animal's willingness to eat a given feed. The reduction in the FI recorded in the FTLP treated groups (5, 6, 7 and 8) and the significant effect of FTLP supplementation in this study suggest the negative effects of the FTLP on the feed acceptability and consumption by the birds during the early state of their life (the starter phase). The reduced FI in the affected treatment groups could be due to the phyto-constituents of the FTLP. For instance, tannin, when present in high levels in diets reduces the voluntary feed intake and nutrient digestibility [23]. The feed conversion ratio (FCR) value ranges from 1.40 – 2.4 and 1.40 – 1.90 for the finisher and overall phases respectively, which falls within the recommended value for broiler birds. The values obtained here were within the optimum value of 2.0 for broilers by [24]. This shows that the inclusion of *F. thonningii* in the diet of experimental birds have effect on the on the overall performance of the birds raised in an enclosed house.

Haematological markers are good indicators of an animal's physiological status, and changes in them are useful in determining how the animal responds to various physiological settings [25]. The protective properties of phytochemicals from plants against reactive oxygen species, as well as their antibacterial action are regarded to be one of their health-promoting impacts and the concentration of phenolic compounds and several vitamins in these medicinal plants may be related to their potential [26]. The PCV aids in determining whether supplemented diets have a negative impact on the health status of the broiler chickens. A high PCV value indicates the presence of toxic factors that may have an adverse effect on blood formation [27]. The values of PCV recorded in this study were within the range recorded by [28], who reported a range of 28 – 35% for broiler chicken aged 5 to 7 weeks. The significant effect of the supplementation of vitamin C and FTLP on the tested birds could be an indication that the interactive effect of vitamin C and FTLP combined could lead to increased feed intake which could lead to increase in the number of RBC. This could be attributed to the increase in feed intake at the finisher phase in the present study. Accumulation of free radicals has been reported to promote oxidative stress which is linked to many chronic diseases and reduced performance of animals [29]. [30] reported that supplementation of diets with vitamin C can increase blood antioxidant levels by up to 30%, thus helping the body's natural defenses against inflammation. The vitamin C and Zinc present in *Ficus thonningii* as reported by [13] could be attributed to the performance of the birds in this regard. This is in agreement with the work of [31] who reported that the interaction of Zinc and ascorbic acid as supplements enhances production performance and immune status.

[26] has reported on the significance of the liver as the coordinating point for different metabolic, productive and digestive functions and it is prone to different levels of biological and chemical damages depending on the amount of specific serum enzymes which has their

originating sources from the liver. These enzymes, depending on their levels may compromise many body functions, thereby resulting in poor immunity status and weak production performance. Aspartate aminotransferase (AST), alkaline phosphatase (ALP), and alanine aminotransferase (ALT) in the blood are key bioindicators of the extent of the function and damage in the liver [32]. A surge in the levels of these enzymes is usually associated with liver or muscle damage, resulting from the body's reaction to stress [33]. The reduction in AST due to inclusion of Vitamin C and FTLP can be deduced as an indication of better liver function due to the activities of the antioxidants and ascorbic acid which reduces oxidative stress induced by hepatotoxicity [34]. **Creatinine** as a nitrogenous organic acid assist in the supply of energy to the cells, it is a by-product of muscle metabolism and it is excreted entirely by the kidney. Increased level of creatinine is an indication of decreased kidney function [35] and indication of a chronic kidney disease [36]. Inclusion of vitamin C and FTLP in this study showed reduced level of creatinine less than the value reported by [27], who reported 79.00 and 81.00% for AB and CO breeds respectively. Cholesterol, produced in the liver is a product of fat metabolism. Increase in the level of cholesterol is associated with hormonal and metabolic diseases, liver disease as well as kidney malfunctioning. [35]. The significantly lower values of cholesterol with the inclusion of FTLP in comparison to the control diets affirmed the believe that chickens fed with herbs have low levels of cholesterol as a result of decreased activity of lipogenic enzymes in the liver [37]. Serum proteins are essentially synthesized in the liver and they perform the function of maintaining blood volume through colloidal osmotic effect, regulate blood pH, transport hormones and drugs, participate in cell coagulation, catalyze chemical reactions, regulate the metabolism and participate in the body defense against foreign objects [38]. The main functions of albumin are the transport of several molecules and the maintenance of blood oncotic pressure [38]. The inclusion of FTLP in diets of tested birds shows a significant effect on the albumin level which could be attributed to the vitamin C content of *Ficus thonningii* as reported by [13]. This agrees with the work of [39] who reported that the inclusion of vitamin C in diets ameliorate the pathological changes in serum protein and body weight.

5. Conclusion

The result of this study revealed the potentials of *Ficus thonningii* leaf powder to enhance performance and ameliorate the effect of heat stress of broiler chickens when used as organic supplement in feed. As observed in this study, vitamin C and FTLP supplementation increased feed conversion ratio as well as improving the haematological profiles of the tested bird. The serum cholesterol and creatinine were reduced as a result of the inclusion of vitamin C and FTLP, suggesting that the supplementation improved the functioning of the liver and kidney of the tested birds, thus mitigating the damaging effects of heat and oxidative stress.

Compliance with ethical standard

The study was carried out with the approval of the committee on ethics for care and use of animal for research of The Federal University of Technology, Akure, Nigeria.

REFERENCES

1. Olanrewaju HA., Purswell JL., Collier SD and Branto SL. "Effect of ambient temperature and light intensity on physiological reactions of heavy broiler chickens". Poultry Science, 2010. 89: 2668– 2677, ISSN: 1525-3171. <https://doi.org/10.3382/ps.2010-00806>.

2. Kpomasse CC, Oke OE, Houndonougbo FM and Kokou Tona. Broiler production challenges in the tropics: A review. *Veterinary Medicine and Science*. 2021. DOI: 10.1002/vms3.435.
3. Suganya, T., Senthilkumar, S., Deepa, K. and Amutha, R. "Nutritional management to alleviate heat stress in broilers". *International Journal of Science, Environment and Technology*, 2015. 4(3): 661– 666, ISSN: 2278-3687.
4. Adeyeye SA., Oloruntola OD., Ayodele, SO., Falowo AB and Agbede JO. "Wild sunflower and goat weed compositemix supplementation in broiler chicken: effects on performance, health status and meat". *Acta Fytotechnica et Zootechnica*, 2020. 23(4): 205-212, ISSN: 1336-9245. <https://doi.org/10.15414/afz.2020.23.04.205-212>.
5. Oloruntola OD, Adu OA, Gbore FA, Falowo AB, Olarotimi OJ. Performance of broiler chicken fed diets supplemented with *Irvingiagabonensis* kernel powder and *Ocimumgratissimum* leaf powder. *Slovakia Journal of Animal Science*. 2021. 54(1):7-20.
6. Gbore FA, Oloruntola OD, Adu OA, Olarotimi OJ, Falowo AB and Afolayan EO. Serum and meat antioxidative status of broiler chickens fed diets supplemented with garlic rhizome meal, moringa leaf meal and their composite. *Tropical Animal Health and Production*. 2021; 53:26. Available: <https://doi.org/10.1007/s11250-020-02438-9>.
7. Adu OA, Gbore FA, Oloruntola OD, Falowo AB and Olarotimi OJ. The effect of *Myristica fragrans* seed meal and *Syzygiumaromaticum* leaf meal dietary supplementation on growth performance and oxidative status of broiler chicken. *Bulletin of National Research Centre*. 2020; 44:149. Available: <https://doi.org/10.1186/s42269-020-00396-8>.
8. Njoronge GN and Kibunga JW. Herbal medicine acceptance, sources and utilization for diarrhea management in cosmopolitan urban area (Thika, Kenya). *African Journal of Ecology*. 2007. 45:65-70.
9. Dangarembizi R, Erlwanger KH, Moyo D. and Chivandi E. Phytochemistry, Pharmacology and Ethnomedicinal uses of *Ficus thonningii* (Blume moraceae): A Review. *African Journal of Traditional Complement Alternative Medicine*. 2013. 10(2): 203-212.
10. Ahur VM, Madubunyi I., Adekonla AY and Udem SC. The effects of acetyl acetate extract of *Ficus thonningii* (Blume) leaves on erythrocyte osmotic fragility and haematological parameters in acetaminophen-treated rats. *Comparative Clinical Pathology*. 2010. 10:1107-1111.
11. Roussan DA., Kwaldeh GY., Haddad RR., Shaheen IA., Salameh GR and Rifai AI. Effect of ascorbic acid, acetylsalicylic acid, sodium bicarbonate, and potassium chloride supplementation in water on the performance of broiler chickens exposed to heat stress. *Journal of Applied Poultry Research*. 17: 141-144.
12. Balvave D. Challenges of accurately defining the nutrient requirements of heat-stressed poultry. In: *World's Poultry Science Association Invited Lecture*. *Poultry Science*. 2004. 83: 5-14.
13. Osowe CO, Olowu OPA, Adu OA, Oloruntola OD and Chineke CA. Proximate and mineral composition, phytochemical analysis, and antioxidant activity of fig trees (*Ficus* spp.) leaf powder. *Asian J. Bioch., Gen. Mol. Bio*. 2021. 9(1): 19-29.
14. NRC. *Nutrient requirements of poultry (9th Revised Ed.)*, 1994. Washington, DC: National Academy Press.
15. Cheesbrough M. *District Laboratory Practice in Tropical Countries*(1st ed.), (2000). Cambridge University Press, UK.
16. Crozier A, Jaganath IB and Clifford MN. Dietary phenolics: Chemistry, bioavailability and effects on health. *Natural Products Reports*, 2009. 26(8); 1001–1043. <https://doi.org/10.1039/b802662a>

17. Hu R, He Y, Arowolo MA, Wu S and He J. Polyphenols as potential attenuators of heat stress in poultry production. *Antioxidants (Basel)*, 2019. 8(3); 67. <https://doi.org/10.3390/antiox8030067>
18. Oke OE, Oyetunji DE, and Daramola JA. Responses of broiler chickens under hot humid tropical climate as affected by *Azadirachta indica* leaf. *Bulletin of Animal Health and Production in Africa*. 2017. 65; 321–329.
19. Abudabos AM, Alyemni AH, Dafalla YM and Khan RU. The effect of phytogenic feed additives to substitute in-feed antibiotics on growth traits and blood biochemical parameters in broiler chicks challenged with *Salmonella typhimurium*. *Environ. Sci Pollut. Res. Int.* 2016. 23; 24151–24157. [CrossRef] [PubMed]
20. Dhama K., Tiwari R., Khan RU., Chakraborty S., Gopi M., Karthik K., Saminathan M., Desingu PA and Sunkara LT. "Growth promoters and novel feed additives improving poultry production and health, bioactive principles and beneficial applications: The trends and advances- A Review". *International Journal of Pharmacology*, 2014. 10(3): 129-159, ISSN: 1812-5700. <https://doi.org/10.3923/ijp.2014.129.159>.
21. Valenzuela-Grijalva NV., Pinelli-Saavedra A., Muhlia-Almazan A., Dominguez-Diaz D. and Gonzalez-Rios H. Dietary inclusion effects of phytochemicals as growth promoters in animal production. *Journal of Animal Science and Technology*. 2017. 59:8.
22. Hassan ZM., Manyelo TR., Salaledi L., Mabelebele M. The effects of tannins in monogastric animals with special reference to alternative feed ingredients. *Molecules*. 2020. 25(20): 4680. <https://doi.org/10.3390/molecules25204680>.
23. Prabakaran R. Good practices in planning and management of integrated commercial poultry production in south Asia. *FAO Animal Production and Health Paper*. 2003. 159:12-24.
24. Attia YA, Bakhawain AA and Bertu NK. Utilization of thyme powder (*Thyme vulgaris* L.) as a growth promoter alternative to antibiotics for broiler chickens raised in a hot climate. *Eur. Poult. Sci.* 2018. 18; 82.
25. Khan TA and Zafar F. Haematological study in response to varying doses of estrogen in broiler chicken. *International Journal of Poultry Science*. 2005. 4(10); 748-751
26. Aikpitanyi KU and Egweh NO. Haematological and biochemical profile of broiler chickens fed diets containing ginger and black pepper additives. *Nigerian J. Anim. Sci.* 2020 Vol 22 (2):114-125 (ISSN:1119-4308)
27. Iyaode II, Ibrahim HO, Uwade F and Shittu MW. Haematology and serum biochemistry of broiler strains (Cobbs and Arbor-acre) fed ginger (*Zingiber officinale*). *GSC Biol. Pharm. Sci.* 2020. 11(02); 320–326. DOI: <https://doi.org/10.30574/gscbps.2020.11.2.0145>
28. Onyishi GC, Oguine CC, Nwani SI, Aguzie IO and Nwani CD. Haematological parameters dynamics of developing *Gallus Gallus domesticus*. *Ani. Res. Int.* 2017. 14(2); 2769 – 2776
29. Pham-Huy LA, He H and Pham-Huy C. Free radicals, antioxidants in disease and health. *Int. Journal of Biomed Sci.* 2008. 4(2):89-96. PMID: 23675073; PMCID: PMC36146967
30. Kim MK, Sasazuki S, Sasaki S, Okubo S, Hayashi M and Tsugane S. Effect of five-year supplementation of vitamin C on serum vitamin C concentration and consumption of vegetables and fruits in middle aged Japanese: a randomized control trial. *J Am Coll Nutr.* 2003. 22(3); 208-216. Doi:10.1080/07315724.2003.10719295. PMID: 12805247.
31. Chand N, Naz S, Khan A, Khan S and Khan RU. Performance traits and immune response of broiler chicks treated with zinc and ascorbic acid supplementation

- during cyclic heat stress. *International Journal of Biometeorology*, 2014. 58(10); 2153–2157. <https://doi.org/10.1007/s00484-014-0815-7>
32. Yildirim EI, Yalchinkaya M, Kanbur MÇ and Oruc E. Effects of yeast lucomannan on performance, some biochemical parameters and pathological changes in experimental aflatoxicosis in broiler chickens. *Révue de MédecineVétérinaire*, 2011. 162: 413–420
 33. Lumeij JT. Avian clinical biochemistry. Pages 839–872 in *Clinical Biochemistry of Domestic Animals*. 6th ed. 2008. In: J. J. Kaneko, J. W. Harvey, and M. L. Bruss, ed. Academic Press, Burlington, MA.
 34. Nayila I. Effect of Ascorbic acid supplementation on liver function tests in hepatitis C patients. *Open Journal of Internal Medicine*. 2020. 10; 263-279. Doi: 10.4236/ojim.2020.103028.
 35. Krista W, Kristiina R and Margo ST. Serum Biochemistry. 2021. VCA Hospitals
 36. Newman T. Symptoms, causes, and treatment of chronic kidney disease. 2021. www.medicalnewstoday.com
 37. Ciftci M, Simsek T, Tilmaz A and Bestami O. Effects of dietary antibiotics and cinnamon oil supplementation on antioxidant enzyme activities, cholesterol levels and fatty acid compositions of serum and meat in broiler chickens. *Acta Veterinaria Scandinavica*.2010. 79:33-40.
 38. Melillo A. Applications of serum protein electrophoresis in exotic pet medicine. *Veterinary Clinics of North America Exotic Animal Practice*. 2013. 16(1): 211-225.
 39. Edoga CO, Njoku OO, Okeke JJ and Ani CE. Effects of vitamin C on serum protein, albumin, beta-globulin profiles and body weight of *Trypanosoma Brucei* infected *Rattus noregicus*. *Animal Research International*. 2013. 10(1): 1685-168

UNDER PEER REVIEW

