

Effect of *Carica Papaya* leaves and seeds on Electrolyte and Hematological Parameters in Albino Wistar Rats Exposed to Lead Nitrate

Abstract

Background: Exposure to lead produces a variety of adverse health effects. This study evaluated the effect of aqueous extracts of *Carica papaya* leaves and seeds on electrolyte and hematological parameters in albino Wistar rats exposed to lead nitrate. **Methods:** Thirty male albino Wistar rats weighing between 174- 196g were assigned to six groups of five animals. Group 1 served as control, group 2 received 50mg/kg body weight lead nitrate (PbNO_3), groups 3 and 4 received 500 mg/kg body weight aqueous extract of *Carica papaya* leaves (CPL) and *Carica papaya* seed (CPS) respectively, groups 5 and 6 received 50mg/kg body weight PbNO_3 + 500mg/kg body weight CPL and CPS respectively for 31 days by oral gavaging. Animals were sacrificed and blood collected for various assay using standard methods. ANOVA on SPSS version 25 was used to determine variations within and between groups. **Results:** Increases in sodium and potassium, and decreases in chloride, calcium and bicarbonate levels were observed in the lead intoxicated group. Treatment with CPL and CPS reversed the alterations in electrolytes. Levels of RBC, PCV, Hb, MCH, MCHC and lymphocytes were lower in the lead intoxicated group relative to the lead intoxicated groups complemented with CPL and CPS, while WBC, MCV and platelet counts were higher in the lead intoxicated group when compared to the group complemented with CPL and CPS. **Conclusion:** This finding suggests that CPL and CPS could be protective against lead induced alterations in electrolytes and hematological parameters and could be harnessed for these potentials.

Keywords: *Carica papaya* leaf and seed; lead nitrate; electrolyte; hematology

Introduction

Heavy metals are environmental pollutants owing to their toxicity, longevity in the atmosphere, and ability to bio-accumulate in the human body. Most heavy metals occur naturally, but their increased atmospheric and environmental levels are as a result of anthropogenic activities [1]. Heavy metals have harmful effects on human health, and exposure to these metals has been increased by industrial and Contamination of water, air and even food by toxic metals is a public health concern [2]. Heavy metals are characterized by their high atomic mass and toxicity to living organisms. Most heavy metals may be lethal to humans [1]. Exposure to heavy metals can occur through ingestion, absorption or inhalation. Exposure to heavy metals result in bioaccumulation in biological systems [3]. Metals may frequently react with biological systems by losing one or more electrons and forming metal cations which have affinity to the nucleophilic sites of vital macromolecules. Simultaneous exposure to two or more metals may have cumulative effects [4,5,6,7]. Low level exposure to lead has been linked with functional and structural impairments in both human and experimental animals [8]. The hematopoietic, nervous and renal tissues are the main targets of lead. Moreover, it hinders the efficiency of the hepatic, reproductive and immune function [9,10]. Approximately 90% of absorbed lead is reported to be stored in the bone with a half-life of 600 - 3000 days. The remaining 10% is stored in soft tissues

like kidney, liver and brain. The half-life of lead in these tissues ranges from 40 - 50 days [11]. Lead is conjugated in the liver and passed to the kidney, where a small quantity is excreted in urine and the rest accumulates in various body organs, affecting many biological activities at the molecular, cellular and intercellular levels, which may result in morphological alterations that can remain even after lead levels have fallen [13,14,15].

Lead toxicity is closely related to its accumulation in various tissues and its interference with the electrolytes that hamper several physiological processes [16]. Electrolytes are structural components of body soft tissues like liver, muscle and also participate in acid-base balance and are components of blood buffers. Electrolytes play significant role in several body processes, such as controlling fluid levels, acid-base balance (pH) nerve conduction, coagulation and muscle contraction [12,17]. Fluid and electrolytes homeostasis is usually maintained within narrow limits and therefore, it must be kept at a level that is suitable for normal biochemical and physiological functions [18]. Therefore, imbalance in serum electrolytes is an important indicator of toxicity in identifying target organs and general health status of animals. Lead is known to substitute for zinc in a number of enzymes in heme biosynthetic pathway and pyrimidine, 5-nucleotidase, important for the correct metabolism of deoxyribonucleic acid (DNA) and can therefore cause fetal damage [19].

About 99% of the lead present in the blood is bound to erythrocytes. They have a high affinity for lead and contain the majority of the lead found in the blood stream, which makes them more vulnerable to oxidative damage than many other cells. Erythrocytes can spread lead to different organs of the body [20]. The alteration in hematological parameters serves as the earliest indicators of the effect of toxins in tissues. Therefore, abnormal alteration in blood parameters is one of the reliable indicators of toxic effects of drugs, chemicals and diseases in the body system [21]. Hematological effects of chronic toxicity of the Pb acetate (0.4%) in drinking water for 12 weeks in adult male rats showed significantly decreased red blood cell (RBC) count, while increasing total and differential WBC counts [22]. In humans, anemia may develop with Pb poisoning via the inhibition of ferrochelatase and δ -aminolevulinic acid dehydratase (ALAD), the two of many enzymes involved in heme biosynthesis. The inhibition of ferrochelatase and ALAD by lead decreases heme synthesis which leads to anemia [23].

The leaves of *C. papaya* have been shown to contain many active components that can increase the total antioxidant power in blood and reduce lipid peroxidation level, such as papain, chymopapain, cystatin, tocopherol, ascorbic acid, flavonoids, cyanogenic glucosides and glucosinolates. *C. papaya* leaf tea or extract also has a reputation as a tumor-destroying agent and antioxidant properties [24] Kadiri et al. [25], in their study on 'chemical composition, functional and antioxidant properties of *Carica papaya* seed flour, protein concentrate and protein isolate', found the plant to possess metal chelating ability which increased as the concentration of the samples increased. The present study, investigated the effect of aqueous extracts of *Carica papaya* leaves and seeds on electrolyte and hematological parameters in albino Wistar rats exposed to lead nitrate.

Materials and Methods

Collection and authentication of plant sample

Carica papaya leaves and fruits were collected from a farm in Ntan Ekere in Ibiono Local Government Area of Akwa Ibom State. The leaves were authenticated by a botanist in the Department of Science Technology, Akwa Ibom State Polytechnic.

Preparation of plant sample

The leaves of *Carica papaya* were plucked and washed with distilled water to be free from dirt or debris and other contaminants while the seeds were gotten from ripe *C. papaya* fruits which were cut into two halves so as to obtain the seeds easily. The leaves and seeds were air-dried separately for 2 weeks and blended into powdered form respectively. The blended leaves and seeds weighed 2492g and 1005g respectively. The blended leaves and seeds of *C. papaya* were soaked in 1500ml of distilled water for 24 hours in an air-tight container and were kept in a room temperature (25°C). The extracts were filtered, separating the residue from the solvent using a funnel and whatman No. 1 filter paper into separate well-labeled beakers. The solvents from both the leaves and seeds were concentrated in a water bath at 50°C for three consecutive days and slurry formed from the leaves and seeds samples were obtained and were preserved in a refrigerator at 4°C to avoid spoilage until use.

Experimental design and animal grouping

Male albino rats weighing between 174g to 196g were obtained from the animal house of the Department of Science Technology, Biochemistry unit, Akwa Ibom State Polytechnic, Ikot Osurua. Animals were housed and maintained in wooden cages (47 by 35.5 by 33cm) with wire mesh covers on the cages at 25°C under a 12:12 light/dark cycle, with free access to food and water. Experiments were carried out during the normal light/dark cycle and always started at the same hour (10 am) [26].

Thirty adult male albino rats were randomly allocated to six groups of five rats each: **Group 1 (control)**: Animals in this group served as control and received only 0.2ml of distilled water.

Group 2 (Pb(NO₃)₂): Animals in this group received 0.2ml lead nitrate at a dose of 50mg/kg body weight for 31 days by oral gavaging. The chosen dose fell within the ranges of doses applied in previous studies [27, 28].

Group 3 (CPL): Animals in this group received 0.2ml aqueous extract of *C. papaya* leaves at a dose of 500mg/kg body weight for 21 days by oral gavaging.

Group 4 (CPS): Animals in this group received 0.2ml aqueous extract of *C. papaya* seeds at a dose of 500mg/kg body weight for 21 days by oral gavaging.

Group 5 (CPL + Pb(NO₃)₂): This group of animals received 0.2ml aqueous extract of *C. papaya* leaves (500mg/kg body weight) beginning eleven (11) days after the start of lead nitrate treatment [29]. An hour after the treatment with *C. papaya* leaves, the animals receive 0.2ml of lead nitrate (50mg/kg) [30].

Group 6 (CPS + Pb(NO₃)₂): This group of animals received 0.2ml aqueous extract of *C. papaya* seeds (500mg/kg body weight) beginning eleven (11) days after the start of lead nitrate

treatment. An hour after the treatment with *C. papaya* seeds, the animals receive 0.2ml of lead nitrate (50mg/kg).

All experimental groups received treatment once daily for 3 weeks following 1 week treatment with lead nitrate for the lead treated groups. No group received above 1ml of extract or lead nitrate.

Collection of blood sample

After 24 hours fast, the animals were anesthetized with chloroform on the morning of the 4th week. The unconscious animals were pinned down on the dissecting board with the help of a tag pin and was dissected medioventrally and whole blood was collected through cardiac puncture using a syringes and needles into properly labeled EDTA sample bottles for hematological analysis. For electrolyte analysis, the blood was transferred into ordinary sample bottles and allowed to clot and serum obtained by centrifugation at 3000 rpm for 15 minutes using a bench top centrifuge.

Determination of electrolytes

Serum sodium, potassium, and bicarbonate concentrations were determined using a spectrophotometer according to the method of Henry et al. [31] while chloride and calcium concentrations were determined according to the methods of Tietz [32] and Ray Sarkar and Chauhan [33] respectively.

Determination of hematological parameters

White blood cell, granulocytes, lymphocytes, monocytes, red blood cell, packed cell volume, hemoglobin, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, mean corpuscular volume and blood platelets were determined digitally with the help of Mythic 18 analyzer. Mythic 18 analyzer is used for assessment of full blood count using whole blood. The Mythic 18 is a fully automated in-house hematology analyzer performing hematological analyses on EDTA anticoagulated blood. For counting the cellular blood components, the Mythic 18 uses the impedance technique. A cyanide-free spectrophotometry method is used to measure hemoglobin by formation of oxyhemoglobin at 555 nm. Hematocrit is measured by volume integration. The sample volume is 10 µl. The instrument can determine 16 parameters in the normal mode and 18 in the research mode: white blood cell count (WBC) with absolute number and percentage of lymphocytes (LYM), monocytes (MONO), and granulocytes (GRAN), red blood cell count (RBC), hemoglobin concentration (Hb), packed cell volume (PCV), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC). For platelet counting, a floating threshold is used whereas for RBC and WBC counts, the thresholds are predefined. Results are provided within 1 min on the liquid crystal display (LCD) which is printed out on a printer and stored in the resident memory or in a USB key.

Determination of body weights of experimental animals

Body weight of animals in each group was determined at the beginning and end of the experiment using a weighing balance. Change in body weight was calculated as: Final body weight- Initial body weight

Determination of organ weights of experimental animals

The weights of the heart, liver, spleen, lung and kidney, liver were carefully removed after the animals were sacrificed using a sensitive weighing balance and filter paper.

Statistical analysis

Results were analyzed using the IBM Statistical Product and Service Solutions (SPSS) version 25 and expressed as mean and Standard Error of Mean (SEM). Multiple Comparisons was made between control and experimental groups and among experimental groups using ANOVA. Significant differences among treatments were detected using LSD and values of $P < 0.05$ were considered statistically significant.

Results

Body weight of the experimental animals

Table 1 presents the body weight of experimental animals after treatment with lead and either *Carica papaya* leaves or seeds. Non-significant body weight gain was observed in all experimental groups except groups treated with *Carica papaya* seeds only and the lead treated group complemented with *Carica papaya* seeds which experienced weight loss.

Table 1
Body weight of the experimental animals

Group	Treatment	Initial weight (g)	Final weight (g)	Change in body weight (g)
1	Control	192.00±21.71	192.80±24.23	0.80±5.41
2	Pb(NO ₃) ₂	178.60±21.88	183.40±21.39	4.80±24.05
3	CPL	196.00±29.59	219.20±32.66	23.20±33.52
4	CPS	185.80±17.32	176.20±13.51	-9.60±6.84
5	Pb(NO ₃) ₂ + CPL	173.20±30.63	184.00±30.15	20.80±7.88
6	Pb(NO ₃) ₂ + CPS	192.80±29.68	182.60±28.67	-10.20±40.43

Pb(NO₃)₂ - Lead nitrate
 CPL - *Carica papaya* leaves
 CPS - *Carica papaya* seeds

Organ weight of experimental animals

Table 2 indicates that the liver weight of animals in all lead treated groups were non-significantly lower ($P > 0.05$) than the control and other groups. There were however no significant changes in organ weights on complementing lead treatment with *Carica papaya* leaves and seeds. The heart and spleen of the groups treated with *Carica papaya* seed were however significantly reduced ($P < 0.05$) compared to the control.

Table 2: Organ weight of the experimental animals

Group	Treatment	Heart (g)	Liver (g)	Spleen (g)	Lung (g)	Kidney (g)
1	Control	0.82± 0.77	6.15 ±0.77	0.88±0.08	1.47±0.14	1.25±0.13
2	Pb(NO ₃) ₂	0.69± 0.05	5.52± 0.48	0.74±0.06	1.31±0.20	1.22±0.07
3	CPL	0.65± 0.72	6.79± 0.92	0.67±0.08	1.33±0.15	1.10±0.16
4	CPS	0.55± 0.47*	6.10±0.73	0.60±0.04*	1.57±0.23	0.99±0.09
5	Pb(NO ₃) ₂ + CPL	0.63± 0.09	5.66±0.75	0.64±0.12	1.36±0.12	1.05±0.15
6	Pb(NO ₃) ₂ + CPS	0.66± 0.03	6.26±0.72	0.67±0.10	1.65±0.15	0.91±0.13

Values with * are significant compared to control (P< 0.05)

Pb(NO₃)₂ - Lead nitrate
 CPL - *Carica papaya* leaves
 CPS - *Carica papaya* seeds

The effect of aqueous extract of *C. papaya* leaves and seeds on Electrolyte concentration in albino Wistar rats exposed to lead nitrate

Table 3 shows a non- significant increase in serum sodium concentration of all the test groups when compared to control except for the group administered 500mg/kg of *C. papaya* leaf extract which was significantly higher when compared to control. Sodium levels were however non-significantly decreased in groups 5 and 6 when compared to group 2. A non-significant decrease (P>0.05) in the chloride levels were observed in all experimental groups when compared to control. A non-significant increase was however recorded in groups 5 and 6 when compared to group 2. Calcium levels were non-significantly increased (P>0.05) in groups 2, 4, and 5 when compared to control and non-significantly decreased (P>0.05) in group 6 when compared to group 2. Potassium levels recorded significant increase (P<0.05) in group 3 when compared to control and other test groups except for group 5. Groups 4, 5 and 6 were non-significantly decreased (P>0.05) when compared to control. Bicarbonate levels were non- significantly increased (P>0.05) in all test groups except group 3 when compared to control. The increases were higher in groups 5 and 6. The level of bicarbonate was decreased in group 2. when compared to all other groups.

Table 3

The effect of aqueous extract of *C. papaya* leaves and seeds on Electrolyte concentration in albino Wistar rats exposed to lead nitrate

Experimental group	Treatment	Electrolyte Concentration				
		Sodium Na ⁺ (mmol/l)	Chloride Cl ⁻ (mmol/l)	Calcium Ca (mmol/l)	Potassium K ⁺ (mmol/l)	Bicarbonate HCO ₃ ⁻ (mmol/l)
1	Control	146.0 ± 12.37 ^{ab}	111.20 ± 3.12 ^a	1.19 ± 0.04 ^a	5.10 ± 0.21 ^a	22.16 ± 1.40 ^a
2	Pb(NO ₃) ₂	158.60 ± 3.14 ^a	106.4 ± 2.06 ^a	1.22 ± 0.2 ^a	5.35 ± 0.08 ^a	21.84 ± 0.74 ^a
3	CPL	177.00 ± 10.17 ^{ac}	107.20 ± 2.57 ^a	1.28 ± 0.10 ^a	4.61 ± 0.09 ^b	22.08 ± 0.77 ^a
4	CPS	162.00 ± 1.02 ^a	106.80 ± 4.07 ^a	1.20 ± 0.01 ^a	5.20 ± 0.25 ^a	22.19 ± 0.96 ^a
5	Pb(NO ₃) ₂ + CPL	152.80 ± 9.7 ^a	109.60 ± 3.58 ^a	1.23 ± 0.1 ^a	4.94 ± 0.50 ^{ab}	22.72 ± 0.92 ^a
6	Pb(NO ₃) ₂ + CPS	150.80 ± 4.73 ^a	108.80 ± 2.17 ^a	1.18 ± 0.1 ^a	5.09 ± 0.16 ^a	22.94 ± 0.84 ^a

Result are expressed as mean ± standard error

Values in the same column with different superscript are significantly different at p < 0.05

Effect of aqueous extracts of *Carica papaya* leaves and seeds on hematological parameters in albino Wistar rats.

The result as presented in table 4, showed that all hematological parameters except MCV, and lymphocytes increased in the lead treated groups when compared to the control. A decrease in WBC, lymphocytes, RBC, PCV, Hb, MCH and MCHC and an increase in MCV and platelet count in the non-complemented lead group- 2, was observed when compared to the complemented lead treated group. The lead treated groups complemented with both *C. papaya* leaves and seeds (Groups 5 and 6) showed increases in WBC, lymphocytes, RBC, PCV, Hb, MCH and MCHC and a decrease MCV and platelet count when compared to the non-complemented lead treated group- 2

Table 4
The effect of aqueous extract of *C. papaya* leaves and seeds on hematological parameters in albino Wistar rats exposed to lead nitrate

Group	Treatment	WBC ($\times 10^3$)	Lymphocytes (%)	RBC ($\times 10^6$)	PCV (%)	Hb (%)	MCH (pg)	MCHC (g/dl)	MCV (fl)	Platelets ($\times 10^3$)
1	Control	7.00 \pm 1.51 ^a	57.82 \pm 7.52 ^b	6.18 \pm 0.73 ^a	33.26 \pm 3.80 ^a	12.52 \pm 1.53 ^a	20.26 \pm 0.39 ^a	37.50 \pm 0.44 ^a	54.04 \pm 0.89 ^a	448.40 \pm 210.63 ^a
2	Pb(NO ₃) ₂	12.28 \pm 1.86 ^a	68.48 \pm 5.70 ^a	8.12 \pm 0.48 ^{bd}	43.18 \pm 0.94 ^b	16.48 \pm 0.54 ^b	19.54 \pm 0.29 ^a	37.54 \pm 0.44 ^a	54.46 \pm 0.90 ^a	999.20 \pm 110.03 ^b
3	CPL	7.42 \pm 1.57 ^a	58.96 \pm 2.57 ^a	7.70 \pm 0.21 ^b	40.38 \pm 1.14 ^b	15.34 \pm 0.51 ^b	19.88 \pm 0.13 ^a	37.96 \pm 0.34 ^a	52.30 \pm 0.25 ^a	1012.80 \pm 98.86 ^b
4	CPS	7.50 \pm 1.97 ^a	65.52 \pm 4.19 ^a	9.64 \pm 0.66 ^{bc}	46.32 \pm 1.22 ^b	17.64 \pm 0.44 ^b	19.96 \pm 0.29 ^a	38.10 \pm 0.32 ^a	52.12 \pm 0.78 ^a	997.20 \pm 111.74 ^b
5	Pb(NO ₃) ₂ + CPL	10.08 \pm 0.88 ^a	72.78 \pm 2.54 ^a	8.45 \pm 0.33 ^b	43.94 \pm 1.46 ^b	16.64 \pm 1.06 ^b	20.46 \pm 0.30 ^{ac}	38.34 \pm 0.36 ^a	53.42 \pm 1.05 ^a	607.60 \pm 109.26 ^{ac}
6	Pb(NO ₃) ₂ + CPS	11.96 \pm 3.47 ^a	71.00 \pm 3.99 ^a	8.16 \pm 0.22 ^{bd}	43.46 \pm 03.02 ^b	16.68 \pm 0.45 ^b	20.44 \pm 0.18 ^{ac}	38.60 \pm 0.47 ^a	53.38 \pm 1.01 ^a	920.00 \pm 42.94 ^b

Values are expressed as mean \pm SEM. Values in the same column with different superscript are significantly different at $p < 0.05$

KEY:

- Pb(NO₃)₂ - Lead nitrate
- CPL - *Carica papaya* leaves
- CPS - *Carica papaya* seeds
- WBC - White Blood Cell
- RBC - Red Blood Cell
- PCV - Packed Cell Volume
- Hb - Hemoglobin
- MCHC - Mean Corpuscular Hemoglobin Concentration
- MCV - Mean Corpuscular Volume
- MCH - Mean Corpuscular Hemoglobin

Discussion

Exposure to lead produces a variety of adverse health effects including disorders of the central nervous, hematopoietic, hepatic and renal systems, especially in sensitive populations. This study evaluated the protective effect of aqueous extracts of *Carica papaya* leaves and seeds against electrolyte and hematological alterations in albino Wistar rats exposed to lead nitrate.

Non-significant body weight gain was observed in all experimental groups except groups treated with *Carica papaya* seeds only and the lead treated group complemented with *Carica papaya* seeds which experienced weight loss. This finding is similar to that of Okon et al. [3], who investigated the bioaccumulation of heavy metals following various oral treatments viz: Lead-exposed group with 60 mg/kg body weight of lead acetate; Cadmium-exposed group with 22.5 mg/kg of body weight of cadmium chloride, while arsenic exposed group with 18.8mg/kg body weight of arsenic trioxide and the co-exposure group with a mixture of lead acetate, (60 mg/kg) body weight, Cadmium chloride, (22.5 mg/kg) body weight, arsenic trioxide, (18.8 mg/kg) body weight for the period of 12 weeks and equally observed body weight gain in the lead intoxicated

group apart from the two groups administered *Carica papaya* seeds. Ayuba et al. [34], also reported that Oral administration of lead acetate at doses of 50mg/kg and 100mg/kg to the rats was observed to have caused an increase of 2.66% and 2.63% in the mean body weight of the rats respectively. Santana et al. [35] reported regarding caloric intake, that it was observed that in the third, fifth, and eighth weeks of supplementation, the animals that received papaya seed oil had lower caloric intake. A reduced caloric intake may be responsible for reduced body weight. The loss of weight in the two groups may also be attributed to loss in muscle and adipose tissue resulting from excessive breakdown of tissue protein and fatty acid [36].

Organ weights were relative to change in body weight of experimental groups. The liver weight of the animals in the group intoxicated with lead was lower compared to all other groups. There were significant reductions in the spleen and heart weights of animals treated with only *Carica papaya* seeds. Seifert et al. [37] in their study, observed that the spleen was found to significantly decrease in size at 24, 48, and 51 h following middle cerebral artery occlusion (MCAO) in rats. They conclude that this is possibly an immune response to tissue injury. There were however no significant changes in organ weights on complementing lead treatment with *Carica papaya* leaves and seeds. Okon et al. [3], observed a significant decrease in liver weights of animals intoxicated with Cadmium and Arsenic and a decrease in the lead (Pb) intoxicated group. The group exposed to a mixture of above three metals had the lowest liver weight.

Decrease in serum electrolytes concentrations of chloride, calcium and bicarbonate was observed in the group administered lead alone. This supports the findings of Okediran et al. [12]. A decreased plasma electrolyte level is an indication in perturbation of electrolytes metabolism due to exposure to lead thus leading to alterations in cellular functions in which these electrolytes are involved. Studies have shown that lead can replace calcium, thereby affecting key neurotransmitters like protein kinase C, which regulates long term neural excitation and memory storage, it can also affect sodium ion concentration which is responsible for numerous vital biological activities like for the purpose of cell to cell communication, uptake of neurotransmitters and regulation of uptake and retention of calcium by synaptosomes [38, 57, 40, 41]. The ionic mechanism of action for lead mainly arises due to its ability to substitute other bivalent cations such calcium, magnesium, iron and monovalent cations like sodium ions, thus affecting various fundamental biological processes of the body by damaging cellular components [39, 40]. However, complementing lead with *C. papaya* leaves and seeds increased the levels of these electrolytes. This increase could be due to the rich array of phyto-nutrients and healing property *C. papaya* leaves on many chronic diseases [42, 25]. Sodium levels were however non-significantly decreased in the groups administered lead complemented with *C. papaya* leaves and seeds when compared to lead alone group. The decrease observed in sodium and potassium levels of the lead groups complemented with *C. papaya* leaves and seeds was also reported by Eze [43] in albino wistar rats when administered with *C. papaya* seeds in a carbontetrachloride induced renal toxicity experiment. This condition occurs in cases of renal failure because the kidney loses the ability to excrete the mineral. Potassium levels however decreased in the group complemented with *C. papaya* leaves and seeds. The decrease in these two groups compared to the lead alone group indicates that *C. papaya* leaf and seed extracts could be used to modulate the alterations in renal function markers. *C. papaya* leaves and seeds may have protective effects

against acute kidney damage especially if the damage is caused by accumulation of heavy metals [44].

All hematological parameters except MCV increased in the lead treated groups when compared to the control. Terayama [45] conducted a study that showed an increase in the levels of WBC when administered with lead. Jassim and Hassan [46], also reported that lead acetate when given orally to female rats at a dose of 10 mg/kg produced significant increase in percentage of WBC count and MCV. In contrast, a research conducted by Enagbonma et al. [47] revealed that when 0.10mg/kg of lead was administered to albino rats, the RBC levels reduced when compared to the control. This could be due to the difference in dosage of lead administered. In the present study, levels of RBC, PCV, Hb, MCH, MCHC and lymphocytes were lower in the non-complemented lead treated group when compared to the complemented lead treated groups, while WBC, MCV and platelet counts were higher in the lead intoxicated group when compared to the lead group complemented with *C. papaya* leaves and seeds. A study conducted by Orji et al. [48] also showed significant reduction ($P < 0.05$) in the red blood cell count following treatment albino of rats with 75mg/kg lead. Okediran et al. [14], also reported inhibition of haem synthesis and reduction of packed cell volume following exposure to lead. In humans, anemia may develop with Pb poisoning via the inhibition of ferrochelatase and δ -aminolevulinic acid dehydratase (ALAD), the two of many enzymes involved in heme biosynthesis. The inhibition of ferrochelatase and ALAD by lead decreases heme synthesis which leads to anemia [24].

Treatment of lead intoxicated groups in the present study with both *C. papaya* leaves and seeds showed increases in RBC, PCV, Hb, MCH, MCHC and lymphocytes, and decreases in WBC, MCV and platelet count when compared to the lead intoxicated group. Sule et al. [49] and Oduola et al. [50] also reported similar trend as the plant *Carica papaya L* produced significant increases in PCV, Hb, RBC, MCV, MCH and lymphocytes. This suggests that the leaves and seeds of *C. papaya* could minimize the alterations of hematological parameters induced by lead intoxication in the albino rats. Consistent with the findings of this study are the reports of Oduola et al. [50] and Sule et al. [49], that WBC levels were decreased in rats further treated with *C. papaya* seeds and leaves respectively.

Terayama [44], Al-Saleh [51], Enagbonma et al. [47], Orji et al. [48] and Savithri et al. [52] all reported that exposure to lead significantly decreased RBC counts values and Hb in rats but when *C. papaya* leaf extract was administered to albino wistar rats, there was a significant increase in the RBC and Hb levels. The decreased RBC in the lead treated group could be as a result of iron deficiency anemia, bone marrow disorder and chronic inflammatory diseases. According to Oduola et al. [50], MCV levels were decreased in rats also treated with 100mg/kg *C. papaya* leaves. The findings of this study corroborates that of Savithri et al. [52], where platelets count revealed a significant increase in intoxicated animals following treatment of albino rat with lead acetate. Platelet level was decreased in rats treated with *C. papaya* leaves [50]. This finding however contrasts a research conducted by Enagbonma et al. [47] who reported that following the administration of 0.10mg/kg of lead to albino rats, the platelet levels reduced when compared to the control. Sule et al. [49], reported that the levels of lymphocytes increased in rats pretreated with *Carica papaya* when compared with rats administered with CCl₄ only. The increases in WBC observed in the lead intoxicated group in present study may be considered a defensive mechanism by the immune system in response to the antigen. However, treatment with *C. papaya* leaves and seeds tended to ameliorate this effect.

This may go a long way to suggesting that this plant may have influenced the defense mechanism of the test rats. So, the continuous exposure of the body systems of animals to *C. papaya* may cause lymphocytosis, which may then account for the use of these plants for medicinal purposes [53]. Earlier report made by Oladunmoye and Osho [54] revealed that the higher values of some hematological indices in rats pretreated with *Carica papaya* may be due to the inability of the toxicant to cause hemolysis resulting from the anti-inflammatory potentials inherent in the herb.

Xenobiotics can cause hemolytic anaemia when sulphhydryl groups of the erythrocyte membrane is oxidized which inflicts injury to the erythrocyte membrane [49]. *Carica papaya* was reported to rank among the first fruits for vitamin C, vitamin A, riboflavin, foliate, calcium, thiamine, iron, niacin, potassium and fibre and may probably be responsible for *Carica papaya* conferring erythropoietic properties on the pretreated rats [54]. The hematological alterations were minimized by the administration of *Carica papaya* leaves and seeds. Flavonoids, reported to be present among other active components are known to be vasculo protector and a powerful antioxidant [55].

Conclusion

The study evaluated the effect of aqueous extracts of *Carica papaya* leaves and seeds on electrolyte and hematological parameters in albino Wistar rats exposed to lead nitrate. **Increases** in sodium and potassium levels, and decreases in chloride, calcium and bicarbonate levels were observed in the lead intoxicated group. Treatment of lead intoxicated groups with CPL and CPS reversed the levels of the altered electrolytes. The hematological analysis revealed an increase in all hematological parameters except MCV when compared to the control. Levels of RBC, PCV, Hb, MCH, MCHC and lymphocytes were lower in the non-complemented lead intoxicated group when compared to the lead intoxicated groups complemented with CPL and CPS, while WBC, MCV and platelet counts were higher in the lead intoxicated group when compared to the lead intoxicated group complemented with CPL and CPS. This finding suggests that *C. papaya* leaves and seeds could be reverse alterations in electrolytes and hematological parameters induced by lead toxicity and could be harnessed for these potentials.

Declarations:

Ethical Approval

The animal study was conducted in accordance with the Ethical Guidelines for the Use of Animals in Research National Committee for Research Ethics in Science and Technology (NENT) (2018).

Data Availability:

All datasets are available on the manuscript

Disclaimer

This paper is an extended version of a **preprint** document of the same author.

References

1. Saikat M, Arka JC, Abu MT, Talha BE, Firzan N, Ameer K, Abubakr MI, Mayeen UK, Hamid O, Fahad AA, Jesus SG. Impact of heavy metals on the environment and human health: Novel therapeutic insights to counter the toxicity. *Journal of King Saud University – Science*. 2022; 34(3):101865. <https://doi.org/10.1016/j.jksus.2022.101865>
2. Balali-Mood M, Naseri K, Tahergorabi Z, Khazdair MR and Sadeghi M. Toxic Mechanisms of Five Heavy Metals: Mercury, Lead, Chromium, Cadmium, and Arsenic. *Front. Pharmacol*. 2021; 12:643972. doi: 10.3389/fphar.2021.643972
3. Okon AJ, Aluko OO, Tom UE, & Essien NE. Bioaccumulation of Heavy Metals in Liver of Albino Wistar Rats Exposed to Single and Heavy Metal Mixture. *African Journal of Environmental Health Sciences*. 2020; 7(November, 2020): 101-109. <http://www.ehsanonline.org>
4. Fernandes Azevedo B, Barros Furieri L, Peçanha FM, Wiggers GA, Frizera Vassallo P, Ronacher Simões, M, et al. Toxic effects of mercury on the cardiovascular and central nervous systems. *Biomed. Res. Int*. 2012; 949048. doi:10.1109/latincloud.2012.6508156
5. Cobbina SJ, Chen Y, Zhou Z, Wu X, Zhao T, Zhang Z, et al. Toxicity assessment due to sub-chronic exposure to individual and mixtures of four toxic heavy metals. *J. Hazard. Mater*. 2015; 294: 109–120. doi:10.1016/j.jhazmat. 2015. 03.057
6. Costa M. Review of arsenic toxicity, speciation and polyadenylation of canonical histones. *Toxicol. Appl. Pharmacol*. 2019; 375: 1–4. doi:10.1016/j.taap.2019. 05.006
7. Gazwi HSS, Yassien EE, Hassan HM. Mitigation of lead neurotoxicity by the ethanolic extract of Laurus leaf in rats. *Ecotoxicol. Environ. Safe*. 2020; 192: 110297. doi:10.1016/j.ecoenv.2020.110297
8. Reza B, Ali N, Azhdar H, Alireza A, Ali K. Effects of low-level lead exposure on blood pressure and function of the rat isolated heart. *Indian J Pharmacol*. 2008; 40(2): 69-72
9. Teijón C, Olmo R, Blanco D, Romer A, Teijón JM. Low doses of lead: effect on reproduction and development in rats. *Biol Trace Elem Res*. 2006; 111: 51- 165.
10. Durgut R, Koc A, Gonenci R., Bal R, Celik S, Guzel M, Altug M, Atesoglu E. Effects of high dose lead intoxication on liver, kidneys, heart, brain and blood in rabbits: an experimental study. *J Appl Biol Sci*. 2008; 2: 11- 18.
11. Garrettson LK. *Lead in Clinical Management of poisoning and drug over-dose*. In: Haddad LM, et al. (Eds.), Philadelphia: W Saunders Company; 1990. p. 1017-23.
12. Okediran BS, Kasali OB, Omotainse SO, Akinloye OA. Haemato-biochemical alterations as Biomarkers of lead induced toxicity in male Wistar rats. *Bang J Vet Med*. 2016; 14(2): 227-232. DOI:10.23880/act-16000150
13. Sidhu P, Nehru B. Lead intoxication: Histological and oxidative damage in rat cerebrum and cerebellum. *J Trace Elem Exp Med*. 2004; 17(1): 45-53.
14. Taib NT, Jarrar BM, Mubarak M. Ultrastructural alterations in hepatic tissues of white rats (*Rattus norvegicus*) induced by lead experimental toxicity. *Saudi J Biol Sci*. 2004; 11(1): 11-20

15. Flora SJS, Flora G, Saxena G. Environmental Occurrence, Health Effects and Management of Lead Poisoning. In: Casas SB (Eds.), *Lead: Chemistry, Analytical Aspects, Environmental Impact and Health Effects*. Netherlands: Elsevier: 2006. P. 158-228.
16. Berrahal AA, Lasram M, El-Elj N, Kerkeni A, Gharbi N, et al. Effect of age-dependent exposure to lead on hepatotoxicity and nephrotoxicity in male rats. *J Environ Toxicol*. 2011; 26(1): 68-78.
17. Hasona NA, Elsbali A. Evaluation of electrolytes imbalance and dyslipidaemia in diabetic patients. *Med Sci*. 2016; 4(2): 1-4.
18. Kaneko JJ, Harvey JW, Bruss ML. *Clinical Biochemistry of Domestic Animals, 6th (Edn.)*. London: Academic press; 2006. p. 117-38
19. Mohammed AA, Mohd NMH, Abd WH, Mohd YMS, Mohd, AR. The detrimental effects of lead on human and animal health. *Veterinary World*. 2016; 9(6): 660-671 doi: 10.14202/vetworld.2016.660-671
20. Sivaprasad R, Nagaraj M, Varalakshmi P. Combined efficacies of lipoic acid and meso-2,3-dimercaptosuccinic acid on lead induced erythrocyte membrane lipid peroxidation and antioxidant status in rat. *Human and Experimental Toxicology*, 2003; 22:183-192.
21. Abbiw OU, Ajali HJ. Effects of Heavy Metals on the Reaction Rate and on the regeneration of Reactive Oxygen Species. *Journal of Biochemical Pharmacology* 2005; 37: 409- 413.
22. Mugahi MN, Heidari Z, Sagheb HM, Barbarestani M. Effects of chronic lead acetate intoxication on blood indices of male adult rat. *DARU J. Pharm. Sci*. 2003;11(4): 147-141.
23. Mense SM, Zhang L. Heme: a versatile signaling molecule controlling the activities of diverse regulators ranging from transcription factors to MAP kinases. *Cell Res*. 2006; 16 (8): 681-692. doi:10.1038/sj.cr.7310086
24. Kong YR, Jong YX, Balakrishnan M, Bok ZK, Weng JKK, Tay KC, Goh BH, Ong YS, Chan KG, Lee LH, Khaw KY. Beneficial Role of *Carica papaya* Extracts and Phytochemicals on Oxidative Stress and Related Diseases: A Mini Review. *Biology*, 2021; 10(4): 287. <https://doi.org/10.3390/biology10040287>
25. Kadiri O, Akanbi CT, Olawoye BT, Gbadamosi SO. Characterization and antioxidant evaluation of phenolic compounds extracted from the protein concentrate and protein isolate produced from pawpaw (*Caricapapaya* Linn.) seeds. *International Journal of Food Properties*, 2017; 20(11): 2423-2436.
26. National Committee for Research Ethics in Science and Technology (NENT). Ethical Guidelines for use of Animals in Research. Oslo; 2018
27. Khan MS, Mostofa M, Jahan MS, Sayed MA, Hossain MA. Effect of garlic and vitamin Bcomplex in lead acetate induced toxicities in mice. *Bangladesh Journal of Veterinary Medicine*, 2008; 6(2): 203-10.
28. Haque MM, Awal MA, Mostofa M, Sikder MM, Hossain MA. Effects of calcium carbonate, potassium iodide and zinc sulphate in lead induced toxicities in rat model. *Bangladesh Journal of Veterinary Medicine*, 2006; 4(2): 23-7.
29. Sharma V, Sharma, A, Kansal L. The effect of oral administration of *Allium sativum* extracts on lead nitrate induced toxicity in male mice. *Food and chemical toxicology*, 2010; 48(3): 928-36.

30. Abdel moneim AE, Dkhil MA, Al-Quaraishy S. The protective effect of flaxseed oil on lead acetate induced renal toxicity in rats. *Food chemistry and toxicology*, 2011; 56 (4): 434-437.
31. Henry RJ, Cannon DC, Winklemen JW. *Clinical Chemistry-Principle and Technique*, 2nd Edition, New York: Harper and Row; 1974. p. 673-676.
32. Tietz NW. *Fundamentals of Clinical Chemistry*, 2nd edition. Philadelphia: Saunders; 1976. p. 876.
33. Ray Sarkar BC, Chauhan UPS. A new method for determining micro quantities of calcium in biological materials, *Analytical Biochemistry*, 1967; 20(1): 155-166.
34. Ayuba Y, Ekanem AU, Garba SH. Effect of Oral Administration of Lead Acetate Exposure on the Histology of the Testis and Testicular Sperm Concentration in Wistar Albino Rats Scholars *Journal of Applied Medical Science*, 2017; 5(6D): 2337-2344.
35. Santana, L.F.; do Espirito Santo, B.L.S.; Tatará, M.B.; Negrão, F.J.; Croda, J.; Alves, F.M.; de Oliveira Filiú, W.F.; Cavalheiro, L.F.; Nazário, C.E.D.; Asato, M.A.; et al. Effects of the Seed Oil of *Carica papaya* Linn on Food Consumption, Adiposity, Metabolic and Inflammatory Profile of Mice Using Hyperlipidic Diet. *Molecules*, 2022; 27: 6705. <https://doi.org/10.3390/molecules27196705>
36. El-Zayat SR., Sibaii H, El-Shamy KA. Physiological process of fat loss. *Bulletin of the National Research Centre*, 2019; 43: 208. <https://doi.org/10.1186/s42269-019-0238-z>
37. Seifert HA, Hall AA, Chapman CB, Collier LA, Willing AE, Pennypacker KR. A transient decrease in spleen size following stroke corresponds to splenocyte release into systemic circulation. *J Neuroimmune Pharmacol*. 2012 Dec;7(4):1017-24. doi: 10.1007/s11481-012-9406-8. Epub 2012 Oct 6. PMID: 23054371; PMCID: PMC3518577.
38. Bressler J, Kim KA, Chakraborti T, Goldstein G. Molecular mechanisms of lead neurotoxicity. *Neurochem Res*. 1999; 24(4): 595-600.
39. Gwaltney-Brant SM. Heavy Metals, *Handbook of Toxicologic Pathology*. 2nd (Edn.), In: Haschek WM, et al. (Eds.), Academic Press; 2002. p. 701-733.
40. Hontela A, Lacroix A. Heavy Metals in endocrine disruptors: Biological Basis for Health effects in Wildlife and Human. In: Carr J, et al. (Eds.), United Kingdom: Oxford University Press; 2006. p. 356-374.
41. Lidsky TI, Schneider JS. Lead neurotoxicity in children: Basic mechanisms and clinical correlates. *Brain*. 2006; 126(1): 5-19.
42. Noriko O, Nam H, Dang E, Akira K, Satoshi I, Chikao M. Aqueous extract of *Carica papaya* leaves exhibit anti-tumor activity and immunomodulatory effects. *Journal of Ethnopharmacology*, 2010; 127: 760-767.
43. Eze KN. Dose dependent Attenuating effects of aqueous extract of *C. papaya* seed on carbon tetrachloede induced renal Toxicity in rats. Delta state: University Department of physiology, 2012; 2(4): 124 – 127.
44. Nor SA, Shafiy YSZ, Ibrahim AT, Morhammed TR. Effect of green and ripe *C. papaya* eapicarp extract on wound healing and during pregnancy. *Food Chemistry Toxicology*, 2008; 46-(7): 2384-2389.
45. Terayama K. Effects of lead on electrophoretic mobility, membrane sialic acid, deformability and survival of rat erythrocytes. *Industrial Health*, 1993; 31: 113-126.

46. Jassim HM, Hassan AA. Changes in some blood parameters in lactating female rats and their pups exposed to lead: Effects of vitamins C and E. *International Journal on Violence and School*, 2011; 25(10): 1-7.
47. Jesuorsewun EB, Ebikere I II, Ozede IN, Eghomwanre AF. Hematobiochemical changes of lead Poisoning and amelioration with Coconut (*Cocosnucifera* L.) Water in wistar albino rats. *J. Appl. Sci. Environ. Manage.* 2016; 20(1): 89- 94. <https://doi.org/10.4314/jasem.v20i1.11>
48. Orji OU, Ibiam UA, Aja PM, Obasi OD, Ezeani N, Alope C, Anayo S, Inya-Agha OR. Effect of Ethanol Extract of *Ruspolia hypocrateriformis* Leaf on Hematological Parameters in Lead Poisoned Albino Rats. *World Journal of Medical Sciences*. 2016; 13 (4): 225-235.
49. Sule OJ, Abdu AR, Kiridi K. Effect of *Carica papaya* (L) Leaves on Hematological Parameters in Ccl4-induced Wistar Albino Rats *British Journal of Medicine & Medical Research*. 2016; 16(3): 1-6, 2016, Article no.BJMMR.13686 DOI: 10.9734/BJMMR/2016/13686
50. Oduola T, Adeniyi FA, Ogunyemi EO, Bello IS, Idowu TO, Subair HG. Toxicity studies on an unripe *Carica papaya* aqueous extract: biochemical and hematological effects in wistar albino rats. *Journal of Medicinal Plants Research*, 2007; 1(1): 001-004.
51. Al-Saleh I. Biochemical and clinical consequences of lead poisoning. *Medicinal Research Review*, 1994; 14: 415-486.
52. Savithri Y, Sekhar PR, Doss PJ. Changes in hematological profiles of albino rats under chlorpyrifos toxicity. *Int. J. Pharma. Bio. Sci.*, 2010; 1: 1-7.
53. Keenwe M, Bekalo I. Ethno Veterinary Medicine in Kenya, a field manual of practical animal health care practices. Nairobi: ITK and IIRR; 1996.
54. Oladunmoye MK, Osho IB. Anti-inflammatory activity of ethanolic leaf extract from *Carica papaya* in rats orogastrically dosed with *Salmonella typhi* and *Staphilococcus aureus*. *Journal of Plant Sciences*, 2007; 2(4): 447-452.
55. Krishna KL, Paridhavi M, Jagruti AP. Review on nutritional, medicinal and pharmacological properties of Papaya (*Carica papaya* Linn.). *Natural product Radiance*. 2008; 7(4): 364-373.