

## HR-Stability of Generalized Quadratic functional equation in Random Norm spaces

**Abstract:** The purpose of this paper to propose a finite variable generalized quadratic functional equation with solution. Also investigate Hyers Ulam stability in Random normed space by direct method.

**Keywords:** Quadratic functional equation, Hyers- Ulam stability, Random normed space.

**Mathematics subject classification:** 54E40, 39B82, 46S50.

**1. Introduction:** Functional equations play an important part in the study of stability. The first question regarding stability is raised by Ulam [15] for group homomorphisms. After that Hyers [1] proved the stability problem for Cauchy’s functional equation in Banach spaces. Hyers’ result further expanded by many authors i.e., Aoki [18], Th.M. Rassias[22], J.M. Rassias[21], Gavruta [12] and many more.

Recently the stability of many functional equations in various spaces like Banach spaces, modular spaces, fuzzy normed spaces and Random normed spaces etc. have been established by researchers[8,9,17-19]. Now we obtain the Hyers- Ulam stability of quadratic function equation  $\sum_{i=1}^n \phi(-2v_i + \sum_{j=1, i \neq j}^n v_j) = (n - 6) \sum_{1 \leq i < j \leq n} \phi(v_i + v_j) - (n^2 - 8n + 3) \sum_{i=1}^n \phi(v_i)$  in Random normed spaces.

We adopt the usual terminology, notions and conventions of the theory of random normed spaces as in [8].

Throughout,  $\Delta^+$  denotes the distribution functions spaces , i.e., the space of all mapping  $f: \mathbb{R} \cup (-\infty, \infty) \rightarrow [0,1]$  such that  $f$  is left continuous and increasing on  $\mathbb{R}$ ,  $f(0)=0$  and  $f(+\infty)=1$ .  $D^+$  subset of  $\Delta^+$  consisting of all functions  $V$  of  $\Delta^+$  for which  $\ell^- V(+\infty)=1$  where  $\ell^- f(s)$  denotes  $\ell^- f(s) = \lim_{t \rightarrow s^-} f(t)$ . The space  $\Delta^+$  is partially order by usual wise ordering of functions, i.e.,

$f \leq g \Leftrightarrow f(s) = g(s), \forall s \in \mathbb{R}$ . The maximal element for  $\Delta^+$  in this order is the distribution function  $\epsilon_0(s) = \begin{cases} 0, & \text{if } s \leq 0 \\ 1 & \text{if } s > 0 \end{cases}$

**Definition 1.1:** A Random Normed space (RN-space) is a triple  $(V, \mu, T)$ , where  $V$  is a vector space,  $T$  is a continuous t-norm and  $\mu: V \rightarrow D^+$  satisfying the following conditions:

(R<sub>1</sub>).  $\mu_v(t) = \epsilon_0(t)$  for all  $t > 0$  if and only if  $v=0$ ,

(R<sub>2</sub>).  $\mu_{av}(t) = \mu_v\left(\frac{t}{|a|}\right)$  for all  $v \in V, t \geq 0$  and  $a \in \mathbb{R}$  with  $a \neq 0$ ,

(R<sub>3</sub>).  $\mu_{v+u}(t+s) \geq T(\mu_v(t), \mu_u(s))$  for all  $v, u \in V$  and  $t, u \geq 0$ .

**Definition 1.2.** Let  $(V, \mu, T)$  be a RN- space.

- (i) A sequence  $\{v_n\}$  in  $V$  is said to be convergent to  $v \in V$  if  $\lim_{n \rightarrow \infty} \mu_{v_n - v}(t) = 1, t > 0$ .
- (ii) A sequence  $\{v_n\}$  in  $V$  is said to be Cauchy sequence if  $\lim_{n \rightarrow \infty} \mu_{v_n - v_m}(t) = 1, t > 0$ .
- (iii) A RN-space  $(V, \mu, T)$  is complete if every Cauchy sequence is convergent in  $V$ .

**2. General solution**

Theorem 2.1. If an even function  $\phi: V \rightarrow W$  satisfies the functional equation (1.1) for all  $v_1, v_2, \dots, v_n \in V$ , then the function  $\phi$  is quadratic.

Proof. Let function  $\phi: V \rightarrow W$  satisfies (1.1). Taking  $v_1 = v_2 = \dots = v_n = 0$  in (1.1), we have  $\phi(0) = 0$ . Replacing  $(v_1, v_2, \dots, v_n)$  by  $(v, 0, \dots, 0)$  in (1.1), we get

$$\phi(-2v) + (n - 1)\phi(v) = (n - 6)(n - 1)\phi(v) - (n^2 - 8n + 3)\phi(v)$$

$$\phi(-2v) = (n^2 - 7n + 6 - n^2 + 8n - 3 - n + 1)\phi(v)$$

$$\phi(-2v) = 4\phi(v)$$

Since  $\phi$  is even mapping, so

$$\phi(2v) = 2^2\phi(v), \text{ for all } v \in V.$$

Now, replace  $v$  by  $2v$ , we have

$$\phi(2^2 v) = 2^4\phi(v)$$

Continue like this, we generalize

$$\phi(2^n v) = 2^{2n}\phi(v)$$

For all  $v \in V$  and for any  $n \geq 0$ .

Similarly, we have

$$\phi(2^{-n} v) = 2^{-2n}\phi(v)$$

Now, replacing  $(v_1, v_2, \dots, v_n)$  by  $(v_1, v_2, \dots, 0)$ , we get our desired result of  $\phi$ .

Remark: Let  $V$  be a linear space and a mapping  $\phi: V \rightarrow W$  satisfies the functional equation (1.1), then

- (i)  $\phi(r^n v) = r^n \phi(v)$  for all  $v \in V, r \in \mathbb{Q}, t \in \mathbb{Z}$ .
- (ii)  $\phi(v) = v \phi(1)$  for all  $v \in V$  if  $\phi$  is continuous.

For notational handiness, we denote  $(V, \Psi, T)$ , and  $(W, \Psi, T)$  are Complete RN spaces and define a mapping  $\phi: V \rightarrow W$  by

$$D\phi(v_1, v_2, \dots, v_n) = \sum_{i=1}^n \phi(-2v_i + \sum_{j=1, i \neq j}^n v_j) - (n - 6) \sum_{1 \leq i < j \leq n} \phi(v_i + v_j) + (n^2 - 8n + 3) \sum_{i=1}^n \phi(v_i) \tag{1.2}$$

for all  $v_1, v_2, \dots, v_n \in V$ .

### 3. Hyers-Ulam Stability : Direct method

**Theorem 3.1.** If an even mapping  $\phi: V \rightarrow W$  with  $\phi(0) = 0$  for which there exists a mapping  $\Phi: V^n \rightarrow D^+$  for some  $0 < \alpha < 4$ ,

$$\Phi_{2v_1, 2v_2, \dots, 2v_n}(\epsilon) \geq \Phi_{v_1, v_2, \dots, v_n} \left( \frac{\epsilon}{\alpha} \right) \tag{3.1}$$

$$\text{And } \lim_{t \rightarrow \infty} \Phi_{2^t v_1, 2^t v_2, \dots, 2^t v_n} (2^{2t} \epsilon) = 1 \tag{3.2}$$

for all  $v_1, v_2, \dots, v_n \in V$  and all  $\epsilon > 0$  such that

$$\mu_{D\phi(v_1, v_2, \dots, v_n)}(\epsilon) \geq \Phi_{v_1, v_2, \dots, v_n}(\epsilon). \tag{3.3}$$

Then there exists a unique quadratic mapping  $Q_2: V \rightarrow W$  satisfying the functional equation (1.1) with  $\mu_{Q_2(v) - \phi(v)}(\epsilon) \geq \Phi_{v, 0, \dots, 0}((2^2 - \alpha)2\epsilon)$  (3.4)

for all  $v \in V$  and all  $\epsilon > 0$ . The mapping  $Q_2: V \rightarrow W$  is defined by

$$\mu_{Q_2(v)}(\epsilon) = \lim_{t \rightarrow \infty} \mu_{\frac{\phi(2^t v)}{2^{2t}}}(\epsilon) \tag{3.5}$$

for all  $v \in V$  and all  $\epsilon > 0$ .

**Proof:** Replace  $(v_1, v_2, \dots, v_n)$  by  $(v, 0, \dots, 0)$  in (3.3), we have

$$\begin{aligned} \mu_{\phi(2v) - 2^2 \phi(v)}(\epsilon) &\geq \Phi_{v, 0, \dots, 0}(\epsilon). \\ \mu_{\frac{\phi(2v)}{2^2} - \phi(v)}(\epsilon) &\geq \Phi_{v, 0, \dots, 0}(2^2 \epsilon). \end{aligned} \tag{3.6}$$

Replace  $v$  by  $2^t v$  in (1.7), we have

$$\mu_{\frac{\phi(2^{t+1} v)}{2^2} - \phi(2^t v)}(\epsilon) \geq \Phi_{2^t v, 0, \dots, 0}(2^2 \epsilon)$$

$$\begin{aligned} \frac{\mu_{\phi(2^{t+1}v)} - \phi(2^t v)}{2^{2(t+1)}}(\varepsilon) &\geq \Phi_{3^t v, 0, \dots, 0}(2^{2(t+1)}\varepsilon) \\ &\geq \Phi_{v, 0, \dots, 0}\left(\frac{2^{2(t+1)}\varepsilon}{\alpha^t}\right). \end{aligned} \tag{3.7}$$

for all  $v \in V$  and all  $\varepsilon > 0$ . Since

$$\frac{\phi(2^m v)}{2^{2m}} - \phi(v) = \sum_{i=0}^{m-1} \frac{\phi(2^{i+1}v)}{2^{2(i+1)}} - \frac{\phi(2^i v)}{2^{2i}} \tag{3.8}$$

From (3.7) and (3.8), we have

$$\begin{aligned} \frac{\mu_{\phi(2^m v)} - \phi(v)}{2^{2m}}\left(\sum_{i=0}^{m-1} \frac{\alpha^i}{2^{2(i+1)}} \frac{\varepsilon}{2}\right) &\geq \Phi_{v, 0, \dots, 0}(\varepsilon). \\ \frac{\mu_{\phi(2^m v)} - \phi(v)}{2^{2m}}(\varepsilon) &\geq \Phi_{v, 0, \dots, 0}\left(\frac{2\varepsilon}{\sum_{i=0}^{m-1} \frac{\alpha^i}{2^{2(i+1)}}}\right) \end{aligned} \tag{3.9}$$

Replace  $v$  by  $2^2 v$ , we got

$$\frac{\mu_{\phi(2^{m+n} v)} - \phi(2^n v)}{2^{2(m+n)}}(\varepsilon) \geq \Phi_{v, 0, \dots, 0}\left(\frac{2\varepsilon}{\sum_{i=n}^{m+n-1} \frac{\alpha^i}{2^{2(i+1)}}}\right)$$

For all  $v \in V$  and all  $\varepsilon > 0$ . As  $\Phi_{v, 0, \dots, 0}\left(\frac{2\varepsilon}{\sum_{i=n}^{m+n-1} \frac{\alpha^i}{2^{2(i+1)}}}\right) \rightarrow 1$  as  $m, n \rightarrow \infty$  then  $\left\{\frac{\phi(2^n v)}{2^{2n}}\right\}$  is a

Cauchy sequence in  $(W, \mu, T)$ , Since  $(W, \mu, T)$  is complete RN- space, thus sequence  $\left\{\frac{\phi(2^n v)}{2^{2n}}\right\}$  converges to some  $Q_2(v) \in W$ . fix  $v \in V$  and put  $n=0$ , we obtain

$$\frac{\mu_{\phi(2^m v)} - \phi(v)}{2^{2m}}(\varepsilon) \geq \Phi_{v, 0, \dots, 0}\left(\frac{2\varepsilon}{\sum_{i=0}^{m-1} \frac{\alpha^i}{2^{2(i+1)}}}\right)$$

And so, for every  $\delta > 0$ , we get

$$\begin{aligned} \mu_{Q_2(v) - \phi(v)}(\varepsilon + \delta) &\geq T\left(\mu_{Q_2(v) - \frac{\phi(2^m v)}{2^{2m}}}(\delta), \mu_{\frac{\phi(2^m v)}{2^{2m}} - \phi(v)}(\varepsilon)\right) \\ &\geq T\left(\mu_{Q_2(v) - \frac{\phi(2^m v)}{2^{2m}}}(\delta), \Phi_{v, 0, \dots, 0}\left(\frac{2\varepsilon}{\sum_{i=0}^{m-1} \frac{\alpha^i}{2^{2(i+1)}}}\right)\right) \end{aligned}$$

For all  $v \in V$  and all  $\varepsilon, \delta > 0$ . Taking the limit  $m \rightarrow \infty$ ,

$$\mu_{Q_2(v)-\phi(v)}(\varepsilon + \delta) \geq \Phi_{v,0,\dots,0}(2(2^2 - \alpha)\varepsilon) \tag{3.10}$$

For all  $v \in V$  and all  $\varepsilon, \delta > 0$ . Since  $\delta$  was arbitrary, taking  $\delta \rightarrow 0$ ,

$$\mu_{Q_2(v)-\phi(v)}(\varepsilon) \geq \Phi_{v,0,\dots,0}(2(2^2 - \alpha)\varepsilon)$$

For all  $v \in V$  and all  $\varepsilon > 0$ . Replacing  $(v_1, v_2, \dots, v_n)$  by  $(2^t v_1, 2^t v_2, \dots, 2^t v_n)$  in (3.7),

$$\mu_{DQ_2(2^t v_1, 2^t v_2, \dots, 2^t v_n)}(\varepsilon) \geq \Phi_{2^t v_1, 2^t v_2, \dots, 2^t v_n}(2^{2t} \varepsilon).$$

Since  $\lim_{t \rightarrow \infty} \Phi_{2^t v_1, 2^t v_2, \dots, 2^t v_n}(2^{2t} \varepsilon) = 1$ , so  $Q_2$  satisfies the functional equation (1.1). To prove the uniqueness of Quadratic mapping  $Q_2$ . Assume that there exists another Quadratic mapping  $Q'_2$ , which satisfies inequality (3.10). Fix  $v \in V$ . Clearly,  $Q_2(2^t v) = 2^{2t} Q_2(v)$  and  $Q'_2(2^t v) = 2^{2t} Q'_2(v)$  for all  $v \in V$ . from (3.10), we have

$$\mu_{Q_2(v)-Q'_2(v)}(\varepsilon) = \lim_{t \rightarrow \infty} \mu_{\frac{Q_2(2^t v)}{2^{2t}} - \frac{Q'_2(2^t v)}{2^{2t}}}(\varepsilon)$$

**Theorem 3.2.** If an even mapping  $\phi: V \rightarrow W$  with  $\phi(0) = 0$  for which there exists a mapping  $\Phi: V^n \rightarrow D^+$  for some  $0 < \alpha < 4$ ,

$$\Phi_{\frac{v_1}{2}, \frac{v_2}{2}, \dots, \frac{v_n}{2}}(\varepsilon) \geq \Phi_{v_1, v_2, \dots, v_n}(\alpha \varepsilon)$$

And  $\lim_{t \rightarrow \infty} \Phi_{\frac{v_1}{2^t}, \frac{v_2}{2^t}, \dots, \frac{v_n}{2^t}}(\frac{\varepsilon}{2^{2t}}) = 1$

for all  $v_1, v_2, \dots, v_n \in V$  and all  $\varepsilon > 0$  such that

$$\mu_{D\phi(v_1, v_2, \dots, v_n)}(\varepsilon) \geq \Phi_{v_1, v_2, \dots, v_n}(\varepsilon).$$

Then there exists a unique quartic mapping  $Q_2: V \rightarrow W$  satisfying the functional equation (1.1) with  $\mu_{Q_2(v)-\phi(v)}(\varepsilon) \geq \Phi_{v,0,\dots,0}(\frac{\varepsilon}{(2^2-\alpha)2})$

for all  $v \in V$  and all  $\varepsilon > 0$ . The mapping  $Q_2: V \rightarrow W$  is defined by

$$\mu_{Q_2(v)}(\varepsilon) = \lim_{t \rightarrow \infty} \mu_{2^{2t} \phi(\frac{v}{2^t})}(\varepsilon)$$

for all  $v \in V$  and all  $\varepsilon > 0$ .

**Corollary 3.3.** If an even mapping  $\phi: V \rightarrow W$  with  $\phi(0) = 0$  for which there exists a mapping  $\Phi: V \rightarrow D^+$  satisfying

$$\mu_{D\phi(v_1, v_2, \dots, v_n)}(\varepsilon) \geq \Phi_{\sum_{i=1}^n \|v_i\|}(\varepsilon).$$

Then there exists a unique quadratic mapping  $Q_2: V \rightarrow W$  satisfying the functional equation (1.1) with  $\mu_{Q_2(v)-\phi(v)}(\varepsilon) \geq \Phi_{\|v\|^\theta}((2^2 - 2^p)2\varepsilon)$

for all  $v \in V$ , where  $p < 2$  and all  $\varepsilon > 0$

We take  $\alpha = 2^{p-2}$  and  $v_1, v_2, \dots, v_n = \sum_{i=1}^n \|v_i\|^\theta$  in the theorem 3.1.

## References

1. D. H. Hyers, "On the stability of the linear functional equation," Proceedings of the National Academy of Sciences of the United States of America, vol. 27, no. 4, pp. 222–224, 1941.
2. D. H. Hyers, G. Isac, and T. M. Rassias, "Stability of functional equations in several variables," in Progress in Nonlinear Differential Equations and their Applications, vol. 34, Birkhäuser Boston, Inc., Boston, MA, 1998.
3. G. Sadeghi, "A fixed point approach to stability of functional equations in modular spaces," Bulletin of the Malaysian Mathematical Sciences Society, vol. 37, no. 2, pp. 333–344, 2014.
4. G. Zamani Eskandani and J. M. Rassias, "Stability of general a cubic functional equations in modular spaces," Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas, vol. 112, no. 2, pp. 425–435, 2018.
5. H. Nakano, *Modular Semi-Ordered Linear Spaces*, Maruzen Co., Ltd., Tokyo, 1950.
6. K. Jun, Y. Lee, On the Hyers-Ulam-Rassias stability of a Pexiderized quadratic inequality, *Math. Inequal. Appl.*, 4, 93–118, 2001.
7. K. Tamilvanan, J. R. Lee, C. Park, Ulam stability of a functional equation deriving from quadratic and additive mappings in random normed spaces, *AIMS Math.*, 6 pp. 908—924, 2020.
8. M. A. Khamsi, "Fixed point theory in modular function spaces," in Recent advances on metric fixed point theory, vol. 48, pp. 31–57, Univ. Sevilla, Seville, 1995, Ciencias.
9. M. A. Khamsi, "Quasicontraction mappings in modular spaces without  $\Delta_2$ -condition," *Fixed Point Theory and Applications*, vol. 2008, no. 1, Article ID 916187, 2008.
10. N. Alessa, K. Tamilvanan, G. Balasubramanian and K. Loganathan, Stability results of the functional equation deriving from quadratic function in random normed spaces, *6(3)*, 2385-2397, 2020.
11. N. Uthirasamy, K. Tamilvanan, H. K. Nashine and R. George, Solution and stability of quartic functional equations in Modular spaces by using Fatou property, *Journal of Function Spaces*, 2022.
12. P. Gavruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, *Journal of Mathematical Analysis and Applications* 184(3), 431 – 436, 1994.
13. S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific Publishing Co., Inc., River Edge, NJ, 2002.

14. S. Koshi and T. Shimogaki, "On F-norms of quasi-modular spaces," *Journal of the Faculty of Science, Hokkaido University. Series, vol. I, no. 15*, pp. 202–218, 1961.
15. S. M. Ulam, "Problems in modern mathematics," in *Science Editions*, John Wiley & Sons, Inc., New York, 1964.
16. S. S. Jin, Y. H. Lee, "On the stability of the functional equation deriving from quadratic and additive function in random normed spaces via fixed point method," *J. Chungcheong Math. Soc.*, 25 , 51–63, 2012.
17. S. Yamamuro, "On conjugate spaces of Nakano spaces," *Transactions of the American Mathematical Society*, vol. 90, no. 2, pp. 291–311, 1959.
18. T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, no. 1-2, pp. 64–66, 1950.
19. T. Dominguez Benavides, M. A. Khamsi, and S. Samadi, "Uniformly Lipschitzian mappings in modular function spaces," *Nonlinear Analysis: Theory, Methods and Applications*, vol. 46, no. 2, pp. 267–278, 2001.
20. T. Dominguez-Benavides, M. A. Khamsi, and S. Samadi, "Asymptotically regular mappings in modular function spaces," *Scientiae Mathematicae Japonicae*, vol. 53, no. 2, pp. 295–304, 2001.
21. T. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
22. Th. M. Rassias, K. Shibata, "Variational problem of some quadratic functionals in complex analysis," *J. Math. Anal. Appl.*, 228 , 234–253, 1998.
23. Th. M. Rassias, "On the stability of the quadratic functional equation and its applications," *Studia Univ. Babeş-Bolyai*, 89–124, 1998.
24. W. M. Kozłowski, "Modular function spaces," in *Monographs and Textbooks in Pure and Applied Mathematics*, vol. 122, Marcel Dekker, Inc., New York, 1988.
25. Z. Gajda, "On stability of additive mappings," *Intern. J. Math. Math. Sci.*, 14 , 431–434, 1991.