

Effect of Nitrogen and Iron on Growth and Yield of Foxtail millet (*Setaria italica*L.)

ABSTRACT:

The Field experiment was conducted during *Zaid* season 2022 at experimental field of Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj and Uttar Pradesh, India. The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.3), low in organic carbon (0.48%), available nitrogen (230 kg/ha), available phosphorus (13.60 kg/ha) and available potassium (215.4 kg/ha). The treatments consist of 3 levels of Nitrogen (40 Kg, 50 Kg, 60 Kg/ha) as basal application and Iron (0.2, 0.4, 0.6 %) as foliar spray along with control. The experiment was layout in Randomized Block Design with Ten treatments each replicated thrice. Growth attributes namely higher plant height (101.57cm), maximum dry weight/plant (14.83 g), more number of tillers/hill (8.87) and yield attributes namely higher panicle length (18.61 cm), grains/panicle (1389.30), grain yield (17.96q/ha) and straw yield (25.32q/ha) were observed with application of nitrogen 60Kg/ha and iron 0.6%.

Keywords: *Foxtail millet, Nitrogen, Iron, Growth Parameters, Yield attributes, Yield.*

Introduction:

Foxtail millet (*Setaria italica* L.) is one of the oldest cultivated millet and most economically important species of the genus *Setaria*. It ranks second in the total world production of millets and it continues to have an important place in world agriculture providing food for millions of people in arid and semiarid regions. It is native to China and regarded as an elite drought-tolerant crop. In India, Andhra Pradesh, Karnataka and Tamil Nadu are the major foxtail millet growing states contributing about 79 per cent of the total cultivated area of about 6 lakh hectares with production of 3 lakh tonnes and productivity of 602 Kg/ha during 2017-2018. It has the total production of 2.29 m.t from the area of 1.057 m.ha in the world.

Foxtail millet has a very good nutritional profile and is ahead of rice and wheat in terms of protein, fiber, minerals and vitamins. It has good nutritive values, proteins (12.3 g), carbohydrates (60.9 g), fat (4.3 g), crude fiber (8.0 g), calcium (3.1 g), vitamins and thiamin (50 mg) per 100g. About 8-14% oil is being extracted from the bran of foxtail millet, which can be used as oil after refinement (Munirathnam *et al.*, 2006). Unlike rice, foxtail millet releases glucose slowly without affecting the metabolism of the human body with low glyce

ric index.

Nitrogen is one of the most important fertilizers for crop growth and is widely used in the farming community. Nitrogen is required for the formation of amino acids, proteins, nucleic acids, enzymes, co-enzymes, and alkaloids. In the presence of solar energy, nitrogen-containing chlorophyll fixes atmospheric carbon dioxide as carbohydrates, improving crop quality. Nitrogen fertilization improves food grain protein quality by increasing the proportion of glutamic acid. Grains should contain more proline, methionine, and less Lysine and Leucine.

Iron is a micronutrient that is required by almost all living organisms because it is involved in metabolic processes such as DNA synthesis, respiration, and photosynthesis. Iron also activates many metabolic pathways and is a prosthetic group constituent of many enzymes. The primary causes of iron chlorosis are an imbalance between the solubility of iron in soil and the plant's demand for iron. Although abundant in most well-aerated soils, iron has low biological activity because it forms highly insoluble ferric compounds at neutral pH levels. Iron is important in many physiological and biochemical pathways in plants. It is required for a

wide range of biological functions because it is a component of many vital enzymes, such as cytochromes of the electron transport chain. Iron is involved in the synthesis of chlorophyll in plants and is required for the maintenance of chloroplast structure and function. **Rout and Sahoo, 2015.**

Materials and Methods

In order to study the two micronutrients with foliar spray, **Iron and Silicon** were taken. The experiment was conducted during *Zaid* 2022, at Crop Research Farm, Naini Agricultural Institute, SHUATS, Prayagraj. The experimental site of the study is geographically located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level (MSL). The soil of the experimental field constituting a part of central Gangetic alluvium is neutral and deep. Pre-sowing soil samples were taken from a depth of 15 cm with the help of an auger. The composite samples were used for the chemical and mechanical analysis. The treatments consist of basal application of nitrogen (40, 50, 60 Kg/ha) and foliar spray of iron (0.2, 0.4, 0.6%). The experiment was layout in randomized block design with ten treatments each replicated thrice and control i.e., recommended N, P and K (50:30:20 Kg/ha). The plots were prepared with dimension of 3m × 3m and seeds were

sown with a spacing of 30cm × 10cm. At 14 DAS plants were thinned to maintain appropriate plant density. Weeds were controlled manually at 12, 21 DAS with the help of Khurpito minimize crop weed competition. Growth characteristics namely plant height (cm), dry weight (g), number of tillers/hill, were recorded. Irrigations were given uniformly and regularly to all plots as per requirement so as to prevent the crop from water stress at any stage. The crop was harvested at right physiological maturity stage and their post-harvest observations such as panicle length (cm), grains/panicle, grain yield (t/ha), straw yield (t/ha) were recorded. The data recorded for different characteristics were subjected to statistical analysis by adopting the method of analysis of variance (ANOVA) as described by Gomez (1984). The crop was harvested separately from each plot taking 1.0 m² area i.e., 80 DAS. Thereafter, the produce from net plot was tied in bundles separately and then tagged. The tagged bundles were allowed for sun drying in field and after drying on the threshing floor, the weight of bundles was recorded for obtaining biological yield. Threshing of foxtail millet was done manually by beating with stick and then seeds were separated by winnowing.

Results and Discussions

Growth parameters

Table.1 Pertaining the details of effect of nitrogen and iron on growth parameters of foxtail millet.

Plant height (cm)

At harvest, higher plant height (101.57 cm) was recorded significantly in the treatment no.9 [Nat60Kg/ha+0.6%Iron]. However, treatment no.8 [Nat60Kg/ha+0.6%Iron] (100.65 cm) was found to be statistically at **far** with treatment no. 9.

Nitrogen application might have resulted in cell elongation and increased chlorophyll content and net photosynthetic rate which attributed to the increase in plant height. Similar findings were reported by **Venkataramon et al., 2000**.

The **increased** in the availability of iron to plant might have stimulated the metabolic and enzymatic activities resulting in increase in the growth of the crop. Similar findings were also reported by **Yadav et al., 2013**.

Dry weight (g)

At harvest, maximum plant dry weight (14.83 g) was recorded significantly in the treatment no.9 [Nat60Kg/ha+0.6%Iron]. However, treatment no.8 [Nat60Kg/ha+0.4%Iron] (14.62 g) was found to be statistically at **far** with treatment no. 9.

Nitrogen is the main component of the protoplasm which **is involves** in metabolic processes like photosynthesis, cell division and cell elongation. Which

have contributed to increase in the dry matter accumulation. Similar findings are reported by **Rekha and Prasad 2020**.

Number of tillers/hill

At harvest, more tillers/hill (8.87) was recorded significantly in the treatment no.9 [Nat60Kg/ha+0.6%Iron]. However, treatment no.8 [Nat60Kg/ha+0.4%Iron] (8.80) was found to be statistically at **far** with treatment no. 9.

Nitrogen involves in the development of strong cell walls and consequently, straw which might **be resulted** into increased tillering. Similar findings were reported by **Ayub et al., 2007**.

Iron **involves in** chlorophyll synthesis, grain formation and dry matter production, which ultimately lead to final yield characters such as number of effective tillers per plant. Similar findings were reported by **Gupta et al., 2002**.

Yield attributes

Table.2 Pertaining the details of effect of nitrogen and iron on yield attributes of foxtail millet.

Panicle length (cm)

Panicle length was significantly influenced by the application of nitrogen and iron. At harvest, the data recorded highest panicle length (18.61 cm) in treatment no.9 [Nat60Kg/ha+0.6%Iron]. However, treatment no.8

[Nat60Kg/ha+0.4%Iron] (18.19 cm) was statistically at par with treatment no.9.

Application of nitrogen fertilizer at right time improves rate of photosynthesis, root growth and development and elongation of structural tissues such as stalk in cereals. Similar findings were reported by **Pandey and Sinha 2010**.

Grains/panicle

The data of grains/panicle was significantly influenced by the application of nitrogen and iron. At harvest, the data recorded more grains/panicle (1389.30) in treatment no.9 [Nat60Kg/ha+0.6%Iron]. However, treatment no.8 [Nat60Kg/ha+0.4%Iron] (1337.97) was statistically at par with treatment no.9.

Higher number of grains/panicle **might due to the application of nitrogen** which increases the fertility of flowers and increase in leaf area and duration and resulted into increase in supplying assimilates for the sink. Similar findings were reported by **Mousavi et al. 2012**.

Grain yield (q/ha)

The data of grain yield was significantly influenced by nitrogen and iron. At harvest, the data recorded higher grain yield (17.96 q/ha) in treatment no.9 [Nat60Kg/ha+0.6%Iron]. However, treatment no.8 [Nat60Kg/ha+0.4%Iron] (17.65 q/ha) was statistically at par with treatment no.9.

Nitrogen being a major nutrient that increases in grain yield as well as improved root growth and development owing to all physicochemical processes impacted. Similar findings were reported by **Ayub et al. 2009**.

Iron plays a major role in biosynthesis of IAA and especially due to its role in initiation of primordial reproductive part and partitioning of photosynthetic products towards them which promotes the yield. Similar result was also observed by **Rao et al. 2019**.

Straw yield (q/ha)

The data of stover yield was significantly influenced by nitrogen and iron. At harvest, the data recorded higher stover yield (25.32 q/ha) in treatment no.9 [Nat60Kg/ha+0.6%Iron]. However, treatment no.8 [Nat60Kg/ha+0.4%Iron] (24.61 q/ha) was statistically at par with treatment no.9.

More uptake of nutrients by nitrogen probably improved growth and development of crops, resulting into improved fodder yield. Similar findings were reported by **Tiwana et al. 2004**.

Conclusion

By the above study we suggest that higher plant height (101.57 cm), maximum dry weight (14.83 g), more number of tillers/hill (8.87) was recorded in treatment no.9 with application of Nitrogen 60 Kg/ha with combination of Iron 0.6%. Maximum

yield attributes viz., panicle length (18.61 cm), grains/ panicle (1389.30), grain yield (1.79 t/ha), straw yield (2.53 t/ha) were recorded in treatment no.9 with the application of Nitrogen 60 Kg/ha and Iron 0.6%.

References

- Ayub, M., Nadeem, M.A., Tanveer, A., Tahir, M. and Khan, R.M.A. (2007). Interactive effect of different nitrogen levels and seeding rates on fodder yield and quality of pearl millet. *Pakistan Journal of Agricultural Science*, **44**(4): 592-596.
- Ayub, M., Nadeem, M. A., Tahir, M., Ibrahim, M., and Aslam, M. N. (2009). Effect of nitrogen application and harvesting intervals on forage yield and quality of pearl millet (*Pennisetum americanum* L.). *Pak. J. Life Soc. Sci.* **7**: 185-189.
- Gomez, K.A. and Gomez, A.A. (1984). Statistical procedures for agricultural research. John Wiley & Sons.
- Gupta, P. K., Sharma, N. N., Acharaya, H. K., Gupta, S. K. and Mali, G. S. (2002). Response of mungbean to zinc and iron on Vertisols of South-Western Plains of Rajasthan. National Symposium on Arid Legumes for Food Security and Promotion Trade, October, 2002. Sponsored by Indian Arid Legumes Society, CAZRI, Jodhpur.
- Mousavi, S. G. R., Seghatoleslami, M. J., & Arefi, R. (2012). Effect of N fertilization and plant density on yield and yield components of grain sorghum under climatic conditions of Sistan, Iran. *Plant Ecophysiology*, **4**, 141-149.
- Munirathnam, P., Sambasiva Reddy, A. and Sawadhkar, S.M. (2006) Evaluation of foxtail millet varieties under low fertility conditions. *Agric. Sci. Digest*, **26** (3): 197 - 199.
- Pandey, S.N and Sinha, B. K., (2010): Plant physiology. (4th Edition). Vikas Publishing House, New Delhi, India.
- Rao, V., Yadav, B. J. K., S, Jeeterwal, R. C., Response of pearl millet (*Pennisetum glaucum* L.) To Integrated nitrogen management. International. *Journal of Current Microbiol Applied Sciences* 2019; **8** (2):429-437.
- Rekha, R. and Prasad, S. K., (2020) Effect of Farm Yard Manure and Nitrogen on Growth and Yield of Pearl Millet under Custard Apple Based Agri-Horti

System. *International Journal of Current Microbiology and Applied Sciences* (2020) Special Issue-11: 1794-1802.

Rout, G. R. and Sahoo, S. (2015). Role of iron in plant growth and metabolism. *Reviews in Agricultural Science*, 3:1-24.

Tiwana, M.S., Puri, K.P., Tiwana, U.S. and Singh, A. 2004. Forage production potential of napier bajra hybrid varieties under different nitrogen levels. *Forage Res.* **30**(2): 83-85.

Venkataramon, N. S., Kempuchetty, N. and Mohadass, S. 2000. Photosynthesis and productivity of direct sow rice under semi dry ecosystem. *J. Ecobiol.* 12: 69-72.

Yadav, G. S., Shivay, Y. S., Kumar, D. and Babu, S. 2013. Enhancing iron density and uptake in grain and straw of aerobic rice through mulching and rhizo-foliar fertilization of iron. *African Journal of Agricultural Research*, **8**(xx): 5447-5454

Table 1.: Effect of Nitrogen and Iron on growth attributes of Foxtail millet.

Sl No.	Treatment combination	Plant height (cm)	Dry weight (g)	No. of tillers/hill	CGR (g/m ² /day)	RGR (g/g/day)
1.	N 40Kg/ha + 0.2% Iron	90.80	12.51	7.20	8.39	0.024
2.	N 40Kg/ha + 0.4% Iron	91.47	12.65	7.33	7.92	0.023
3.	N 40Kg/ha + 0.6% Iron	92.64	12.98	7.33	8.28	0.023
4.	N 50Kg/ha + 0.2% Iron	93.90	13.25	7.47	8.11	0.022
5.	N 50Kg/ha + 0.4% Iron	94.83	13.60	8.13	7.54	0.019
6.	N 50Kg/ha + 0.6% Iron	97.03	13.85	8.33	7.88	0.020
7.	N 60Kg/ha + 0.2% Iron	99.15	14.35	8.40	8.28	0.020
8.	N 60Kg/ha + 0.4% Iron	100.65	14.62	8.80	8.16	0.020
9.	N 60Kg/ha + 0.6% Iron	101.57	14.83	8.87	7.96	0.018
10.	Control (RDF-50:30:20 NPK Kg/ha)	91.20	13.19	7.27	9.16	0.026
	F Test	S	S	S	NS	NS
	SEm (±)	0.35	0.07	0.08	0.30	0.094
	CD (p=0.05)	1.04	0.21	0.23	-	-

Table 2.: Effect of Nitrogen and Iron on yield attributes and yield of Foxtail millet.

Sl No.	Treatment combination	Yield attributes and Yield					
		Panicle length (cm)	Grains/panicle (no.)	Test weight (g)	Grain yield (t/ha)	Straw yield (t/ha)	Harvest index (%)
1.	N 40Kg/ha + 0.2% Iron	16.19	1220.00	3.00	1.43	2.14	40.19
2.	N 40Kg/ha + 0.4% Iron	16.32	1223.80	3.19	1.47	2.17	40.36
3.	N 40Kg/ha + 0.6% Iron	16.45	1240.90	3.22	1.48	2.20	40.31
4.	N 50Kg/ha + 0.2% Iron	16.74	1255.27	3.23	1.50	2.21	40.13
5.	N 50Kg/ha + 0.4% Iron	17.23	1276.33	3.32	1.54	2.27	40.48
6.	N 50Kg/ha + 0.6% Iron	17.51	1299.57	3.33	1.61	2.30	41.14
7.	N 60Kg/ha + 0.2% Iron	17.83	1316.73	3.39	1.69	2.36	41.78
8.	N 60Kg/ha + 0.4% Iron	18.19	1337.97	3.46	1.76	2.46	41.75
9.	N 60Kg/ha + 0.6% Iron	18.61	1389.30	3.50	1.79	2.53	41.50
10.	Control (RDF-50:30:20 NPK Kg/ha)	16.34	1222.47	3.21	1.46	2.16	40.40
	F Test	S	S	NS	S	S	NS
	SEm (\pm)	0.20	17.54	0.10	0.24	0.41	0.59
	CD (p=0.05)	0.60	52.12	-	0.71	1.23	-