

Original Research Article

An Error Analysis of Implicit Finite Difference Method with Mamadu-Njoseh Basis Functions for Time Fractional Telegraph Equation

Abstract

In this paper, we proposed and analyzed the error estimate of an implicit finite difference method with Mamadu-Njoseh as basis functions for time fractional telegraph equation. To enhance the efficacy of the method we first transform the Caputo type fractional derivative into Riemann-Liouville derivatives. The error analysis of the method is stated and proven. Also, the optimal results for scalars unknown in L_∞ norm were derived for the two-dimensional case. Numerical illustrations are presented to test the reliability of the analytical and computed results. The resulting numerical evidence shows that the proposed method convergences more rapidly than the standard finite difference method. MAPLE 18 is used for all mathematical procedures in this paper.

Keywords: Riemann–Lionville derivatives, Quadrature formula, Orthogonal collocation method, Mamadu-Njoseh polynomials, Sobolev space, Finite difference method.

AMS Subject Classification 2010: 65L60, 26A3, 33D52

1. Introduction

The telegraph equation (otherwise known as the transmission line model) is a coupled partial differential equation that models the flow of voltage and current on a transmission line in time and distance. The equation was designed by Oliver Heaviside in 1876 in the course of developing the transmission line model. The equation has revolved, over the years with direct applications to transmission lines involving all frequencies, such as telephone lines, radio frequency, telegraph wires, power line and wire radio antenna [1].

In view of Wang *et. al.* [2], a typical time fractional telegraph equation is given as

$${}_0^C D_t^\beta u(x, t) + u_t(x, t) - u_{xx}(x, t) = g(x, t), x \in [0, T], t > 0, \quad (1.1)$$

with the initial conditions

$$\left. \begin{aligned} u(x, 0) &= u_0(x) \\ u_t(x, 0) &= u_1(x) \end{aligned} \right\} x \in [0, T], \quad (1.2)$$

and boundary conditions

$$\left. \begin{aligned} u(0, t) &= 0 \\ u(T, t) &= 0 \end{aligned} \right\} t > 0, x \in [0, T] \quad (1.3)$$

where $1 < \alpha < 2$ and $g(x, t)$ is the source term.

In recent literature, there exist some numerical techniques for solving partial differential equations (PDEs), which include the spectral method [3], finite volume method [4], finite difference method [5], mixed finite element method [6], finite element method [7] and $H^1 -$

Galerkin mixed finite element method [2]. Yazdani *et. al.* [8] studied the numerical solution of space fractional advection-diffusion equation adopting the finite volume-element method. The work expressed the fractional derivative composition in the Grunwald-Letnikov form. The convergence and stability of the method was also studied which resulted to the conclusion that full discretization is possible and stable in as much as the mesh graded size is sufficiently small. In like fashion, Hao *et. al.* [9] considered the Galerkin finite element method (GFEM) for the solution of two-sided one-dimensional diffusion equation with variable coefficients. They reformulated the governing problem into a low-ordered term that is fractional by mere introduction of an extra parameter. It was argued that the GFEM is far superior to that of the Petro-Galerkin method in the sense that the GFEM can easily be extended to three-dimensional variable coefficients. However, the stability and convergence analysis of this method was not treated. Superconvergence of the finite element method as applied to time-fractional diffusion equation (TFDE) governed by a time-space diffusivity was studied by An [10]. Weak singularity of the model problem was studied at $t = 0$. Also, the fully discrete scheme on a bounded mesh, and fully discrete conforming finite element method was investigated. The author in conclusion remarked that superconvergence is achievable if temporary mesh pints are set at $r \geq (2 - \alpha)/\alpha$, where r is graded mesh size and $0.5 < \alpha < 1$. Liu *et. al.*, [6] also studied the numerical solution of time-fractional partial differential equations. The mixed element method (MEM) was adopted as the numerical solver of the problem. The work of [6] is quiet fascinating in the sense that the Caputo fractional derivative was discretized in time via the two-step method (otherwise, finite difference method), and spatial direction was discretized using the mixed finite element method. There are few works existing in literature on the numerical methods of the telegraph equation. For instance, Wang *et. al.* [2] applied the H^1 – Galerkin mixed finite element method (H^1 -GMFEM) for the solution of time fractional telegraph equation. In line with the approach of Liu *et. al.* [6], the authors also fully discretized in time the Caputo fractional derivative using the finite difference method, and discretized in space using the H^1 -GMFEM. For more on this, see Wei *et. al.* [11], and Zhao and Li [7].

Mamadu-Njoseh polynomials (MNP) are orthogonal polynomials developed by Njoseh and Mamadu [12] for seeking the approximate solution of many linear and nonlinear problems in the field of applied mathematics. The polynomials were constructed in the interval [-1,1] with respect to the weight function $w(x) = 1 + x^2$. These polynomials so far has contributed so much in seeking the approximate solutions of integral equations, boundary value problems, singular initial value problems in ordinary differential equation, integro-differential equation, delay differential equations. However, these polynomials are implemented through an appropriate numerical scheme as basis functions. For instance, the MNP were adopted by Ahmed and Singh [13] for the solution of integral equation via the Galerkin method. In like manner, Al-Humedi and KadhimMunaty [14] studied comparatively the MNP alongside Chebyshev and Laguerre polynomials for the solution of first kind intergral equation by the spectra petro-Galerkin method. Montazer *et. al.* [15] studied the MNP alongside non-uniform Haar wavelets for the numerical treatment of linear Volterra integral equations. Problems involving fractional order have equally solved applying these polynomials as seen in the literature Xie [16]. Njoseh and Mamadu [17] applied the MNP as trial functions for the solution of fifth order boundary value problems via the power series approximation method. These polynomials were also used by Mamadu and Njoseh [18] for the solution of Votterra integral equation via the Galerkin Method. Mamadu and Njoseh [19] considered the Mamadu-Njoseh polynomials in orthogonal collocation methods for the solution of integro-differential equations. Ogeh and Njoseh [20] constructed a modified variational iteration method for the solution of fifth and sixth order

boundary value problems adopting the Mamadu-Njoseh polynomials as trial functions. In like manner, Njoseh and Musa [21] adopted these polynomials for the solution of pantograph-type delay differential equation in a variational iteration approach. Also, Mamadu and Ojarikre [22] proposed a reconstructed Elzaki transform method (RETM) for the solution of delay differential equation using Mamadu-Njoseh polynomials as basis functions. A perturbation by decomposition technique was considered by Mamadu and Tsetimi [23] adopting the MNP as basis functions for the solution of singular initial value problems.

This paper will centre on Mamadu-Njoseh polynomials as basis functions in a discretization scheme. Thus, a novel finite difference method with Mamadu-Njoseh basis functions for the time-fractional telegraph equation (1.1) will be proposed. An optimal error analysis for scalar unknowns in L_∞ -norm will be established for the equation (1.2).

2. Preliminaries

Let equation (1.1) can be transformed into a fractional differential equation with the Riemann – Lionville derivatives [24]

$${}^R D_t^\beta [u - u_0](x, t) + u_t(x, t) - u_{xx}(x, t) = g(x, t) \quad (1.4)$$

By definition, ${}^R D_t^\beta (u_0) = \frac{d}{dt} \frac{1}{\Gamma(1-\beta)} \int_0^t (t-s)^{-\beta} u_0 ds = \frac{u_0}{\Gamma(1-\beta)} \frac{d}{dt} \left(\frac{1}{1-\beta} t^{1-\beta} \right) = \frac{u_0}{\Gamma(1-\beta)} t^{-\beta}$

Thus,

$${}^R D_t^\beta u(x, t) = \frac{1}{\Gamma(-\beta)} \int_0^t (t-s)^{-\beta-1} u(s) ds \quad (1.5)$$

Let $[0, 1]$ be partitioned as $0 = t_0 < t_1 < \dots < t_n = 1$. By using $nt_j = j, j = 1, 2, \dots, n$, we approximate (1.4) in time step as

$${}^R D_t^\beta u(x, t) = \frac{1}{\Gamma(-\beta)} \int_0^{t_j} (t_j - s)^{\beta+1} u(s) ds.$$

Let $t_j = t_j w + s$, we obtain,

$${}^R D_t^\beta u(x, t_j) = \frac{t_j^{-\beta}}{\Gamma(-\beta)} \int_0^1 \frac{u(t_j - t_j w) - u(0)}{w^{\beta+1}} dw = \frac{t_j^{-\beta}}{\Gamma(-\beta)} \int_0^1 F(w) w^{-\beta-1} dw, \quad (1.6)$$

where $F(w) = u(t_j - t_j w) - u(0)$.

Using the quadrature formula [24], the integral is replaced with $t_j = n/j, n = 0$, for each j , to obtain

$${}^R D_t^\beta u(x, t) = \frac{1}{\Gamma(-\beta)} \left[\sum_{r=0}^j \beta_{rj} u(t_j - t_r) + G_j(g) \right],$$

where

$$\|G_j(g)\| \leq k_j^{\beta-2} \sup_{0 \leq t \leq T} \|u''(t_j - t_j t)\|.$$

Thus,

$$\begin{aligned}
{}^R D_t^\beta u(x, t_j) &= \frac{\Delta t^{-\beta}}{r(2-\beta)} \sum_{r=0}^j (-\beta)(1-\beta)j^{-\beta} \beta_{rj} u(t_j - t_r) + \frac{t_j^{-\beta}}{\Gamma(-\beta)} G_j(g) \\
&= \Delta t^{-\beta} \sum_{r=0}^j \frac{(-\beta)(1-\beta)j^{-\beta} \beta_{rj}}{r(2-\beta)} u(t_j - t_r) + \frac{t_j^{-\beta}}{\Gamma(-\beta)} G_j(g) \\
&= \Delta t^{-\beta} \sum_{r=0}^j w_{rj} u(t_j - t_r) + \frac{t_j^{-\beta}}{\Gamma(-\beta)} G_j(g),
\end{aligned}$$

where

$$\Gamma(2-\beta)w_{rj} = (-\beta + \beta^2)j^{-\beta} \beta_{rj},$$

where w_{rj} and β_{rj} satisfies [24]

$$\begin{aligned}
w_{rj} &= \frac{1}{\Gamma(2-\beta)} \begin{cases} 1, & r = 0 \\ -2r^{1-\beta} + (r-1)^{1-\beta} + (r+1)^{1-\beta}, & r = 1, 2, \dots, j \\ -(\beta-1)r^{-\beta} + (r-1)^{1-\beta} - r^{1-\beta}, & r = j \end{cases} \\
\beta_{rj} &= \frac{1}{\beta(1-\beta)j^{-\beta}} \begin{cases} -1, & r = 0 \\ -2r^{1-\beta} - (r-1)^{1-\beta} - (r+1)^{1-\beta}, & r = 1, 2, \dots, j \\ (\beta-1)r^{-\beta} - (r-1)^{1-\beta} + r^{1-\beta}, & r = j \end{cases}
\end{aligned}$$

3. Discretization in Time

Let consider the finite difference method [25] of (1.4) at $t = t_j$, we get

$${}^R D_t^\beta [u(x, t) - u_0(x, t)]|_{t=t_j} + u_t(x, t_j) = g(x, t_j) + u_{xx}(x, t_j) \quad (1.7)$$

$$\Rightarrow \Delta t^{-\beta} \sum_{r=0}^j w_{rj} [u(t_j - t_r) - u(0)] + \frac{t_j^{-\beta}}{\Gamma(-\beta)} G_j(g) + u_t(x, t_j) = g(x, t_j) + u_{xx}(x, t_j),$$

or

$${}^R D_t^\beta [u(x, t) - u_0(x, t)]|_{t=t_j} = \Delta t^{-\beta} \sum_{r=0}^j w_{rj} [u(t_j - t_r) - u(0)] + \frac{t_j^{-\beta}}{\Gamma(-\beta)} G_j(g). \quad (1.8)$$

Denote $u_j \approx u(x, t_j)$, we obtain,

$$\Delta t^{-\beta} \sum_{r=0}^j w_{rj} [u_{j-r} - u_0] + \frac{u_j - u_{j-1}}{\Delta t} = G(x, t_j) + u_{xx}(x, t_j).$$

Let $r = 0$, we obtain,

$$\begin{aligned}
&\Delta t^{-\beta} (w_{0j} u_j - w_{0j} u_0) + \Delta t^{-\beta} \sum_{r=1}^j w_{rj} [u_j - r] + \Delta t^{-\beta} \sum_{r=0}^j w_{rj} u_0 + \frac{u_j - u_{j-1}}{\Delta t} = G(x, t_j) + \frac{u_{j+1} - 2u_j + u_{j-1}}{\Delta t^2} \\
\Rightarrow &(u_j - u_0)w_{0j} + \sum_{r=1}^j w_{rj} u_{j-r} + \sum_{r=0}^j w_{rj} u_0 + \Delta t^{-\beta} \left[\frac{u_j - u_{j-1}}{\Delta t} - \frac{u_{j+1} + 2u_j - u_{j-1}}{\Delta x^2} \right] = g(x, t_j).
\end{aligned}$$

But from, we find [30]

$$\sum_{r=0}^j w_{rj} = \frac{-\beta(1-\beta)j^{-\beta}}{(2-\beta)} \left(\frac{-1}{\beta}\right) = j^{-\beta} h_{\beta},$$

where $h_{\beta} = \frac{1}{\Gamma(1-\beta)}$.

Thus, the implicit difference formula for (1.4) is given as

$$(u_j - u_0)w_{0,j} + \sum_{r=1}^j w_{rj}u_{j-r} + j^{-\beta}h_{\beta}u_0 + \Delta t^{\beta} \left(\frac{u_j - u_{j-1}}{\Delta t} - \frac{u_{j+1} + 2u_j - u_{j-1}}{\Delta x^2} \right) = g(x, t_j). \quad (1.9)$$

3.1 Discretization in Time with Mamadu-Njoseh Polynomials

Suppose that an approximation to (1.4) is declined by

$$u_j(x, t) \cong u(x, t) = \sum_{r=0}^j a_r(t) \varphi_r(x), \quad (1.10)$$

where $a_r(t)$, $r = 0(1)j$, are unknown constants to be estimated, and $\varphi_r(x)$, $r = 0(1)j$, are Mamadu – Njoreh basis functions.

Using (1.10) on (1.9), we obtain a new implicit implicit scheme given by

$$\begin{aligned} & \sum_{r=0}^j a_r(t) \varphi_r(x) w_{0,j} + \frac{j^{-\beta}}{\Gamma(1-\beta)} a_0(t) + \sum_{r=1}^j w_{rj} \left(\sum_{r=0}^{j-r} a_r(t) \varphi_r(x) \right) + \\ & \frac{\Delta t^{\beta}}{\Delta t} \sum_{r=0}^j a_r(t) \varphi_r(x) - \frac{\Delta t^{\beta}}{\Delta t} \sum_{r=0}^{j-1} a_r(t) \varphi_r(x) - \frac{\Delta t^{\beta}}{\Delta t} \sum_{r=0}^j a_r(t) \varphi_r(x) + \frac{2\Delta t^{\beta}}{\Delta x^2} \sum_{r=0}^j a_r(t) \varphi_r(x) + \\ & \frac{\Delta t^{\beta}}{\Delta x^2} \sum_{r=0}^{j-1} a_r(t) \varphi_r(x) = g(x, t) \end{aligned} \quad (1.11)$$

Equation (1.11) is collocated orthogonally ([26]-[28]) for any $j > r$ to obtain a system of $(j + 1)$ linear equations which can be solve for the unknowns $a_r(t)$, $r = 0(1)j$, via a suitable mathematical software with estimate $\beta, \Delta t, \Delta t^2, w_{rj}$ clearly given. The approximate solution to (1.4) is obtain by substituting the known estimate into (1.10).

4. Main Results

In this section, we carry out a precise analysis of error estimate on the derived implicit formula (1.11).

Define $\beta = u_t(x, t) - u_{xx}(x, t)$, $\Delta(\beta) = H_0^1 \cap H^2 = \{u: u(0) = u(T) = 0, u', u'' \in L_2(0, T)\}$, then the equation (1.4) can be reformulated in abstract sense as

$${}^R D_t^{\beta} [u(t) - u_0(t)]|_{t=t_j} + Bu(t) = g(t), 0 \leq x \leq T, t > 0, \quad (1.12)$$

Using compound quadratic formula on (1.12), we have

$$\frac{t^{-\beta}}{\Gamma(-\beta)} \sum_{r=0}^j \beta_{rj} [u(t_j - t_r) - u_0] + G_j(g) + Bu(t_j) = g(t_j) \quad (1.13)$$

Evaluating (1.13) at $r = 0$ yields

$$[\beta_{0j} + t_j^{\beta} \Gamma(-\beta) \beta] u(t_j) = t_j^{\beta} \Gamma(-\beta) g_j - \sum_{r=1}^j \beta_{rj} (t_j - t_r) + \sum_{r=0}^j \beta_{rj} u_0 - G_j(g) \quad (1.14)$$

$$\Rightarrow [\beta_{0j} + t_j^\beta \Gamma(-\beta) \beta] u(t_j) = t_j^\beta \Gamma(-\beta) - \sum_{r=0}^j \beta_{rj} u_{jor} + \sum_{r=0}^j \beta_{rj} u_0, \quad (1.15)$$

where

$$u_j \approx u(t_j) = \sum_{r=0}^j a_r(t) u_r(x).$$

Let $e_j = u - u_j$ be the error function, where u is the analytic solution and u_j , then (1.15) can also reformulated in reference to (1.11) as

$$[\beta_{0j} + t_j^\beta \Gamma(-\beta) \left(\frac{u_j - u_{j-1}}{\Delta t} - \frac{u_{j+1} + 2u_j - u_{j-1}}{\Delta x^2} \right)] u_j = t_j^\beta \Gamma(-\beta) - \sum_{r=0}^j \beta_{rj} u_{j-r} + \sum_{r=0}^j \beta_{rj} u_0 \quad (1.16)$$

Lemma 1 [24]: For $\beta \in (0,1)$, let $\{h_j\}_{j=1}^\infty$ be defined by $h_1 = 1$, then

$$1 \leq h_j \leq \frac{\sin \pi \beta}{\pi \beta (1-\beta)} j^\beta, j = 1(2)\infty.$$

Lemma 2: Suppose $p_0 = 1, p_j = \beta(1-\beta)j^{-\beta} \sum_{r=1}^j \beta_{rj} p_{j-r}, j = 1(2)\infty$, then, $p_j \leq 1$.

Theorem 1: Let U_j and u_j denotes the solution of (1.4) with the prescribed conditions, then

$$e_j \leq k \Delta t^{2-\beta} + e_0, \text{ where } e_0 = \|u_0 - u\|.$$

Proof: Let subtract (1.15) from (1.13) to obtain the error equation as

$$\begin{aligned} (\beta_{0j} + t_j^\beta \Gamma(-\beta) B) e_j &= -(\sum_{r=0}^j \beta_{rj} e_{j-r} + G_j(g)) \\ \Rightarrow e_j &= (-\beta_{0j} - t_j^\beta \Gamma(-\beta) B)^{-1} \left(\sum_{r=0}^j \beta_{rj} e_{j-r} + G_j(g) \right) \end{aligned} \quad (1.17)$$

By definition of L_∞ norm, we obtain

$$\|e_j\| \leq \|(-\beta_{0j} - t_j^\beta \Gamma(-\beta) B)^{-1}\| (\sum_{r=0}^j \beta_{rj} \|e_{j-r}\| + \|G_j(g)\|) \quad (1.18)$$

where B is symmetric and positive definite operator [29]. Now for any function $f(x)$, we have via spectral method that $\|f(B)\| = \sup_{\tau > 0} |f(\tau)|$.

Hence, using the definition of β_{rj} and (1.18), we have

$$\begin{aligned} \|(-\beta_{0j} - t_j^\beta \Gamma(-\beta) B)^{-1}\| &= \left\| \left(\frac{1}{\beta(1-\beta)j^{-\beta}} - t_j^\beta \Gamma(-\beta) B \right)^{-1} \right\| \\ &= \left\| \beta(1-\beta)j^{-\beta} (1 - \beta(1-\beta)j^{-\beta} t_j^\beta \Gamma(-\beta) B)^{-1} \right\| \\ &= \beta(1-\beta)j^{-\beta} \sup_{\tau > 0} (1 - \beta(1-\beta)j^{-\beta} t_j^\beta \Gamma(-\beta) \tau)^{-1}. \end{aligned} \quad (1.19)$$

By definition of gamma function, it is obvious $\Gamma(-\beta) > 0$. This implies

$$\sup_{\tau > 0} (1 - \beta(1-\beta)j^{-\beta} t_j^\beta \Gamma(-\beta) \tau)^{-1} \leq 1.$$

Therefore,

$$\left\| (-\beta_{0j} - t_j^\beta \Gamma(-\beta) B)^{-1} \right\| \leq \beta(1 - \beta) j^{-\beta}.$$

Thus, (1.18) becomes

$$\|e_j\| \leq \beta(1 - \beta) j^{-\beta} \left[\sum_{r=0}^j \beta_{rj} \|e_{j-r}\| + K j^\beta m^{-2} \sup_{0 \leq t \leq T} \|u_{tt}''(t_j - t_r)\| \right], \Delta t = 1, t_m = m. \quad (1.20)$$

Equation (1.20) can be reformulated into the form

$$\|e_j\| \leq b + \beta(1 - \beta) j^{-\beta} \sum_{r=0}^j \beta_{rj} \|e_{j-r}\|, \quad (1.21)$$

where $b = \beta(1 - \beta) j^{-\beta} K m^{-2} \|u''\|_{L_\infty}$.

Let $j = n$, then (1.21) becomes

$$\begin{aligned} \|e_n\| &\leq b + \beta(1 - \beta) n^{-\beta} \left[\sum_{r=1}^{n-1} \beta_{rn} \|e_{n-r}\| + \beta_{nn} \|e_0\| \right] \\ &\leq b + \beta(1 - \beta) n^{-\beta} \left[\sum_{r=1}^{n-1} \beta_{rn} (b h_{n-r} + p_{n-r} \|e_0\|) + \beta_{nn} \|e_0\| \right] \\ &= a h_n + p_n \|e_0\| \end{aligned}$$

where

$$\begin{aligned} h_n &= 1 + \beta(1 - \beta) n^{-\beta} \sum_{r=1}^{n-1} \beta_{rn} h_{n-r}, \quad n = 2, 3, \dots, \\ p_n &= \beta(1 - \beta) n^{-\beta} \sum_{r=1}^n \beta_{rn} p_{n-r}, \quad n = 1(2)\beta, p_0 = 1. \end{aligned}$$

Now using lemma 1 and lemma 2, we have from (1.21),

$$\begin{aligned} \|e_j\| &\leq b h_j + p_j \|e_0\| \leq \beta(1 - \beta) K m^{-2} \|u''\|_{L_\infty} \cdot h_j + p_j \|e_0\| \\ &\leq \beta(1 - \beta) K m^{-2} \|u''\|_{L_\infty} \frac{\sin \pi \beta}{\pi \beta (1 - \beta)} j^\beta + \|e_0\| \\ &\leq 1 \\ &\leq k \Delta t^{2-\beta} + \|e_0\|. \end{aligned}$$

5. Numerical Illustration

To test the reliability and accuracy of the proposed method, we consider the example below:

$${}_0 D_t^\beta u(x, t) + u_t(x, t) - u_{xx}(x, t) = 2(x^2 - x) t \left(\frac{\Gamma(3-\beta) + t^{1-\beta}}{\Gamma(3-\beta)} \right) - 2t^2, \quad x \in [0, 1], \quad t \in (0, 1], \quad (1.22)$$

with the initial conditions

$$\left. \begin{aligned} u(0, 0) &= 0 \\ u_t(x, 0) &= 0 \end{aligned} \right\}'$$

and boundary conditions

$$\left. \begin{aligned} u(0, t) &= 0 \\ u(1, t) &= 0 \end{aligned} \right\} t > 0.$$

Applying the scheme (1.11) on (1.22) at $j = 3$ with parameters $\beta = 1.5$, $\Delta x = 1/64$, $\Delta t = 1/1000$ at $t = 0$, and values of w_{rj} , $r = 0(1)3$, estimated as $w_{0,3} = 1$, $w_{1,3} = -0.7294368868$, $w_{2,3} = 0.09204003089$ and $w_{3,3} = 0.1817856084$. Results are presented **Table 1** and **Figure 1** using MAPLE 18.

Table 1: Maximum error at $\beta = 1.5$

j	L_∞ Error (Proposed method)	L_∞ Error (Li and Cao, 2012)
20	5.6445E-008	9.877022E-003
40	3.5198E-006	3.477002E-003
80	1.5441E-005	1.232302E-003
160	6.3327E-005	4.249358E-004




Figure 1: Comparison of computed solutions and exact solutions

Similarly at $j = 3$ with parameters $\beta = 1.8$, $\Delta x = 1/32$, $\Delta t = 1/1000$ at $t = 0$, and values of w_{rj} , $r = 0(1)3$, estimated as $w_{0,3} = 1$, $w_{1,3} = -0.3105422252$, $w_{2,3} = 0.05806019726$ and $w_{3,3} = 0.06480727693$. Results are presented in **Table 2** and **Figure 2** using MAPLE 18.

Table 2: Maximum error at $\beta = 1.5$

j	L_∞ Error (Proposed method)	L_∞ Error (Li and Cao, 2012)
20	6.6376E-004	1.1484477E-002
40	1.4188E-006	5.148878E-003
80	2.4280E-007	1.998394E-003
160	3.9604E-008	8.980387E-004




Figure 2: Comparison of computed solutions and exact solutions

6. Discussion of Results

Numerical evidences to test the reliability and accuracy of the proposed method are presented in tables and figures. **Tables 1** and **2** shows the maximum error of the proposed method at $j = 20, 40, 80,$ and 160 . In comparison with the finite difference method in [30] show the superiority of the proposed method with maximum errors of order 10^{-8} and 10^{-8} , respectively. Results are also presented in graphs showing the comparison of results as shown in the **figures 1** and **2**.

5. Conclusion

In this paper, we have successively proposed an implicit finite difference method with Mamadu-Njoseh polynomials as basis functions for the time fractional telegraph equation. Numerical illustration of the proposed method showed convergence and accuracy than the standard finite difference method. The optimal error analysis of the proposed method was investigated in L_∞ norm for two dimensional case. The result showed that the method is of order $(2 - \beta)$.

References

- [1] V. K., Srivastava, M. K., Awasthi, R. K., Chaurasia, and M. Tamsir, the telegraph equation and its solution by reduced differential transform method, *Modelling and Simulation in Engineering*, Article ID 746351, 6 pages, 2013. <http://dx.doi.org/10.1155/2013/746351>
- [2] J. Wang, M. Zhao, M. Zhao, Y. Liu, Y. and H. Li., Numerical analysis of an H^1 -galerkin mixed finite element method for time fractional telegraph equation, *The Scientific World Journal*, Article ID 371413, 14 pages, 2014 <http://dx.doi.org/10.1155/2014/371413>.
- [3] Y. M. Lin, and C.J. Xu, Finite difference/spectral approximations for the time-fractional diffusion equation, *Journal of Computational Physics*, 225(2): 1533–1552, 2007.
- [4] Q. Q. Yang, I. Turner, T. Moroney and F.W. Liu, A finite volume scheme with preconditioned Lanczos method for two-dimensional space-fractional reaction-diffusion equations, *Applied Mathematical Modelling*, 2014.
- [5] A. Atangana and D. Baleanu, D., Numerical solution of a kind of fractional parabolic equations via two difference schemes, *Abstract and Applied Analysis*, Article ID 828764, 8 pages, 2013.
- [6] Y. Liu, H. Li, W. Gao, S. He, S. and Z.C. Fang, A new mixed finite element method for a class of time-fractional partial differential equations, *The Scientific World Journal*, Vol 2014, Article ID 141467, 8 pages, 2014.
- [7] Z. G. Zhao and C. P. Li, Fractional difference/finite element approximations for the time-space fractional telegraph equation, *Applied Mathematics and Computation*, 219(6): 2975–2988, 2012.
- [8] A. Yazdani., N. Mojahed, A. Babaei, and V. Cendon, Using finite volume-element method for solving space fractional advection-dispersion equation, *Progr. Fract. Differ. Appl* 6(1):55-66, 2020.
- [9] Z. Hao, M. Park, G. Lin and Z. Cai, Finite element method for two-sided fractional differential equations with variable coefficients: Galerkin approach, *Journal of Scientific Computing*, 2018. <https://doi.org/10.1007/s10915-018-0869-5>
- [10] N. An, Superconvergence of a finite element method for the time-fractional diffusion equation with a time-space dependent diffusivity, *An advances in Differential Equations*, 2020. <https://doi.org/10.1186/s13662-020-02976-4>
- [11] L. Wei, H. Dai, D. Zhang, and Z. Si, Fully discrete local discontinuous Galerkin method for solving the fractional telegraph equation, *Calcolo*, 51(1):175–192, 2014.
- [12] E.J. Mamadu and Njoseh, I. N, Power series variational iteration method for the numerical treatment of fifth order boundary value problems, *PK ISSN 0022- 2941; CODEN JNSMAC* 57:10-18, 2017.
- [13] N. Ahmadi and B. Singh, Numerical solution of integral equation using galerkin method with hermite, chebyshev and orthogonal polynomials, *Journal of Science and Arts Year* 20, 1(50): 35-42, 2020.
- [14] H.O. Al-Humedi and A. KadhimMunaty, The spectral petrov-galerkin method for solving integral equations fo the first kind, *Turkish Journal of computer and Mathematics Education*, 12(13): 7856- 7865, 2021.
- [15] M. Montazer1, R. Ezzati1 and M. Fallahpour, Numerical solution of linear volterra integral equations using non-uniform haar wavelets, *Kragujevac Journal of Mathematics*, 47(4): 599–612, 2023. https://imi.pmf.kg.ac.rs/kjm/pdf/accepted-finished/e297c72509880e6c79bfd0f8b082e4ba_2536_12162020_043209/kjm_47_4-9.pdf

- [16] J.Q. Xie, [Numerical Computation of Fractional Constitutive Model for Potential and Non-stationary Heat Transfer Problems](https://elibrary.ru/item.asp?id=42679289), 2018. <https://elibrary.ru/item.asp?id=42679289>
- [17] I.N. Njoseh and E.J. Mamadu, Numerical solutions of fifth order boundary value problems using Mamadu-Njoseh polynomials. *Science World Journals*, 11(4): 21-24, 2016.
- [18] E.J. Mamadu and I.N. Njoseh, Numerical solutions of Volterra equations using Galerkin method with certain orthogonal polynomials. *Journal of Applied Mathematics and Physics*, **4**:376-382, 2016. <http://dx.doi.org/10.4236/jamp2016.420454>.
- [19] E.J. Mamadu and I.N. Njoseh, Certain orthogonal polynomials in orthogonal collocation methods of solving integro-differential equations (fides). *Transactions of the Nigeria Association of Mathematical Physics*, **2**:59-64, 2016.
- [20] K.O. Ogeh and I.N. Njoseh, Modified variational iteration method for solving boundary value problems using Mamadu-Njoseh polynomials. *International Journal of Engineering and future technology*. 16(4): 24 – 36, 2019.
- [21] I.N. Njoseh and A. Musa, Numerical solution of Pantograph-type delay differential equation using variational iteration method with Mamadu-Njoseh polynomials. *International Journal of Engineering and future technology*. 10(3), 2019.
- [22] E.J. Mamadu and H.I. Ojarikre, Reconstructed Elzaki transform method for delay differential equations with Mamadu-Njoseh Polynomials. *Journals of Mathematics and System Science* **9**:41-45, 2019.
- [23] J. Tsetimi, and E.J. Mamadu, Perturbation by decomposition: A new approach to singular initial value problems with Mamadu-Njoseh basis functions, *Journal of Mathematics and System Science*, 2020. <http://dx.doi.org/10.17265/2159-5291/2020.01.003>
- [24] K. Diethelm, Generalized compound quadrature formulae for finite integral, *IMA Journal of Numerical Analysis*, 17:479-493, 1997.
- [25] K. Diethelm, *Fractional differential equations, theory and numerical treatment*, TU Braunschweig, Braunschweig, 2003.
- [26] B.A. Finlayson, Orthogonal collocation on finite elements - progress and potential. *Mathematics and Computer in simulation*, **22**(1):11-17, 1980.
- [27] L. Fox and I.B. Parker, *Chebyshev Polynomials in Numerical Analysis*. Oxford University Press, Oxford, London, 1968.
- [28] C. Lanczos, Trigonometric interpolation of empirical analytical functions . *J. Math. Phys.*, **17**: 123-199, 1938.
- [29] P.G. Ciale, *Finite element method for elliptic problems*, North-Holland Publishing Company, Amsterdam, 1987.
- [30] C. Li and J. Cao, A finite difference method for time-fractional telegraph equation, *IEEE*, 2012.