

Original Research Article

EFFECT OF MULCHING ON GROWTH, YIELD AND QUALITY ATTRIBUTES OF (*Fragaria x ananassa*)” IN DEHRADUN VALLEY OF UTTARAKHAND

Abstract

Present experimental research was conducted during the year 2020-21 at Horticulture Research block, School of Agriculture Sciences, SGR University, Dehradun, Uttarakhand to study the “Effect of Mulching on Growth, Yield and Quality Attributes of Strawberry (*Fragaria X ananassa*) in Dehradun Valley of Uttarakhand”. The field experiment was laid out in randomized block design with three replications and three treatments. The treatments were comprised with different mulching materials viz. control as without mulching (T1), Straw mulching (T2) and Polythene mulching (T3). The planting of cultivar “Chandler” was done on first week of November, 2020 and observations were recorded at 30 DAT, 60 DAT, 90 DAT, 120 DAT, 150 DAT and at final harvest. Significant results were obtained for different attributes of growth, yield, quality as well as in terms of economics under the treatment of black polythene mulching (T3). The results showed that the Number of Flowers per Plant (24.08), Plant spread (48.06 cm), Fruit length (4.22 cm), Fruit diameter (3.80 cm), Fruit texture (5.9 lb), T.S.S. (8.92 °Brix) and Total juice content (84.05 %) However, the control (T1) reported maximum Titrable acidity (0.92 %).

Keywords- Chandler, Economics, Mulching, Randomized, Runners.

Introduction

Strawberry (*Fragaria x ananassa*) belongs to the family Rosaceae, is one of the important soft fruit of the world and has resulted from a cross between two wild strawberries: *Fragaria virginiana* (Meadow strawberry) and *Fragaria chiloensis* (Chilean strawberry). It is a herbaceous perennial plant which can be successfully cultivated in wide range of Agroclimatic conditions from subtropics to temperate climate.

It is a rich source of Vitamin A (60 IU/100 g of edible fruit), Vitamin C (30-120 mg/100g of edible portion), Vitamin B1, niacin, Proteins and minerals like Phosphorous, Potassium, Calcium and Iron. The presence of ellagic acid, which prevents cancer and occurrence of heart diseases and the abundance of anthocyanins have made it a more valuable fruit. The heart-shaped silhouette of the strawberry is the first clue that this fruit is good for health. These potent little packages protect your heart, lower your blood pressure, and guard against cancer. Packed with Vitamins, Fibre, and particularly high levels of antioxidants known as polyphenols helps in Wrinkle-prevention, strawberries are a sodium-free, fat-free, cholesterol-free, low-calorie food. They are among the top 20 fruits in antioxidant capacity and are a good source of manganese and potassium (S.K. Singh 2019).

The plastic materials used as mulch are poly vinyl chloride or polyethylene films. Owing to its greater permeability to long wave radiation it can increase temperature around the plants during night in winter. Hence, polyethylene film mulch is preferred as mulching material for crop production (Nautiyal *et al.*, 2017a). Now a day application of black plastic mulch film is becoming popular and very good results have been achieved particularly in rain fed agriculture (Nautiyal *et al.*, 2017b). The black polyethylene mulch also checks all types of weeds in addition to soil moisture conservation therefore, black plastic mulch is more beneficial (McCann *et al.*, 2007).

Mulching reduces soil temperature in summer and raises it in winter. It prevents the extremes of temperatures. During summer, mulching conserves the soil moisture due to

reduced evaporation. The cooling effect of soil promotes root development. In general, the effect of mulching on the temperature regime of the soil varies according to the capacity of the mulching material to reflect and transmit solar energy. Mulches results in greater water content and lower the evaporation (Semwal *et al.*, 2022). White mulches decrease soil temperature while clear plastic mulches increase soil temperature. The soil temperature can be higher up to 70C under clear mulch compared to bare soil. At night, condensation on the underside of the mulch absorbs the long wave radiation emitted by the soil thereby slowing cooling of the soil (Lamont, W. J. 2005). The ability of clear mulches to produce soil temperatures high enough to control weeds, plant pathogens and nematodes forms the basis for the soil solarization process.

Materials and Method

The field experiment was conducted on 2020-21 in the Horticulture Research Block of School of Agricultural Sciences, SGRR University, Dehradun, Uttarakhand. The experiment was laid out in Randomized block design with three replications. Each replication consisted of three treatments. All the treatments were randomized separately in each replication.

List 1 : Treatment Details

Treatment Symbol	Treatment Combination
T ₁	Control (without any mulching)
T ₂	Paddy straw mulching
T ₃	Black polythene mulching

Result and Discussion

The findings of the present investigation were recorded and are thoroughly discussed below:

Number of Flowers

Bothstraw and black polythene mulches recorded higher number of flowers as compared to control. However, the maximum number of flowers per plant (62.71) was obtained under black polythene mulching (T₃). This might be attributed to more number of flowers produced by vigorous plants under black polythene mulch, which is in conformity with the findings of Nagalakshmi *et al.*, (2002).

Plant Spread (cm)

Plant spread was found significantly maximum (48.06) under black polythene mulching (T₃) as compared to rest of the treatments. This might be due to the presence of adequate moisture in the soil which is vital for plant growth not only because plant needs

water for their physiological processes but also for nutrient solubility and their availability in soil solution (**Kumar and Dey, 2011**).

Fruit Length (cm) and Fruit Diameter (cm)

Fruit length (4.22 cm) and fruit diameter (3.80 cm) under black polythene mulch (T₃) was found significantly higher in comparison to all other treatments. The larger fruit size under black polythene mulch was attributed to more plant growth and development under micro-climatic condition resulting in better nutrient uptake. Increase in fruit size due to mulching had also been reported by **Nagalakshmi et al., (2002)**.

T.S.S. (°Brix) and Total Juice Content (%)

The data depicts that the application of black polythene mulch recorded maximum T.S.S. (8.92°B) and total juice content (84.05 %). Higher fruit quality is related to weed free environment, higher moisture conservation and maximum nutrient uptake under black polythene mulch treatment. These results are in conformity with the findings of (**Mathad and Jholgiker, 2005**) and **Singh et al., (2007)**.

Titration Acidity (%)

The titration acidity was high in fruits harvested from the control treatment (T₃) (0.92%). The variation in titration acidity of strawberry fruits harvested from the plants applied with different mulches might be due to variation in the translocation of photo assimilates from source to sink. These results were in conformity with the findings of **Verma et al., (2005)** and **Das et al., (2010)** in guava.

The candidate manuscript does not have a robust scientific discussion. I suggest the authors incorporate the suggested paragraphs, in this way it would improve the scientific quality of the manuscript.

Formatted: Justified

Mulching can have a significant impact on plant growth by modifying soil temperature and moisture. In tropical climates, mulching can help to reduce soil temperature and conserve soil moisture, which can benefit plant growth (Rey et al. 2022). Several studies have demonstrated that mulching can increase the growth of strawberry plants by improving their root development, nutrient uptake, and photosynthetic efficiency (Moor et al. 2004).

Formatted: Font: (Default) Arial, 10 pt, Not Bold, English (United States)

Mulching can also have a positive effect on crop yield. By modifying soil temperature and moisture, mulching can improve the availability of nutrients and water to the plant, which can enhance fruit development and increase yield. Several studies have reported that mulching can significantly increase the yield of strawberry crops in tropical regions. For example, a study conducted in Brazil found that plastic mulching increased strawberry yield by 20% compared to non-mulched plants (Coelho et al. 2008).

Formatted: English (United States)

Mulching can also affect the quality attributes of strawberry fruits. By modifying soil moisture and nutrient availability, mulching can influence the concentration of sugars, acids, and other compounds that contribute to fruit flavor and nutritional value. Several studies have reported that mulching can improve the quality attributes of strawberry fruits, including their sugar content, antioxidant activity, and vitamin C concentration.

The effect of mulching on strawberry growth, yield, and quality attributes is closely related to environmental factors such as climate and soil. In tropical regions, where high temperatures and low soil moisture can limit crop growth and yield (Hernández and Olivares, 2019), mulching can help to create a more favorable growing environment for plants (Orlando and Hernandez, 2020). The choice of mulching material and application rate should be based on the specific environmental conditions of the region. For example, in regions with high rainfall, organic mulches may be more appropriate to prevent soil erosion (Olivares et al. 2011; Olivares et al. 2015) and maintain soil structure (Olivares et al. 2022; Orlando et al. 2020), while in regions with low rainfall, inorganic mulches such as plastic film may be more effective in conserving soil moisture (Olivares et al. 2016; Campos and López, 2019; Hernandez et al. 2018).

In conclusion, mulching can have a significant positive effect on the growth, yield, and quality attributes of *Fragaria x ananassa* in tropical crops. Mulching can modify soil temperature and moisture, which can benefit plant growth and development. The effect of mulching on strawberry crops is closely related to environmental factors such as climate and soil, and the choice of mulching material and application rate should be tailored to the specific environmental conditions of the region. Mulching is a simple and effective agricultural practice that can improve the productivity and sustainability of tropical agriculture.

Conclusion

On the basis of present experimental research on “Effect of Mulching on Growth, Yield and Quality Attributes of Strawberry (*Fragaria X ananassa*) in Dehradun Valley of Uttarakhand” in cultivar “Chandler”, it can be concluded that among different mulching materials, black polythene mulch (T₃) was found to be most effective for influencing various growth parameters such as- number of flowers /plant, plant spread (cm), yield parameters like fruit length (cm), fruit diameter (cm), fruit texture (lb) and quality parameters like T.S.S. (°Brix)

and total juice content (%). However, the titrable acidity (%) was found maximum in control (T₁)

Comment [A1]: this is a result, not a conclusion

References

suggest adding recent references which address the issue in question, suggested citations are for genuine scientific reasons that emphasize the current topic of study in context.

[Campos, B., and López, M. 2019. Normalized Difference Vegetation Index \(NDVI\) applied to the agricultural indigenous territory of Kashaama, Venezuela. UNED Research Journal. 11\(2\): 112-121. <https://doi.org/10.22458/urj.v11i2.2299>](#)

Formatted: English (United States)

[Coelho, M. V., Palma, F. R., and Cafe-Filho, A. C. 2008. Management of strawberry anthracnose by choice of irrigation system, mulching material and host resistance. International Journal of Pest Management, 54\(4\), 347-354.](#)

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States)

Formatted: English (United States)

Das, B.C, Maji, S. and Mulieh, S.R. 2010. Response of soil covers on guava cv.L-49. Journal of Crop and Weed. 6 (2): 10-14.

Formatted: Font: (Default) Times New Roman, 12 pt, English (United States)

Formatted: Font: (Default) Times New Roman, 12 pt

[Hernández, R. and Olivares, B. 2019. Ecoterritorial sectorization for the sustainable agricultural production of potato \(*Solanum tuberosum* L.\) in Carabobo, Venezuela. Agricultural Science and Technology. 20\(2\): 339-354. \[https://doi.org/10.21930/rcta.vol20_num2_art:1462\]\(https://doi.org/10.21930/rcta.vol20_num2_art:1462\)](#)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Formatted: Font: (Default) Times New Roman, 12 pt

Kher, R, Baba, J. A, Bakshi, P. and Wali, V.K. 2010. Effect of planting time and mulching material on quality of strawberry. Journal of Research. SKUAST-Jammu 9 (1): 54-62.

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic

Kumar S, Dey P 2011. Effect of different mulches and irrigation methods on root growth, nutrient uptake, water use efficiency and yield of strawberry. Sci Hort. 127(3):318-324.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: Italic

Lamont, W.J. 2005 Plastics: Modifying the microclimate for the production of vegetable crops. Hort Technology 15:477-48.

Mathad JC, Jhologiker P 2005 Effect of synthetic and organic mulches in improving growth, yield and quality of strawberry under subtropical ecosystem. Acta Hort 696:56-61.

McCann, I. Kee, E. Adkins, J. Ernest, E. and Ernest, J. 2007. Effect of irrigation rate on yield of drip-irrigated seedless watermelon in humid region. Science Horticulture, 113:155-161.

[Moor, U., Karp, K., and Pöldma, P. \(2004\). Effect of mulching and fertilization on the quality of strawberries.](#)

Nagalakshmi S, Palanisamy D, Eswaran S, Sreenarayanan VV 2002. Influence of plastic mulching in chilli yield and economics. South Indian Hort. 50(1/6):262-265.

Nautiyal P, Sachan, V K, Papnai, G, Tiwari, R K, Manisha. 2017. Improving growth, yield and profitability in apple through mulching in rainfed condition in hilly region of Uttarakhand. J Krishi Vigyan 6 (1): 101-104.

Nautiyal P, Papnai, G, Manisha. 2017. Impact of adoption of mulching technology in higher apple production in Uttarakhand. Progressive Horticulture, 49 (2): 204-206.

Semwal A, Singh S, Nautiyal P, Pandey S. 2022. Effect of Mulching on Growth, Yield and Economics of Strawberry (*Fragaria* × *ananassa*) under Subtropical Conditions of

Uttarakhand. International Journal of Agriculture, Environment and Biotechnology, 15(004): 819-825

[Olivares B, Verbist, K., Lobo, D., Vargas, R., Silva, O. 2011. Evaluation of the USLE model to estimate water erosion in an Alfisol. Journal of Soil Science and Plant Nutrition of Chile. 11 \(2\):71-84. <http://dx.doi.org/10.4067/S0718-95162011000200007>](#)

Formatted: Portuguese (Brazil)

[Olivares, B.O., Calero, J., Rey, J.C., Lobo, D., Landa, B.B., Gómez, J. A. 2022. Correlation of banana productivity levels and soil morphological properties using regularized optimal scaling regression. Catena, 208: 105718. <https://doi.org/10.1016/j.catena.2021.105718>](#)

Formatted: Spanish (Spain, International)

[Olivares, B., Lobo, D., Verbist, K. 2015. Aplicación del modelo USLE en parcelas de erosión bajo prácticas de conservación de suelos y aguas en San Pedro de Melipilla, Chile. Revista Ciencia e Ingeniería. 36 \(1\):3-10.](#)

Formatted: Hyperlink, Font: (Default) +Body, 11 pt, Font color: Auto, English (United States)

Field Code Changed

[Olivares, B. 2016. Descripción del manejo de suelos en sistemas de producción agrícola del sector Hamaca de Anzoátegui, Venezuela. La Granja: Revista de Ciencias de la Vida. 23\(1\): 14-24. <https://n9.cl/ycp08>](#)

[Hernández, R; Olivares, B., Arias, A; Molina, J.C., Pereira, Y. 2018. Identificación de zonas agroclimáticas potenciales para producción de cebolla \(*Allium cepa* L.\) en Carabobo, Venezuela. Journal of the Selva Andina Biosphere. 6 \(2\): 70-82. \[http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf\]\(http://www.scielo.org.bo/pdf/jsab/v6n2/v6n2_a03.pdf\)](#)

Formatted: Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: Spanish (Spain, International)

Formatted: Font: Italic, Spanish (Spain, International Sort)

Formatted: Spanish (Spain, International)

Formatted: English (United States)

[Orlando, B., Hernández, R. 2020. Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. Tropical and Subtropical Agroecosystems, 23\(2\):1-12. <https://n9.cl/zeedh>](#)

Formatted: Portuguese (Brazil)

[Orlando, B.; Hernandez, R.; Arias, A; Molina, J.C., Pereira, Y. 2020. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. Idesia. 38\(2\):95-102. <http://dx.doi.org/10.4067/S0718-34292020000200095>](#)

Formatted: Portuguese (Brazil)

Formatted: No underline, Portuguese (Brazil)

Formatted: Portuguese (Brazil)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

[Rey, J.C., Olivares, B.O., Perichi, G., Lobo, D. 2022. Relationship of Microbial Activity with Soil Properties in Banana Plantations in Venezuela. Sustainability 14, 13531. <https://doi.org/10.3390/su142013531>](#)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto, Portuguese (Brazil)

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto, Portuguese (Brazil)

S.K. Singh 2019. Handbook of Horticulture Volume 1, Indian council of Agricultural Research, New Delhi. ISBN: 978-81-7164-187-1.

Formatted: No underline, Font color: Auto, Portuguese (Brazil)

Singh R, Sharma RR, Goyal RK 2007. Interactive effects of planting time and mulching on 'Chandler' strawberry Sci. Hort. 111:344-351.

Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto, Portuguese (Brazil)

Singh, A. K, Singh, S, Rao, V.V. A, and More, T.A. 2010. Efficiency of organic mulches on soil properties, earthworm population, growth and yield of aonla cv. NA7 in semi-arid ecosystem. Indian Journal of Horticulture. 67: 124-128.

Formatted: Font: (Default) Times New Roman, 12 pt

Verma, M.L, Bhardwaj, S.P, Thakur, B.C.S and Bhandari, A. R. 2005. Nutritional and mulching studies in apple. Indian Journal of Horticulture. 62 (4): 332-335.

Table 1: GROWTH PARAMETERS

Treatment	Number of Flowers/ Plant	Plant Spread (cm)
T ₁	15.30	29.624
T ₂	16.97	30.946
T ₃	24.08	39.994
C.D. (P=0.05)	5.91	1.97
SE(m) ±	1.78	0.59
SE(d) ±	2.52	0.84
C.V.	19.25	3.96






Table 2 : YIELD PARAMETERS

Treatment	Fruit Length (cm)	Fruit Diameter (cm)	Fruit Texture (lb.)
T ₁	3.16	2.68	5.1
T ₂	3.40	3.49	5.5
T ₃	4.22	3.8	5.9
C.D. (P=0.05)	0.24	0.19	N/A
SE(m) ±	0.06	0.04	0.32
SE(d) ±	0.08	0.06	0.46
C.V.	2.89	2.53	10.36










Table 3 : QUALITY PARAMETERS

Treatment	T.S.S. (°Brix)	Juice Content (%)	Titration Acidity (%)
T ₁	6.88	67.86	0.92
T ₂	7.54	72.14	0.81
T ₃	8.92	84.05	0.72
C.D. (P=0.05)	0.80	0.22	0.085
SE(m) ±	0.20	0.05	0.021
SE(d) ±	0.28	0.07	0.030
C.V.	4.46	0.12	4.471





UNDER PEER REVIEW