

Yield and Economics of Pea (*Pisum Sativum* L.) as Influence by Spacing and Nipping in Manipur

ABSTRACT

Aims: To study the yield and economics of pea (*Pisum Sativum* L.) as influence by spacing and nipping in Manipur

Study design: Factorial randomized block design (FRBD)

Place and Duration of Study: Experimental field of College of Agriculture, Central Agricultural University, Imphal, Manipur, *rabi* season of 2013-14 and 2014-15.

Methodology: The treatment combinations involving two factors were S₁N₁, S₁N₂, S₂N₁, S₂N₂, S₃N₁, S₃N₂, S₄N₁ and S₄N₂ (Spacing i.e., S₁-20X15cm, S₂-20X20cm, S₃-30X10cm and S₄-30X15cm and nipping i.e., N₁-No nipping and N₂-Nipping).

Results: The pooled results of two years of study revealed that the yield differed significantly at a 5% probability level with different treatment combinations. The number of branches at harvest was significantly affected by both spacing and nipping with maximum branches in treatment combination S₄N₂ resulting in highest number of pods observed in S₄N₂ with 11.02 (pooled) and the lowest in S₁N₁. Spacing and nipping treatment had no significant influence on the pod length. The maximum stover yield was recorded in S₁N₂ with 27.35 q/ha and minimum in S₄N₁ with 22.02 q/ha. The highest seed yield of 16.05 q/ha was recorded in S₄N₂ with highest net return (₹ 2,05392/ha) and cost-benefit ratio (1:4.69). The present study suggests that the spacing of 30x15cm with nipping was found to be the best amongst the treatment combinations in regards to yield and higher feasible net returns.

Keywords: Pea; yield; economics; spacing; nipping.

1. INTRODUCTION

Pea (*Pisum Sativum* L.) is an important pulse crop of India grown during *rabi* season. Peas are cultivated for the fresh green seeds, tender green pods, dried seeds and foliage (Duke)[3]. Like any legume crop, pea is an integral component of sustainable agriculture due to its soil enriching and conditioning properties. Pulses are important sources of dietary protein for a majority of the Indian population. They are highly nutritive and contain high proportion of digestible protein, carbohydrates, minerals and vitamins. It is the richest source of protein, i.e., 18-35 % in dry seeds. It contains a high proportion of minerals and a good source of vitamins A, B and C (Khvostova)[5].

In Manipur, since time immemorial, as a tradition nipping of peas was done at initial stage of growth which improve the yield and provides additional income by selling nipped shoots as leafy vegetable. Some of the local cultivar of peas in Manipur are *Makuchabi*, *Makhyatmubi*, *Ningtekpi*, etc. They are grown mainly during winter. However, in the Hilly tracts of the state where cool weather prevails throughout the year, pea is traditionally cultivated all the year round for a sustainable crop production. It is essential to provide optimum plant population density per unit area by adjusting the spacing levels in pea crop unlike in normal spacing the plants grown in closer spacing exhibited more vertical growth but give less yield and poor-quality seeds for need of sufficient space, light, nutrient and moisture due to heavier plant population pressure. Among various components of agro-techniques, identification of optimum plant population per unit area plays a vital role in maximizing pea seed yield as well as its quality (Saimbhi and Gill) [10]. Among several seed production approaches, apical bud nipping is being commonly practiced in several crops to increase the seed yield and quality. The studies on influence of nipping on seed yield and economics are absolutely very scanty. Keeping in view of the above aspects, the present investigation has been undertaken to find out the yield and economics of pea (*Pisum Sativum* L.) as influence by spacing and nipping.

2. MATERIAL AND METHODS

A field experiment was conducted at College of Agriculture, Central Agricultural University, Imphal during the *rabi* seasons of 2013-14 and 2014-15, situated at about 24°46' N latitude and 93°54' E longitude and an altitude of about 790 metre above MSL. The soil was clay in texture, acidic soil pH, medium in organic carbon (0.58%) content, medium in available nitrogen (280.53 kg/ha), medium in available P₂O₅ (18.45 kg/ha) and medium in K₂O. During the period of growing season, minimum temperature was found to be 4.56°C in the month of January for the first year (2013-14) and 6.2°C in the month of December for the second year (2014-15) and a maximum temperature of 27.97°C (2013-14) and 28.5°C (2014-15) in the month of March. A maximum rainfall of 47.5mm (2013-14) and 213.4 (2014-15) was recorded in the month of April but there was no rain during the month of November to January. A recommended dose of fertilizers 20:40:20 in the form urea, single super phosphate and muriate of potash were applied as basal. The experiment was laid out in FRBD with 8 treatments and 3 replications. The treatment combinations involving two factors i.e., 1) Spacing: S₁-20x15cm, S₂-20x20cm, S₃-30x10cm and S₄-30x15cm and 2) Nipping: N₁-No nipping and N₂-Nipping were S₁N₁, S₁N₂, S₂N₁, S₂N₂, S₃N₁, S₃N₂, S₄N₁ and S₄N₂. The selected cultivar for the experiment was Pea (*Pisum sativum* L. subsp. hortense, local cultivar-*Makhyatmubi*). Five plants from each plot were selected randomly excluding border rows plants and were tagged properly for taking observations.

3. RESULTS AND DISCUSSION

3.1 Influence of spacing on yield and attributes

In crops like pea (*Pisum sativum*), seed yield is a function of plant growth, number of productive branches, number of pods per plant, seeds per pod, seed yield, etc. Effect of spacing and nipping influenced the yield and yield attributes of pea and are presented in Table 1. On an average, plants of wider spacing 30x15cm (S₄) recorded significantly more number of pods per plant of 9.58 (pooled data) while minimum number (6.50) were recorded in the spacing 20x15cm (S₁). The increase in the number of pods per plant in wider row spacing may be due to vigorous plant growth producing more number of productive branches which resulted in higher pod yield. On the other hand, in closer row spacing, the plant growth was limited resulting in less number of pods per plant. These results are in agreement with the findings of Monalisa *et al.* [8] in pigeon pea, Shajid *et al.* [11] in pea, Sharma *et al.* [12] in pigeon pea and Subedi *et al.* [14] in pea. They found the highest number of pods per plant in wider row spacing as compared to closer spacing.

The data on seed yield revealed that significantly maximum yield was recorded in the spacing 30x15cm (S₄) with 15.09 q/ha while minimum seed yield (10.01) was recorded in the spacing 20x15cm (S₁). The superior values of yield and its component per plant noticed under wider spacing may be attributed to be better growth and development of plants under less plant population density and it resulted into better source to sink relationship due to availability of balanced and adequate nutrients and better light, space and moisture unlike in narrow spacing. These results are in conformity with those of Ali *et al.* [1] in pea, Monalisa *et al.* [8] in pigeon pea and Subedi *et al.* [14] in pea.

Stover yield in pea plant differ significantly due to spacing. It was observed that the maximum stover yield was given by the spacing 20x15cm (S₁) with 26.93 q/ha while the minimum was recorded in the spacing of 30x15cm (S₄) with 23.09q/ha. The higher plant density per unit area and the higher plant height attributed to maximum stover yield in closer spacing among all the spacing treatments. The similar increase in stover yield with narrow spacing was observed by Derya [2] in peas, Luikham *et al.* [6] in broad bean and Taipodia and Nabam [15] in cowpea.

Table 1. Effect of spacing and nipping on number of pods per plant, pod length, yield and stover yield of pea

	Number of pods/plant	Pod length (cm)	Yield (q/ha)	Stover yield (q/ha)
--	----------------------	-----------------	--------------	---------------------

Treatment	Year 1	Year 2	Pooled	Year 1	Year 2	Pooled	Year 1	Year 2	Pooled	Year 1	Year 2	Pooled
Spacing												
S ₁ (20x15cm)	6.47	6.53	6.50	7.83	7.92	7.88	10.05	9.96	10.01	26.81	27.04	26.93
S ₂ (20x20cm)	7.4	7.78	7.59	7.95	7.9	7.93	11.19	11.32	11.26	25.62	25.7	25.66
S ₃ (30x10cm)	8.02	8.1	8.06	7.98	7.9	7.94	11.96	11.92	11.94	23.63	24.01	23.82
S ₄ (30x15cm)	9.53	9.62	9.58	7.92	7.83	7.88	15.03	15.14	15.09	22.86	23.32	23.09
S.E. d (±)	0.09	0.11	0.06	0.04	0.05	0.19	0.17	0.12	0.15	0.53	0.34	0.25
CD (P=0.05)	0.37	0.34	0.17	NS	NS	NS	0.36	0.24	0.34	1.13	0.72	0.54
Nipping												
N ₁ (Non-nipped)	6.43	6.64	6.54	7.88	7.88	7.88	10.96	11.02	10.99	23.9	24.35	24.13
N ₂ (Nipped)	9.28	9.38	9.33	7.96	7.89	7.93	13.16	13.15	13.16	25.55	25.68	25.62
S.E. d (±)	0.09	0.08	0.04	0.03	0.04	0.2	0.12	0.08	0.1	0.37	0.24	0.17
CD (P=0.05)	0.26	0.24	0.12	NS	NS	NS	0.25	0.17	0.24	0.8	0.51	0.36
Treatment combinations												
S ₁ N ₁	5.20	5.10	5.15	7.80	7.90	7.85	9.04	8.69	8.87	26.16	26.82	26.49
S ₁ N ₂	7.73	7.97	7.85	7.87	7.93	7.90	11.05	11.23	11.14	27.45	27.25	27.35
S ₂ N ₁	5.83	6.37	6.10	7.90	7.87	7.89	10.44	10.63	10.54	24.99	25.06	25.03
S ₂ N ₂	8.97	9.20	9.09	8.00	7.93	7.97	11.93	12.01	11.97	26.25	26.34	26.30
S ₃ N ₁	6.67	6.87	6.77	7.97	8.00	7.99	10.29	10.56	10.43	22.72	23.24	22.98
S ₃ N ₂	9.37	9.33	9.35	8.00	7.80	7.90	13.63	13.27	13.45	24.53	24.78	24.66
S ₄ N ₁	8.03	8.23	8.13	7.87	7.77	7.82	14.04	14.20	14.12	21.75	22.29	22.02
S ₄ N ₂	11.03	11.00	11.02	7.97	7.90	7.94	16.02	16.08	16.05	23.98	24.34	24.16
S.E. d (±)	0.17	0.16	0.08	0.05	0.08	0.09	0.23	0.16	0.21	0.75	0.48	0.35
CD (P=0.05)	0.51	0.49	0.24	NS	NS	NS	0.5	0.34	0.49	1.61	1.03	0.75

3.2 Influence of nipping on yield

The advantage of nipping to increase the number of productive branches, pods per plant and higher yield have been noticed in several field crops. In the present study, the terminal shoot tips of plants of the nipped treatments were nipped at 30 DAS and the resulting influence on the yield parameters and monetary benefits were studied. In the present investigation, nipped plants recorded significantly higher values for almost all the seed yield parameters. The higher number of pods per plant with 9.33, higher seed yield of 13.16 q/ha was recorded in the nipped treatment (N₂) compared to the non-nipped plants (N₁) giving 6.54 number of pods per plant and 10.99 q/ha seed yield. The increase in seed yield and yield attributing parameters noticed with nipping was attributed due to production of more number of productive branches in chickpea by Khan *et al.* [4]. Similar increase in seed yield and yield parameters with nipping were also reported by Reddy [9] in cowpea.

3.3 Interaction effect of spacing and nipping on yield

In the present investigation, the seed yield and almost all the yield parameters are significantly influence by the interaction between spacing and nipping. The maximum seed

yield of 16.05 q/ha was obtained in the treatment combination of spacing 30x15cm (S₄) with nipping (N₂) and minimum (8.87 q/ha) at spacing 20x15cm (S₁) without nipping (N₁) amongst all the treatment combinations. The higher seed yield may be attributed due to more number of branches in the wider spacing and the nipped which encourage vigorous vegetative growth increasing the quantum of photosynthates production and translocation to sink which resulted in increased number of pods per plant. The finding is in agreement with Mazoka *et al.* [7]

The stover yield of pea crop increases with narrow spacing and with nipping. The maximum stover yield of 27.35q/ha (pooled data) were recorded in the treatment combination of spacing 20x15cm (S₁) with nipping (N₂) compared to all other treatment combinations. This may be due to markedly increased in plant density at narrow spacing and the increased in number of branches per plant on account of nipping at that particular spacing. These results are in accordance with the findings of Luikham *et al.* [6]

3.4 Effects of spacing and nipping on economics

Effect of spacing and nipping on economics of pea are presented in Table 2. The treatment combinations involving wider spacing and nipping practice showed higher gross return, net returns and cost-benefit ratio over the narrow spacing without nipped treatments. In the present study, maximum gross return of ₹ 8,480 from shoot yield and ₹2,40,750 from seed yield per hectare, net returns and benefit cost ratio of ₹ 2,05,392 and 1:4.69 respectively were obtained in the treatment combination of spacing 30x15cm (S₄) with nipping (N₂). This may be attributed to the fact that at wider spacing, lower seed rate is required compared to narrow spacing under the same agronomic practices and nipping. This finding is also supported by Singh and Devi [13] in pea.

Table 2. Effect of spacing and nipping on economics of pea

Treatment combinations	Nipped shoot yield (q/ha)	Return from nipped shoot (₹/ha)	Seed yield (q/ha)	Return from seed yield (₹/ha)	Total gross return (₹/ha)	Total cost of cultivation (₹/ha)	Net returns (₹/ha)	Benefit cost ratio
S ₁ N ₁	0	0	8.87	1,33,050	1,33,050	48538	84,512	1.74
S ₁ N ₂	2.08	8320	11.14	1,67,100	1,75,420	49663	1,25,757	2.53
S ₂ N ₁	0	0	10.54	1,58,100	1,58,100	44338	1,13,762	2.57
S ₂ N ₂	2.07	8280	11.97	1,79,550	1,87,830	45238	1,42,592	3.15
S ₃ N ₁	0	0	10.43	1,56,450	1,56,450	48338	1,08,112	2.24
S ₃ N ₂	2.1	8400	13.45	2,01,750	2,10,150	49238	1,60,912	3.27
S ₄ N ₁	0	0	14.12	2,11,800	2,11,800	42938	1,68,862	3.93
S ₄ N ₂	2.12	8480	16.05	2,40,750	2,49,230	43838	2,05,392	4.69

4. CONCLUSION

The result of this study reveals that the number of pods per plants recorded significantly higher at wider spacing with nipping treatment. The maximum numbers of pods and seed yield were recorded in the treatment combination S₄N₂ (30x15cm spacing with nipping) among all the treatment combinations and that of stover yield in the treatment combination S₂N₂ (20x20cm spacing with nipping). Careful study of data from the present investigation suggests that the spacing of 30x15cm with nipping was found to be the best amongst the treatment combinations in regards to better yield and higher feasible net returns.

REFERENCES

1. Ali EH, Abdulla AA, Obaid AK. Effect of plant spacing on growth and yield of two pea cultivars (*Pisum sativum* L.) sown in southern of Iraq. *Basrah J. Agric. Sci.*, 2007;20(2).
2. Derya OY. Impact of plant density on yield and yield components of pea (*Pisum sativum* ssp. *sativum* L.) cultivars. Asian Research Publishing Network (ARPN). *J. Agric. Biol. Sci.*, 2013;2(8):169-174.
3. Duke JA. Hand book of legumes of world economic importance. Plenum Press, New York. 1981;199-265.
4. Khan H, Latif A, Mahmood S, Khan MSS. Effect of nipping at various stages on yield and yield components of chickpea (*Cicer arietinum* L.). *J. Res. Sci.*, 2006; Bahauddin Zakariya University, Multan, Pakistan.
5. Khvostova VV. Genetics and Breeding of Peas: Published by Oxonian Press PVT. Ltd., New Delhi, India. 1983.
6. Luikhram E, Phulchand M, Anal PS. Response of broad bean (*Vicia faba* L.) to row spacing and phosphorus under late sown rainfed condition of Manipur. *Agric. Sci. Digest*, 2009;29(1):54-56.
7. Majoka M, Panghal VPS, Duhan DS. Effect of nipping and plant spacing on seed production of cowpea in Haryana condition. *Forage Res.*, 2021;46(4):343-347.
8. Monalisa S, Wadile SC and Patil HM. Effect of spacing and seedling age on growth and yield of transplanted pigeon pea. *Bulletin of Env., Pharmacology and Life Sci.* 2017;6(5):226-235.
9. Reddy P. Effect of growth retardants and nipping on growth and yield parameters in cowpea (*Vigna unguiculata* L.). M.Sc. (Ag.) thesis, 2005; University of Agricultural Sciences, Dharwad.
10. Saimbhi MS, Gill BS. Effect of phosphorus and row spacing on green pod yield in garden pea. *Vegetable Sci.* 1986;13(1):22-24.
11. Sajid M, Rab A, Jan I, Ahmad I, Ahmad Khan I, Anwar Khan M. Effect of herbicides and row spacing on the growth and yield of pea. *Pakistan J. Weed Sci. Res.*, 2012;18(1).
12. Sharma A, Potdar MP, Pujari BT, Dharmaraj PS. Studies on response of pigeon pea to canopy modification and plant geometry. *Karnataka J. Agric. Sci.*, 2001;16(1):1-3.
13. Singh MS, Devi KS. Profitability of nipping in cultivation of pea (*Pisum sativum*)—an indigenous agro-technique in Manipur. *Indian J. Agronomy*, 2006;51(3):206-208.
14. Subedi PB, Srivastava A, Giri HN, Lohani S. Performance of pea varieties at different plant spacing for seed production at Rampur, Chitwan, Nepal. *Nepalese J. Agric. Sci.*, 2019;18:122-132.
15. Taipodia R, Nabam AT. Impact of time of sowing, spacing and seed rate on potential seed production and fodder quality of cowpea [*Vigna unguiculata* (L.) Walp.]. *J. Agriculture and Veterinary Sci.*, 2013;4(4):61-68.