

**Effect of Bio fertilizers and Phosphorus on Growth and Yield of Pearl millet**  
**(*Pennisetum glaucum* L.)**

**ABSTRACT**

A research trail was conducted at Crop Research Farm in the Department of Agronomy during *kharif* season of 2022 on pearl millet. SHUATS, Prayagraj (U.P). To study the effect of bio fertilizers and Phosphorus on growth and yield of Pearl millet. The treatments consisted of 3 levels of Biofertilizers (20 g/kg *Azotobacter*, 20 g/kg *Azospirillum*, 20 g/kg *Azotobacter*+*Azospirillum*) and phosphorus (35, 40, 45 kg/ha) and a control. The experiment was laid out Randomized Block Design with ten treatments each replicated thrice. The soil in experimental field was sandy loam texture, having alkaline reaction (pH 7.7) with very low organic carbon (0.44%), available higher N (171.48 kg/ha), P (27.0 kg/ha) and higher level of K (291.2 kg/ha). The results showed that application of 20 g/kg *Azotobacter*+*Azospirillum*+ 45 kg/ha Phosphorus was recorded significantly higher plant height (200.13 cm), Plant dry weight (40.33 g/plant), Ear head length (25.25 cm), grains/earhead (2010.9), Grain yield (3.25 t/ha), Stover yield (6.27 t/ha) as compared to other treatments.

**Keywords:** *Azospirillum*, *Azotobacter*, phosphorus, Growth attributes and Yield attributes.

**INTRODUCTION**

Pearl millet (*Pennisetum glaucum* L.) is the staple cereal of arid and semi-arid drier regions of the country. India is the largest Pearl millet growing country contributing 42 percent of production in the world. In India, pearl millet is pre-dominantly cultivated as a rainfed crop in diverse soils, climatic condition and indispensable arid zone. In India pearl millet was cultivated in 7.12 million hectares with 8.06 million tonnes production and productivity of 1132 kg/ha during 2015-16. The major pearl millet producing states in India are Rajasthan, Maharashtra, Gujarat, Uttar Pradesh and Haryana. Therefore, there is need to improve fertility management along with optimum plant density of current hybrids for sustainable production and productivity (Venkata et al., 2001). Bio fertilizers are commonly called microbial inoculants which are capable of mobilizing important nutritional elements in the soil from non-usable to usable form by the crop plants through their biological processes.

Azotobacter and Azospirillum are the two examples of these microorganisms. Mycorrhizal symbiosis increases absorption of some elements such as phosphorus, nitrogen and micronutrients, improves water uptake, produces hormones, reduces damages caused by environmental stress. Improves quality of soil aggregate.

Azospirillum enhance the growth of the plant by influencing the mineral uptake, enhances the dry matter production. Further it also helps in improving the uptake of water and increase the yield of crops. Azotobacter is a N fixing bacteria which improves the crop growth through absorption of N which is fixed near the soil root zone (Goud *et al.*, 2021). *Azospirillum* is a rhizosphere bacterium colonizing the root of the crop plants making use of root exudates and fixes substantial amount of nitrogen and is benefit to plants by mechanisms related to enhancement of plant growth, increase the mineral uptake, increase the dry matter, improve the water absorption and improve the yield. Azotobacter is a free-living nitrogen bacterium which has been reported to fix about 20 kg N/ha in non-legumes (Goud *et al.*, 2021). Phosphorus is widely called as “Bottleneck of world hunger” and an essential element with plays vital role in plant’s growth and development. Adequate phosphorus nutrition enhances many aspects of plant growth development including flowering, fruiting, roots growth and yield components of different crops. P uptake in plants is often constrained by the very low solubility of P in the soil. In agricultural systems, Phosphorus in the harvested crops is removed from the system, resulting in P deprived soils if no P is supplemented as fertilizer (Jain 2005).

Phosphorus is essential for all living organisms. Plants must have phosphorus for normal growth and maturity. Phosphorus plays a role in photosynthesis, respiration, energy, storage and transfer, cell division enlargement and several other processes in plants. A Plant must have phosphorus to complete its normal production cycle. Phosphorus is a vital component of DNA, the genetic memory unit of all living things. It is also component of RNA, the compound that reads the DNA genetic transfer Anonymus, (2016).

## MATERIALS AND METHODS

The experiment was conducted during *kharif* season of 2022. The experiment was conducted in randomized block design and it consist of 10 treatment combinations with

three replications and was laid out variously with different treatments assigned randomly in each replication. The soil in experimental field was sandy loam texture, having alkaline reaction (pH 7.7) with very low organic carbon (0.44%), available higher N (171.48 kg/ha), P (27.0 kg/ha) and higher level of K (291.2 kg/ha). treatment combination were T<sub>1</sub> - 20 g/kg Azotobacter + 35 kg/ha Phosphorus; T<sub>2</sub> - 20 g/kg Azotobacter + 40 kg/ha Phosphorus ; T<sub>3</sub> - 20 g/kg Azotobacter + 45 kg/ha Phosphorus; T<sub>4</sub> - 20 g/kg *Azospirillum* + 35 kg/ha Phosphorus ; T<sub>5</sub> -20 g/kg *Azospirillum* + 40 kg/ha Phosphorus; T<sub>6</sub> - 20 g/kg *Azospirillum* + 45 kg/ha Phosphorus ; T<sub>7</sub> - 20 g/kg Azotobacter + *Azospirillum* + 35 kg/ha Phosphorus ; T<sub>8</sub> - 20 g/kg Azotobacter + *Azospirillum* + 40 kg/ha Phosphorus ; T<sub>9</sub> - 20 g/kg Azotobacter + *Azospirillum* + 45 kg/ha Phosphorus ; T<sub>10</sub> - Control (RDF – 80 – 40 - 40 NPK kg/ha) . The observations were recorded on growth parameters at harvest yield i.e. plant height(cm), dry weight (g/plant), Ear head length (cm), Number of Grains/ear head, Test weight (g), Grain yield (t/ha), Stover yield (t/ha).

### 3. RESULTS AND DISCUSSIONS

#### Growth attributes

##### 3.1 Plant height (cm)

At 80 DAS, significantly highest plant height (200.13 cm) was observed in the treatment with 20 g/kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorus over all the other treatments. However, the treatments with application of 20 g/kg *Azotobacter* + *Azospirillum* + 40 kg/ha Phosphorus (196.40 cm) which were found to be at par with treatment 20 g/kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorus.

The probable reason for increase in plant height due to application of bio fertilizer and phosphorus were bio fertilizer play crucial role in a free-living nitrogen-fixing bacterium termed *Azotobacter* has been reported to fix about 20 kg/ha of nitrogen in non-legumes. The results were found in accordance with **Ramdev et al (2017)**. The significant increase in the height may be due to inoculation of bacterial preparation accelerate plant growth provide biologically fixed nitrogen to the inoculated plant and also stimulate plant growth by excreting plant growth promoting substances like auxins, kinetins, vitamins and gibberellins as similarly observed by **Patidar and Mali (2004)**. Phosphorus plays a pivotal role in extensive root systems efficient utilization of nutrients under conditions of sufficient P application similar results was observed by **Lakhan et al (2017)**

##### 3.1.2 Dry weight (g/plant)

At 80 DAS, treatment with 20 g/kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorus was recorded with significantly maximum dry weight (40.33 g/plant) over all the treatments. However, the treatments with 20 g/kg *Azotobacter* + *Azospirillum* + 40 kg/ha Phosphorus (37.67 g/plant) which were found to be statistically at par with 20 g/kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorus.

The significant increase in a plant dry weight at different stages of growth due to application of biofertilizers and phosphorus was might be due to crucial role of *Azotobacter* can transform elemental nitrogen into Ammonical form ( $\text{NH}_4^+$ ), which plant life uses. In addition, *Azotobacter* has a distinct benefit due to its capacity to synthesize auxins, vitamins, growth factors, and antifungal antibiotics. The roots absorbed the nitrogen that was being fixed by *Azotobacter* in the soil near the root zone (rhizosphere), which may have increased the crop's growth parameters. These findings closely match those of **Kumar et al.,**

(2012) Inoculation of biofertilizers stimulates activation of hormones which helps in shoot and root elongation and high dry matter production, similar results were observed by **Rathore et al. (2006)**

## 3.2 Yield attributes

### 3.2.1 Earhead length (cm)

The application of 20 g/kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorous significantly increased the maximum ear head length (25.25 cm). However, the treatments of 20 g/kg *Azotobacter* + *Azospirillum* + 40 kg/ha Phosphorous (23.87 cm) and 20 g/kg *Azotobacter* + *Azospirillum* + 35 kg/ha Phosphorous (22.55 cm) were statistically at par with 20 g/kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorous.

The yield characteristics were significantly enhanced, and the bio fertilizer and phosphorus application significantly increased the ear head length response. *Azotobacter* that helped roots grow better. Accordingly, better root advancement and energetic plant development, which thus came about to more dry matter lastly better blossoming and ear head improvement, may have been helped by the expanded accessibility of nitrogen. Similar findings were obtained (**Patel et al 2014**).

### 3.2.2 Grains /earhead

The application of 20 g/kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorous produced the highest grains/ear head (2010.9). However, the treatment 20 g/kg *Azotobacter* + *Azospirillum* + 40 kg/ha Phosphorous (1981.0) which were found to be statistically at par with 20 g/kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorous. A significant increase in the number of grains per ear head is the result of an increase in the availability of nitrogen through bio fertilizer inoculation, which also results in an increase in the number of ear heads produced as a result of increased rates of spikelet primordial production; **Marnagar and Dawson (2017)** found results that were comparable.

### 3.2.3 Test weight(g)

There was no significant difference among the treatments. However, highest test weight (6.63 g) was recorded with the treatments 20 g/ kg *Azotobacter* + *Azospirillum* + 35 kg/ha Phosphorous whereas, minimum Test weight (5.47g) was recorded with 20 g/ kg *Azotobacter* + 35 kg/ ha Phosphorus.

### 3.2.4 Grain yield (t/ha)

Significantly highest Grain yield (3.25 t/ha) was recorded with the treatment application of 20 g/ kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorous. However, the treatments with (3.15 t/ha) in 20 g/ kg *Azotobacter* + *Azospirillum* + 40 kg/ha Phosphorous which were found to be statistically at par with 20 g/ kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorous.

Phosphorus recorded significantly higher grain yield, this could mainly be ascribed to the increased availability of the nitrogen to the plants through biological nitrogen fixation in rhizosphere by *Azotobacter* that caused better root development. Thus, the greater availability of nitrogen might have helped in better root proliferation and vigorous plant growth, resulting in more dry matter and ultimately better flowering and ear head development. The increase in yield might be due to the cumulative effect of increased growth and yield attributes noted under this treatment. The results obtained are in close agreement with the findings of **Sushila and Giri (2000)**

### 3.2.5 Stover yield (t/ha)

Significantly highest Stover yield (6.27 t/ha) was recorded with the treatment application of 20 g/ kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorous overall the treatments. However, the treatments with (5.64 t/ha) in 20 g/ kg *Azotobacter* + *Azospirillum* + 40 kg/ha Phosphorous (5.43 t/ha) which were found to be statistically at par with 20 g/ kg *Azotobacter* + *Azospirillum* + 45 kg/ha Phosphorous.

Straw yield is dependent on vegetative growth as use of balanced and optimum use of fertilizer increased plant height, green leaves per hill, and dry matter production, which finally resulted in higher stover yield, similar results were obtained by **Zothanmawii et al (2018)**

### 3.2.6 Harvest index (%)

There was no significant difference among the treatments. However, highest Harvest index (36.94 %) was recorded with the treatments 20 g/ kg Azotobacter + 35 kg/ha Phosphorus whereas, minimum Harvest index (32.55 %) was recorded with Control (RDF 80-40-40 N-P-K kg/ha).

### CONCLUSION

From the results, application of biofertilizers and P can significantly improve the growth and yield of pearl millet. The results of this study support the use of biofertilizers and P as a sustainable agricultural practice for improving soil fertility and crop productivity in tropical regions. It is concluded that with the application of 20 g/kg Azotobacter + Azospirillum and 45 kg/ha Phosphorus (treatment 9) was found more productive (3.25 t/ha) in Pearl millet crop and it can be recommended to farmers after further trials.

### REFERENCES:

- Anonymous, (2016). What's the function of Phosphorus (P) in plants <https://www.greenwaybiotech.com>.
- Goud, B. S. S., Singh, R., & Kan, W. (2021). Effect of seed inoculants and organic manures on growth and yield of organic pearl millet (*Pennisetum glaucum* L.). *Asian J Microbiol Biotechnol Environ Sci*, 23(2), 253-7.
- Jain, N. K., & Dahama, A. K. (2005). Residual effect of phosphorus and zinc on yield, nutrient content and uptake and economics of Pearl millet (*Pennisetum glaucum*)-wheat (*Triticum aestivum*) cropping system. *Indian journal of agricultural science*, 75(5), 281-284.
- Kumar, P., Hooda, R. S., Kumar, S. U. R. E. S. H., & Singh, K. A. R. M. A. L. (2012). Effect of nitrogen levels and biofertilizer strains on dry matter production and attraction index in pearl millet. *Haryana Journal of Agronomy*, 28(1/2), 34-37.
- Lakhan, S., PK Sharma, M Jajoria, P, D., and R Verma. (2017). Effect of Phosphorus and Zinc Application on Growth and Yield Attributes of Pearl millet (*Pennisetum glaucum* L.) under Rainfed Condition. 6(1): 388-391.
- Marnagar, E. and Dawson, J. 2017. Effect of biofertilizers, level of nitrogen and zinc on growth and yield of hybrid maize (*Zeamays* L.). *International Journal of Current Microbiology and Applied Sciences*, 6(9):3614-3622.

- Patidar, M. and Mali, A.L. 2004. Effect of Farmyard manure, fertility levels and biofertilizers on growth, yield and quality of sorghum. *Indian Journal of Agronomy* 49(2):117-120.
- Patel PR, Patel BJ, Vyas KG, Yadav BL. Effect of integrated nitrogen management and bio-fertilizer in *Kharif* pearl millet (*Pennisetum glaucum* L.). *Adv. Res. J. Crop Improv.* 2014; 5(2):122-125.
- Ramdev, T., LR Yadav, SL Choudhary and GV Shisuvinahalli. (2017). Effect of Azotobacter on growth, yield and quality of pearl millet (*Pennisetum glaucum* L.). *Journal of Pharmacognosy and Phytochemistry*, 6(4): 889-891.
- Rathore, V.S., Singh, P. and Gautam, R.C. 2006. Productivity and water-use efficiency of rainfed pearl millet (*Pennisetum glaucum*) as influenced by planting patterns and integrated nutrient management. *Indian J. Agron.*, 51(1):46-48.
- Sushila, R. and Giri, G. 2000. Influence of farmyard manure, nitrogen and biofertilizer on growth, yield attributes and yield of wheat (*Triticum aestivum* L.) under limited water supply. *Indian Journal of Agronomy*, 45(3):590-595.
- Venkata Lakshmi, K., Integrated nutrient management for dryland fodder sorghum (*Sorghum bicolor* L. Moench)- chickpea (*Cicer arietinum* L.) + coriander (*Coriander sativum* L.) cropping system. *M.Sc. (Ag.) Thesis. Tamil Nadu Agricultural University, Coimbatore, India.* (2001).
- Zothanmawii., Edwin Luikham and Mariam Anal, P.S. 2018. Growth and yield of hybrid maize as influence by levels of nitrogen and biofertilizer. *International Journal of Current Microbiology and Applied Sciences*, 7(8):

**Table 1. Effect of Bio fertilizers and Phosphorus on Growth attributes of Pearl millet**

Sr.no	Treatments	Plant height (cm) 80 DAS	Dry weight (g/plant) 80 DAS
1	20 g/ kg <i>Azotobacter</i> + 35 kg/ ha Phosphorus	171.23	30.70
2	20 g/ kg <i>Azotobacter</i> + 40 kg /ha Phosphorus	179.80	31.19
3	20 g/ kg <i>Azotobacter</i> + 45 kg/ ha Phosphorus	185.43	33.59
4	20 g /kg <i>Azospirillum</i> + 35 kg /ha Phosphorus	175.03	34.22
5	20 g/ kg <i>Azospirillum</i> + 40 kg/ ha Phosphorus	183.17	34.30
6	20 g/ kg <i>Azospirillum</i> + 45 kg/ ha Phosphorus	188.70	33.88
7	20 g/ kg <i>Azotobacter</i> + <i>Azospirillum</i> + 35 kg/ha Phosphorous	192.10	34.87
8	20 g/ kg <i>Azotobacter</i> + <i>Azospirillum</i> + 40 kg/ha Phosphorous	196.40	37.67
9	20 g/ kg <i>Azotobacter</i> + <i>Azospirillum</i> + 45 kg/ha Phosphorous	200.13	40.33
10	Control (RDF 80-40-40 N-P-K kg/ha )	168.13	31.87
	F-test	S	S
	SEm(±)	1.40	0.93
	CD (p=0.05 )	4.15	2.75

**Table 2. Effect of Bio fertilizers and Phosphorus on Yield attributes and Yield of Pearl millet**

Sr.no	Treatments	Ear head Length (cm)	No.of Grains/Ear head	Test Weight (g)	Grain yield (t/ha)	Stover yield (t/ha)	Harvest index (%)
1	20 g/ kg <i>Azotobacter</i> + 35 kg/ ha Phosphorus	17.29	1597.2	5.47	2.38	4.15	36.94
2	20 g/ kg <i>Azotobacter</i> + 40 kg /ha Phosphorus	19.44	1682.9	6.03	2.31	4.30	35.49
3	20 g/ kg <i>Azotobacter</i> + 45 kg/ ha Phosphorus	20.81	1796.17	6.13	2.22	4.47	33.66
4	20 g /kg <i>Azospirillum</i> + 35 kg /ha Phosphorus	18.05	1613.4	6.00	2.70	4.98	36.22
5	20 g/ kg <i>Azospirillum</i> + 40 kg/ ha Phosphorus	19.85	1747.5	6.43	2.75	5.15	35.23
6	20 g/ kg <i>Azospirillum</i> + 45 kg/ ha Phosphorus	21.52	1891.8	6.40	2.61	5.20	33.80
7	20 g/ kg <i>Azotobacter</i> + <i>Azospirillum</i> + 35 kg/ha Phosphorous	22.55	1902.0	6.63	2.98	5.32	36.79
8	20 g/ kg <i>Azotobacter</i> + <i>Azospirillum</i> + 40 kg/ha Phosphorous	23.87	1981.0	6.53	3.15	5.64	36.05
9	20 g/ kg <i>Azotobacter</i> + <i>Azospirillum</i> + 45 kg/ha Phosphorous	25.25	2010.9	6.27	3.25	6.27	34.26
10	Control (RDF 80-40-40 N-P-K kg/ha )	15.84	1404.8	5.47	1.96	4.13	32.55
	F-test	S	S	NS	S	S	NS
	SEm(±)	0.27	32.00	0.61	0.04	0.27	1.17
	CD (p=0.05 )	2.33	95.06	-	0.13	0.80	-