

# A case of highly located thyroglossal duct cyst

## Abstract:

Congenital anterior neck masses comprise a rare group of lesions that are frequently found in children. The most frequent abnormalities are ectopic thyroid tissue and malformations of the thyroglossal duct, particularly the thyroglossal duct cyst.

The most frequent congenital neck mass is a thyroglossal duct cyst (TGDC), which frequently manifests in young patients as a painless lump in the middle of the neck. When an adult develops a lateralized neck lump for the first time, it may raise more concern for malignancy.

A thyroglossal duct cyst is usually suspected based on clinical examination, imaging can confirm the diagnosis, assess the extent, and evaluate for associated complications. We report the case of a sixty years old female patient presenting with a submental swelling, suspected to be a laryngocele initially. Surgery and histology revealed a giant and high located thyroglossal duct cyst.

**Keywords:** Thyroglossal, Cyst, Laryngocele, Thyroid, neck mass

## Introduction:

“Thyroglossal duct cysts (TGDC) represent 70% of all congenital neck anomalies and are the most common type of head and neck

congenital cysts” [1,11].

It often manifests as a midline neck lump that is asymptomatic and moves upward with swallowing and tongue protrusion [2,10].

TGDC typically manifests in kids and teenagers. However, those 20 years of age or older are the ones who receive the diagnosis in up to one-third of cases. Equal amounts of both genders are impacted [3,13].

There are a number of differential diagnoses that should be taken into account, including branchial cleft cysts, dermoid/epidermoid cysts, laryngocele, thymic cysts, lymphatic malformations, and metastasis.

In order to correctly diagnose an anterior neck mass, it is essential to have a thorough understanding of the embryologic process of thyroid development and be able to identify key landmarks such the foramen cecum, hyoid bone, thyroid cartilage, and strap musculature.

## **Case report**

It's about a 60-year-old woman, with a medical past history of cholecystectomy, that was referred to our department for a 4-year developing submental mass, increasing in volume progressively. The evolution was marked one year ago by the apparition of dyspnea in the dorsal decubitus, along with nocturnal snoring.

Clinical examination revealed a

submental anterior mass, renitent, mobile, ascending during swallowing, measuring 5 cm.

Nasofibroscopy revealed a bulging in both laryngeal ventricles, suspecting a laryngocele.

Ultrasound showed a submental cystic mass, with no vascular characteristic in the doppler.

The CT scan showed an anterior cystic mass, occupying the hyo-thyro-epiglottic loge, limited, measuring 49 x 35 mm. The CT scan was in favor of a thyroglossal duct cyst.

The patient underwent surgery, and intra-operative exploration revealed a thyroglossal duct cyst. The cystic lesion was dissected from the surrounding tissues and fully excised together with the center of the hyoid bone. Tissues from the superior aspect of the hyoid bone up to the foramen cecum were removed. The cyst was not ruptured and was removed completely.



Cases typically present with a mobile, painless midline neck mass that is inferior to the hyoid bone (about 75% of cases), moving with a tongue protrusion.

10% of patients show signs of infection, including the development of fistulas, however children are more likely to do so than adults. The precise site of involvement is revealed through imaging studies.

The Sistrunk surgery, which also includes a cylinder of tissue at the base of the tongue and the middle region of the hyoid bone, is the preferred course of treatment for TGDC [9].

“According to histology, the lining of TGDCs is made up of either respiratory epithelium, squamous epithelium, or a combination of the two” [5,6].

The epithelium may become obscured by granulation-type tissue or a foreign-body giant cell reaction when there is an infiltration of inflammation, which is always present. In 70% of cases, microscopic foci of ectopic thyroid gland tissue will be seen inside the cyst's wall or inside the nearby soft tissue. But sometimes, the foci are little and necessitate cautious

examination, which may occasionally involve a TTF1 immunohistochemistry investigation.

“An associated thyroid gland cancer may be present in about 3% of cases; the majority (>99%) are papillary thyroid carcinomas. The anatomic site, histologic lining, and recognition of thyroid tissue aid in this differentiation between a branchial cleft cyst, a bronchogenic cyst, an epidermal inclusion cyst, or a dermoid cyst, which are all possible differential diagnoses” [5,6,12].

Several studies focused on adults with TGDC who presented as asymptomatic lateral neck masses; the diagnosis was unclear in all of these cases until pathological assessment following cyst removal. These examples highlight the idea that doctors examining asymptomatic lateral neck tumors in adults should consider TGDC [7,8].

## **CONCLUSION:**

Thyroglossal duct cysts (TGDCs) may occur in adults and may be lateralized, however they most frequently manifest in children as

midline masses.

Surgery for TGDC requires the Sistrunk surgery, which involves removing the hyoid bone's center.

It should be noted that lesions that have TGDC on the differential diagnosis should undergo additional preoperative examination, in order to perform appropriate surgical treatment.

#### **Consent**

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

### **ETHICAL APPROVAL:**

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

### **REFERENCES:**

1. D.L. Reede, R.T. Bergeron, P.M. Som CT of thyroglossal duct cysts Radiology, 157 (1) (1985), pp. 121-125
2. K.P. Oomen, V.K. Modi, J. Maddalozzo Thyroglossal duct cyst and ectopic thyroid: surgical management Otolaryngol Clin, 48 (1) (2015), pp. 15-27
3. F.M. Gioacchini, *et al.* Clinical presentation and treatment outcomes of thyroglossal duct cysts: a systematic review Int J Oral Maxillofac Surg, 44 (1) (2015), pp. 119-126
4. Ren W, Zhi K, Zhao L, Gao L. Presentations and management of thyroglossal duct cyst in children versus adults: A review of 106 cases. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2011;111(2):e1-6.
5. Thompson LD, Herrera HB, Lau SK. A clinicopathologic series of 685 thyroglossal duct remnant cysts. Head Neck Pathol 2016;10(4):465-74.
6. Wei S, LiVolsi VA, Baloch ZW. Pathology of thyroglossal duct: An institutional experience. Endocr Pathol 2015;26(1):75-9.
7. Yamada S, Noguchi H, Nabeshima A, et al. Papillary carcinoma arising in thyroglossal duct cyst in the lateral neck. Pathol Res Pract 2013;209:674
8. Wakasa Y, Oishi S, Nara M, et al. [Papillary carcinoma arising in thyroglossal duct cyst in the right lateral neck]. Gan To Kagaku Ryoho 2017;44:1152-4.
9. Quah HM, Lim DT. Unusual endolaryngeal extension of a large thyroglossal duct cyst. Ann Acad Med Singapore 2002;31:238-40
10. Kacker A, Komisar A. Imaging quiz case 1. Thyroglossal duct cyst with intralaryngeal extension. Arch Otolaryngol Head Neck Surg 1996;122:1266-8

11. Slotnick D, Som PM, Giebfried J, Biller HF. Thyroglossal duct cysts that mimic laryngeal masses. *Laryngoscope* 1987;97:742-5
12. Brousseau VJ, Solare CA Xu M, Krakovitz P, Koltai PJ. Thyroglossal duct cyst. Presentation and management in children versus adults. *Int J Paediatric Otorhinolaryngology* 2003; 67: 1285-90
13. Mondin V, Ferlito A, Muzzi E, Silver CE, Fagan JJ, Devaney KO, Rinaldo A. Thyroglossal duct cyst: personal experience and literature review. *Auris Nasus Larynx*. 2008 Mar 1;35(1):11-25.

