

## Original Research Article

### Aviary caged hens with unusual lymphocytes and normal H/L ratios

#### ABSTRACT

The aim is to describe hemograms of hens that bear upon interpretation of the traditional H/L stress determination method. Hemograms from aviary caged (AV) commercial hens sampled at 18 wk with low to moderate heterophil/lymphocyte ratios (H/L, range 0.02 – 0.66) were accompanied by slightly elevated total white cell counts (TWBC ~ 30K/ $\mu$ L) to leukocytosis (TWBC ~ 120K/ $\mu$ L) levels. Atypical cells were commonly seen in Wright-Giemsa stained blood by standard differential counts (SDC). Heterophils exhibited signs of toxicity affecting both the nucleus, the cytoplasmic granules, and cell membranes. Many forms of atypical lymphocytes were also detected. These were aggregated small lymphocytes (Ls), medium sized reactive lymphocytes (Lm) plasmacytes (PC) including Mott cells, and other atypical forms. Some PCs contained pink cytoplasmic vacuoles (Russell bodies) indicating they are IgA types. Developmental PCs rarely found in the circulation of homeostatic avians were also present. Bacteria were commonly associated with the atypical cells. These were both free swimming and cell associated (CAB) types. Fungal forms were also present including yeast-like conidia and hyphae. When microorganisms were in a field the background erythrocytes were in Rouleaux formation, an indication of an inflammatory milieu. In conclusion, it is demonstrated that in the absence of detailed cytological descriptions the H/L ratio can be misleading. Given its wide usage in stress and welfare determinations these observations emphasize some of the difficulties of the simple H/L method. They reinforce earlier observations and draw attention to the necessity of cytological detail before the computed H/L ratio can be accepted as a stress measure.

**Key words:** commercial hens, heterophil lymphocyte ratio, atypical lymphocytes, plasmacytes

#### ABBREVIATIONS

$A_c$ ,  $R_c$ : Area of cell in  $\mu\text{m}^2$ , cell radius in  $\mu\text{m}$

AV: aviary cage

CAB: cell associated bacteria

H/L: heterophil to lymphocyte ratio

IgA: immune globulin A

Ls, Lm: small, medium size lymphocytes

PC: plasmacyte

SDC: standard differential count

TWBC: total white blood count

**Comment [R11]:** Abbreviations are important, however - they must be established as they appear in the text - their use in this way (in list) is not recommended

#### 1. INTRODUCTION

Hemogram data derived from either hemocytometer or standard differential counts (SDC) are widely used to estimate stress. Attention is directed toward the ratio of heterophils to lymphocytes (H/L). The

theory is based on a series of studies by Davidson and Rowell (1) and Gross and Siegel (2,3, 4) who observed increased H/Ls in experimental chickens treated with corticosterone, exposed to social stress, or injected with bacteria. H/L values about 0.5 accompanied by total white blood counts (TWBC) about 30K/ $\mu$ L would indicate homeostasis. However, in many instances H/Ls are reported without detailed descriptions of the cells used for its computation or the TWBC (5). A low H/L accompanied by a high TWBC cannot receive the same physiological interpretation as one with a normal TWBC (6,7).

As neither heterophils nor lymphocytes are homogeneous (8, 9) it is crucial to define which are included in the H/L computation. Heterophils of caged hens and ducks were sorted into 3 categories based on cytoplasmic granules and nuclear configuration. Typical heterophils (HT) with faint granules, the most frequent type, are distinct from classic heterophils (HC) with deep red spindle shaped granules. A third heterophil (variant type, HV) with orange spherical granules also occurs (9,10,11) Therefore, it is possible that the same H/L value may be associated with distinct heterophil spectra.

Lymphocyte counts are used as the H/L denominator often without regard to their origin or if they are reactive. In an SDC from a homeostatic source most lymphocytes should be small resting (T-cell) types. Reactive types or plasmacytes are not suitable for use in the H/L. Therefore, the aim of this manuscript is to expand observations of leukocytes whose cytology would render the simple H/L an inappropriate stress measure. The source are commercial hens housed in aviaries and sampled at 18 wk. The information should be of interest to those who rely on the H/L computation to assess stress and establish welfare status.

## 2. MATERIALS AND METHODS

### 2.1 Chickens and Welfare

Chicks of a Lohmann White Egg commercial type (LSL) were housed in AV at 850 to 1,700 hens per compartment. They were vaccinated at the hatchery for laryngotracheitis and Marek's disease. They were managed according to a typical program designed for commercial poultry. The experimental protocol was approved by the Michigan State University Institutional Animal Care and Use Committee.

Comment [R12]: Reference?. It does not appear in the reference list?

### 2.2 Blood, Stain Procedure, H/L and TWBC Computations

Blood samples were collected from wing veins into EDTA tubes and stained later using Wright-Giemsa. Differential counts (SDC): a minimum of 200 leukocytes per slide were sorted into categories: small or medium lymphocytes, monocytes, heterophils, basophils, or eosinophils. Morphological criteria for sorting were as described in (9-11). Division of the sum of all heterophil types (typical, variant, and classic) by the number of small "resting" lymphocytes is H/L 1. Division of the same heterophil value by the sum of all lymphocyte types, (resting, reactive, and atypical) is H/L 2. The difference, H/L 1 – H/L 2 is expressed as  $\Delta$ H/L. The average number of white blood cells in five 40 $\times$  microscopic fields multiplied by 4,000 provides the total white blood cell count (TWBC/  $\mu$ L) estimate, after a modification of a method described by Campbell and Ellis (12).

Comment [R13]: what is the name of the vein?

### 2.3 Light Microscopy and Photomicrographs

An Olympus CX-41(Olympus America, Center Valley, PA) fitted with Plan N 40×, 0.65 Numerical Aperture dry, and Plan N, 1.25 N.A. 100× (oil) objectives. The images were captured by an Infinity2 1.4 Megapixel CCD USB 2.0 Camera and processed with Infinity Analyze software (Release 6.5, Lumenera Inc., Ottawa, Ontario, Canada). Magnification was 100× (oil).

### 3. RESULTS

Standard differential counts (%) total white cell counts, and H/L ratios for the hens of Figures 1-5 and the average values for all hens in the study are in Table 1. For simplicity all heterophil types (typical, classic and variant) are combined into one category "H" and "Bst" represents a combination of granuloblasts (meso and metamyelocytes).

As heterophils represent the numerator of the H/L computation examples of types seen in the source population are in Figure 1. The cells of both panels A and B are standard size band (young) cells [HT, R<sub>c</sub> 6.7 μm, A<sub>c</sub> 141 μm<sup>2</sup>] with a single lobe usually the youngest types. The cytoplasmic granules of each are poorly stained suggesting toxicity. The HT of panel C has 3 well-stained lobes but its cm is irregular. The variant heterophil of panel D is smaller [HV, R<sub>c</sub> 4.5 μm, A<sub>c</sub> 64 μm<sup>2</sup>] It has 3 faint nuclear lobes and orange spherical cytoplasmic granules. Collectively each cell of figure 1 exhibits some form of atypia.

Reactive lymphocytes and plasmacyte(13) examples are in Figure 2. Panel A. Binuclear bi/PCs with patchy cytoplasm, a cell type suggesting stress and possible viremia (14) and more traditional PCs with well developed Hof. Panel B. Reactive small lymphocytes (Ls) in the form of multicellular aggregates. Panel C. Ls with scanty cytoplasm and larger reactive Lm cells with vacuolated cytoplasm; likely a plasmacyte. Panel D. An atypical plasmacyte (\*PC) with paracrystalline Russell bodies; an indication of a defective antibody. Either free or cell associated bacteria (CAB) are at the arrows of panels B, C, and D.

Additional examples of reactive lymphocytes are in Figure 3. Panel A. a reactive lymphocyte (Lm) with plasmacyte cytoplasmic characteristics (Hof) is in a field with a Ls whose cm contains blebs (zeiosis) an indication of toxicity. Panel B. A second example of a plasmacytoid Lm in a field contrasts with a (resting) Ls (inset). The background RBC of both panels A and B are Rouleaux (adhering) forms. Panel C. An Ls is adhering to a Lm/PC whose cytoplasm is fenestrated with small and large clear vacuoles. Remnants of 2 lysed nuclei (?RBC) are also present (lower right). Panel D. A solitary Lm/PC (right arrow) is attached to RBCs (left and down arrows).

Examples of atypical plasmacytes with pink vacuoles are in Figure 4. Panel A. Solitary PC with an irregular cm and cytoplasm with pink and orange vacuoles (Russell bodies) in a field with aggregated

**Comment [R14]:** In the results- It is not recommended that references appear.

thrombocytes (Th) and RBCs (15). Panel B. Two cells that are mutually adhering plasmacytes with pink Hof's and patchy cytoplasm. Panel C. A PC with a pink Hof is in a field with an elongated bacillus containing a terminal endospore (arrow). Panel D. A proplasmacyte (Türk cell) with a giant nucleolus. A pseudopod is extending from the cytoplasm at the lower left. The inset is a 2.5x enlargement of a mixed species microcolony seen to the left of the Türk cell.

Additional examples of reactive cells from the study flock are in Figure 5. Panel A. A large plasmacytoid cell (Lm/PC) is associated with mesomyelocytes: developmental cells of the granulocytic series. Small lymphocytes (Ls) and a lysed nucleus of uncertain origin are nearby. Panel B. A pair of monocytoïd cells (Mn) with reactive cytoplasmic characteristics. Panel C. A (?mixed) bacterial microcolony composed of bacilli of varying lengths is attached to a RBC, itself attached to another RBC. The predominantly magenta color of the bacteria suggests these are Mollicutes, bacteria with thin cell walls. Panel D. Conidia at different stages of development are attached to a Lm.

A scatter plot of H/L ratios for all hens of the flock is in Figure 6. The large data points are from the hens providing the figures. The open circles are the other hens of the flock. Reference lines, H/L 1 at 0.5, and H/L 2 at 0.4 represent theoretical homeostasis (low or no stress) values.

#### 4. DISCUSSION

The objective of this study is to expand on observations of the robustness of traditional H/L stress detection method. In earlier work the H/L was divided into two categories H/L 1 and H/L 2 as an attempt to recognize the importance of cytology in establishing stress levels (5,8,11,16). The value of H/L 1 is determined by using all heterophil categories as the numerator and only small lymphocytes (Ls) in the denominator as these should be more common in the absence of high stress levels. In contrast H/L 2 is computed using both small (Ls) and large lymphocytes (Lm) (see Cotter, 5). If there is minimal stress Lm cells are few and there is little difference in the two measurements. Therefore, the  $\Delta H/L$  would be  $\leq 0.1$ ; as was true for the hens of this study (Table 1). However, in many reports the H/L value is given in the absence of the total white blood count (TWBC) and too few cells counted for the SDC (see Archer, as recent examples (17,18)). Here the TWBC ranged from the homeostatic level ( $\sim 30k/\mu L$ ) to well into leukocytosis (TWBC  $> 100k/\mu L$ ; Table 1). Remarkably atypical cells of all series were found in the subject hens independent of either H/L or TWBC values (Figure 6).

Furthermore, many authors do not recognize that multiple heterophil types exist (Lucas and Jamroz, 1961, (9) as do lymphocytes. This is despite their possession of multiple granule types and physiologic functions (as reviewed in Genovese et al., 19). This deficiency includes some who take extraordinary care in experimental protocol but fail to consider cytology (see Lentfer, et al. 20 as an

Comment [R15]: review and adapt the writing to be able to make the appointment correctly

example). Therefore, some of these H/Ls might be contaminated by including toxic and atypical cells. A situation that immediately suggests stress, inflammation, and disease.

Reactive and apoptotic heterophils were described in hens from the same population as the subjects of this study but housed in conventional cages (20). Such cells indicate inflammation and infection (Latimer, et al, 21). Here bacteria and fungi were present in many of the blood films (Figures 2-5) at levels high enough to indicate bacteremia/fungemia rather than as casual contaminants of blood. Not only were microorganisms found in the blood but many were attached to RBCs and other cells. These types are referred to as cell associated bacteria (CAB) are indications that erythrocytes should be given consideration in interpretation of hemograms.

In addition, plasmacytes were common. These cells are designed to secrete antibody but are rarely found among circulating leukocytes. Not only were PCs frequent but many were atypical; including Mott cells, a PC known to be defective (8,22).

## 5. CONCLUSION

The cells of a homeostatic SDC should display a standard non-reactive appearance. Size, shape, N/C ratio, and an Romanowsky-Giemsa effect (RGE, staining) must be appropriate for each series. This is a critical issue in evaluating the H/L. Developmental stages and dividing cells may be found among the circulating community but these should be rarely seen in a sample from an avian at homeostasis.

Atypia, especially heterophils, indicate a complex hemogram, already beyond the quiescent (homeostatic) stage. The question of the relation of stress to excess monocytes, is not considered here or are reactive basophils (23) both need further attention.

## DECLARATION

The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## REFERENCES

1. Davidson TF, JG Rowell, and J Rea. 1983. Effects of corticosterone on peripheral lymphocyte and granulocyte populations in immature domestic fowl. Res. Vet. Sci. 14:236-239. PMID: 6856997
2. Gross WB and HS Siegel, 1983. Evaluation of the heterophil lymphocyte ratio as a measure of stress in chickens. Avian Dis., 27: 972-979. PMID: 6360120
3. Gross WB and PB Siegel, 1986. Effects of initial and second periods of fasting on heterophil/lymphocyte ratios and body weight. Avian Dis., 30: 345-346. PMID: 3729879
4. Gross WB, 1989. Factors affecting chicken thrombocyte morphology and the relationship with heterophil:lymphocyte ratios. Brit. Poultry Sci. 30:919-925. doi: 10.1080/00071668908417218. PMID: 2697458

5. Cotter PF, 2022a. Letter to the Editor regarding, Research Note: Sex difference in changes in heterophil to lymphocyte ratios in response to acute exposure of both corticosterone and cortisol in the Pekin duck. *Poultry Science* [https://authors.elsevier.com/sd/article/S0032-5791\(22\)00349-2](https://authors.elsevier.com/sd/article/S0032-5791(22)00349-2)
6. Cotter PF, 2015a. An examination of the utility of heterophil lymphocyte ratios in assessing stress of caged hens. *Poult. Sci.* 94:512–517.
7. Cotter PF, 2015b. Atypical lymphocytes and leukocytes in the peripheral circulation of caged hens. *Poult. Sci.* 94:1439– 1445.
8. Cotter PF, 2015c. Are peripheral Mott cells an indication of stress or inefficient immunity? *Poult. Sci.* 94:1433–1438.
9. Lucas AM and C Jamroz, 1961. *Atlas of Avian Hematology*. U.S.D.A. Agricultural Monograph 25 Washington
10. Cotter PF, 2021. Apoptosis of circulating heterophils; implications for the interpretation of the heterophil/lymphocyte ratio. *J Immunological Sci.* 5(4): 1-8 [DOI:10.29245/2578-3009/2021/4.1222](https://doi.org/10.29245/2578-3009/2021/4.1222)
11. Cotter, PF and ED Heller, 2016. Complex Hemograms of Isolator Raised Specific Pathogen Free (SPF) Chicks. *International Journal of Poultry Science* 15 (6): 211-217 [ISSN 1682-8356](https://doi.org/10.1186/s12917-016-0835-6)
12. Campbell TW and CK Ellis, 2007. *Avian and Exotic Animal Hematology and Cytology*. 3rd Ed., Blackwell Publishing, Ames, Iowa
13. Cotter PF, 2022b. Plasmacyte heterogeneity of lame ducklings. *J Immunological Sci.* 6(1): 7-14 [doi: 10.29245/2578-3009/2022/1.1225](https://doi.org/10.29245/2578-3009/2022/1.1225) <https://www.immunologyresearchjournal.com/public/>
14. Cotter PF, 2022c. A microscopic study on morphology of reactive thrombocytes in duckling. *J. World Poult. Res.*, 12 (3): 142-150. [DOI: https://dx.doi.org/10.36380/jwpr.2022.16](https://dx.doi.org/10.36380/jwpr.2022.16)
15. Cotter PF, 2023. P230. Avian polyclonal B-cell lymphocytosis-a stress indication or a consequence of infection? *International Poultry Scientific Forum*, Atlanta, Georgia, January 23-24
16. Cotter PF, 2022d. Stress assessment by the hemogram method - circulating cells complicating reliance on Heterophil/Lymphocyte (H/L) Ratio. *J Vet Med Res* 9(1): 1224 <https://doi.org/10.47739/2378-931X/1224>
17. Archer GS, 2018. Color temperature of light-emitting diode lighting matters for optimum growth and welfare of broiler chickens. *Animal* 12: (5) 1015-1021. <https://doi.org/10.1017/S1751731117002361>
18. Archer GS, 2019. How does red light affect layer production, fear, and stress? *Poult Sci.* 98(1):3-8. [doi: 10.3382/ps/pey302](https://doi.org/10.3382/ps/pey302). [PMID: 30690610](https://pubmed.ncbi.nlm.nih.gov/30690610/).
19. Genovese KJ, et al, 2013. The avian heterophil. *Dev. Comp. Immunol.* 41(3):334-340 [http://dx.doi.org/10.1016/j.dci.2013.03.021](https://dx.doi.org/10.1016/j.dci.2013.03.021)
20. Lentfer TL et al. 2015. H/L ratio as a measurement of stress in laying hens – methodology and reliability, *British Poultry Science*, 56(2): 157-163. [DOI: 10.1080/00071668.2015.1008993](https://doi.org/10.1080/00071668.2015.1008993)

21. Latimer KS et al. 1988. Leukocyte changes associated with acute inflammation in chickens. Avian Diseases 32(4): 760-772 <http://www.jstor.org/stable/1590996>
22. Cotter PF, MR Bakst, 2017. A comparison of Mott cell morphology of three avian species. II. - Bad behavior by plasmacytes? Poult Sci. 96(2):325-331. doi: 10.3382/ps/pew288.
23. Cotter PF, 2017. Basophilia and basophilosis in caged hens at 18 and 77 weeks. Int. J. Poult. Sci.16:23-30. DOI: 10.3923/ijps.2017.23.30
24. Swanson JC, JA Mench JA, D Karcher, 2015. The coalition for sustainable egg supply project: An introduction. Poult Sci.94(3):473-474. doi: 10.3382/ps/peu012.

UNDER PEER REVIEW

**TABLE 1.** Unusual lymphocytes and normal H/L ratios

Source	No.	H <sup>1</sup>	Ls	Lm		Bst	Mn	Ba	Eo	TWBC(K)	H/L 1	H/L 2	ΔH/L
Figures	7	14.1	67.8	9.3		0.2	3.3	4.3	0.9	81.1	0.27	0.20	0.06
Flock	41	14.3	73.7	4.4		0.2	3.6	3.4	0.5	69.8	0.21	0.19	0.01

**Comment [R16]:** Take care of the style: by the use of different type and size of letter

1. Cells: H, heterophil (Σ [HT typical, HC classic, HV variant.]) Ls small lymphocyte ~6 μm diameter, Lm medium, large (diameter 8–10 μm) Bst, granulocyte developmental cell, Mn, monocyte, Ba, basophil, Eo, eosinophil. TWBC (K), total white blood cells per cubic μL in thousands (K).  $H/L\ 1 = (HC + HT + HV) / Ls$ ;  $H/L\ 2 = (HC + HT + HV) / (Ls + Lm)$ ,  $\Delta H/L = H/L1 - H/L2$ .

UNDER PEER REVIEW

UNDER PEER REVIEW





Figure 1. Examples of atypical (A, B band stages; C, 3 lobe stage) heterophils (HT) characteristic of the SDC from the subject hens. A variant heterophil (HV) is in panel D. Additional descriptions are given in the text.



**Comment [R17]:** It is recommended - to eliminate the text: Additional descriptions are given in the text / since a figure (the title) must be explicit by itself, unless it is required to highlight more or other relevant aspects - it is advisable to make a detailed description.

Figure 2. Panel A. Binuclear bi/PC with patchy cytoplasm and a large reactive lymphocyte (Lm/PC). Panel B. A group of 9 Ls and a larger reactive lymphoid cell. The remnants of a lysed basophil (Ba) are among the Ls. A solitary encapsulated bacterium is at the arrow. Additional descriptions are given in the text.




Figure 3 | Panel A. An example of a reactive lymphocyte with plasmacyte cytoplasmic characteristics. Arrows locate bacteria. Panel B. A second example of a plasmacytoid Lm in a field with a (resting) Ls (inset). Panel C. A Ls is adhering to a Lm/PC. Remnants of 2 lysed nuclei (RBC) are also present (lower right). Panel D. A solitary Lm/PC (right arrow) are attached to RBCs (left and down arrows). Additional descriptions are given in the text.




Figure 4. Panel A. Solitary PC with an irregular cm and cytoplasm with pink and orange vacuoles (Russell bodies) thrombocytes (Th) and RBCs. Panel B. Two adhering plasmacytes. Panel C. Plasmacyte with a pink Hof in a field with an elongated bacillus containing a terminal endospore (arrow). Panel D. Proplasmacyte (Türk cell). Inset a 2.5x enlargement of a mixed species microcolony. Additional descriptions are given in the text.




Figure 5. Panel A. A large plasmacytoid cell (Lm/PC) mesomyelocytes, small lymphocytes (Ls) and a lysed nucleus. Panel B. Monocytoid cells (Mn). Panel C. A (?mixed) bacterial microcolony attached to a RBC. Panel D. Conidia at different stages of development are attached to a Lm. Additional descriptions are given in the text.





Figure 6. Scatter plot of H/L 2 vs H/L 1 ratios for all AV housed hens in the study. Filled circles locate hens of hematology figures. Broken lines are expected values for homeostasis. Reference lines (H/L 1 = 0.5; H/L 2 = 0.4) are expected values of homeostatic samples. Additional descriptions are given in the text.



UNDER PEER REVIEW