

Characterization and Study of Dominated Alluvial Soil Profile of Mirzapur in Indo-Gangetic Plain (IGPs) of India

Abstracts

This provides the basic information necessary to create functional soil classification schemes and assess soil fertility in order to unravel some unique soil problems in an ecosystem. Mirzapur district is situated in the southern part of Uttar Pradesh and is located between 25.8° N to 25.15° N latitude and 82.34° E to 82.58° E longitude covering an area of 4952.5 km². The soils of the study region comprised of alluvial soil representing entisol and soils which are formed on granitic parent material representing Alfisols. The coupling of soil characterization, soil classification and soil mapping provides a powerful resource for the benefit of mankind especially in the area of food security and environmental sustainability. Among the cationic micronutrients, Zn content varied between 1-3 ppm in the Mirzapur soil profile. Copper content varied between 1-7 ppm in Jamalpur, 1-4 ppm in Narayanpur. The available Cu content is categorized as sufficient if it is > 0.2 ppm. Available iron content varied between 2-40 ppm in Jamalpur soil profile and 8-63 ppm in the Narayanpur soil profile. Available Mn content varied between 2-40 ppm in Jamalpur, 5-18 ppm in Narayanpur. Soils are generally categorized as having enough manganese in the DTPA extractable Mn content is > 0.2 ppm. Hence all soils of Jamalpur and Narayanpur had sufficient available soil manganese.

Keywords- Soil profile; Soil fertility, soil characterization, soil classification, environmental sustainability

1. Introduction

There is an increasing demand for information on soils as a means to produce food (Fasinaet *al.*, 2007). Agriculture is the predominant economic activity in India and because of agricultural development and increasing demand in India, much work is carried out on soil characterization. This provides the basic information necessary to create functional soil classification schemes, and assess soil fertility in order to unravel some unique soil problems in an ecosystem (Lekwaet *al.*, 2004). The coupling of soil characterization, soil classification and soil mapping provides a powerful resource for the benefit of mankind especially in the area of food security and environmental sustainability. Soil use refers to the conversion of a whole or a part of the soil for specific purposes like agricultural activities, house building, or industry purposes etc. (Rai, 2015). It also provides adequate information in terms of landform, natural vegetation as well as characteristics of soils which can be utilized for land

resources management and development (Manchanda *et al.* 2002). The alluvium deposit of both old and new types by the two great rivers of India, the Ganga and the Yamuna are found in this region, only the southern uplands, touching the Vindhya hills are not covered by such deposits. Sandstone, shale, conglomerate and limestone are found in the SunValley region of Mirzapur. The sands of the Kaimur group are of two to three-metre thick friable sandstone. The silica sand found in Chandauli district is fine to medium-grained, brown and buff in colour. Most of the glass factories of northern India get their supply from this region. Similar sands are reported from Robertsganj plateau of Mirzapur district. Besides kankar, which is found in abundance in various parts of the study area, suitable limestone occurs in some places of Mirzapur district (Agarwal, *et al.*, 1952). The rivers rising from the mountains during the period of great gradational activity deposited the detritus brought down by them in their long journey and in this way the plain was formed (Wadia, 1975). The sediments deposited at the bed of the Tethys Sea was folded and warped due to the northward drift of peninsula (Khullar, 2005). This rolling upland, touching the Vindhya hills exhibits a complex and heterogeneous nature of topography with detached hills, flat-topped ridges, summit plains and entrenched narrow as well as broad valleys almost reaching the base level. In the eastern part, the east-west trending Vindhayan range composed of shallow marine deposits of proterozoic age, divides the Ganga plain in the north and vast pediplain exposing metamorphosed sequence of Archaean age in the south (NATMO, 2008). Based on the micro-level topographic facets, the Ganga plain can be divided into three sub-regions, viz. (i) the older alluvium or *Bhangar*; (ii) the newer alluvium or *khaddar* and (iii) the *terai* adjoining to the *bhabar* area is the *terai*. Geologically Eastern Uttar Pradesh is made up of diverse rock types, ranging in age from the oldest.

Owing to the fact that Mirzapur district is an agrarian community outskirt and not much study has been done on the soils of the area, characterization and classification will help reveal information that could be useful in the management and use of the soils on a sustainable manner. The objective of this research therefore, is to characterize and classify the soils of Mirzapur District of Uttar Pradesh.

2. Material and methods

2.1. Study area

Mirzapur district is situated in the southern part of Uttar Pradesh and is located between 25.8° N to 25.15° N latitude and 82.34° E to 82.58° E longitude covering an area of

4952.5 km². The soils of the study region comprised of alluvial soil representing entisol and soils which are formed on granitic parent material representing Alfisols.

2.2. Method of soil collection and preparation

A total of five profiles were dug and samples were collected from each location. To collect a soil sample, surface litter was gently scraped off with a khurpi. A rectangular pit was dug to a depth of 1m x 1 m x 1m. Soil samples were collected from the wall of the rectangular pit using stainless steel auger from a depth of 0-5, 5-15, 15-30, 30-60 and 60-100 cm. In addition, to measure the bulk density, core samplers having cores measuring 5 cm in length and 5 cm in diameter were used. Soil samples were dried in shade and brought to the laboratory where they were ground to pass through 2 mm sieve, tagged and stored in plastic containers for analysis.

2.3. Analysis of samples

The bulk density was estimated using the soil core sampler method (Blake, 1965). Soil colour was measured using a Munsell's soil colour chart. For this air-dried soil was taken on plastic sheets and the colour matched with the colour notation of the chart. The Munsell system divides colour into hue; value; and chroma. Hue is the wavelength of the colour, value is the tone (from dark to light), and chroma is the colour saturation. The Pycnometer method was used to estimate the particle density of soil using water as the displacing liquid as the soils were salt free and non-swelling type. The water holding capacity (WHC) of the soils was measured in the laboratory using keen- Rackzowski box (Black, 1965). Particle size analyses of soils were estimated by hydrometer method as described by Bouyoucous (1927). The soil pH and electrical conductivity (EC) were recorded in 1:2.5 soil to water suspension (Jackson, 1973). Exchangeable bases were collected using neutral normal ammonium acetate and the exchanged ion measured following the procedure outlined in Hesse (1970). The total Ca⁺⁺ and Mg⁺⁺ was determined by complexometric titration, involving ethylene diamine tetra acetic acid (EDTA). Different soil chemical properties determined were: soil organic carbon (OC) content by chromic acid wet oxidation method (Walkley and Black, 1934), Carbon stocks were determined using the formula,

$$CarbonStock = \frac{OC \times BD \times D \times 10,00}{100}$$

Where Carbon Stock is Mgha⁻¹, OC is the organic carbon expressed as percentage, BD id bulk density in Mgm⁻³ and D is the depth of soil in m.

Available nitrogen (N) was estimatedby alkaline potassium permanganate method (Subbiah and Asija, 1956), available phosphorus (P) spectrophotometrically (Olsen *et al.*,

1954), available potassium (K) by flame photometrically (Hanway and Heidel, 1952), soil available micronutrients iron (Fe), manganese (Mn), copper (Cu) and zinc (Zn) by DTPA-extraction method (Lindsay and Norwell, 1978). Water-soluble ions were determined in 1:2.5 soil to water suspension, whereas, exchangeable ions in neutral normal ammonium acetate extract (Richards, 1954). The data on soil properties were correlated with the Pearson Correlation analysis by colour matrix using R square (R version 3.5.1) to signify the relationship among different soil parameter at different depths.

The collected data were subjected to statistical analysis using analysis of variance and LSD at 5% level with MSTAT computer program according to, Gomez and Gomez (198 ???????????/(comment please suggest me, how can do and may be possible correlation with chemical and physical properties)

3. Results and Discussions

3.1. Physical properties of young alluvial soil

Data in Table No. 1 show that the young alluvial soil of Mirzapur is found in the northern part of the district adjoining the Ganga basin. Soil profile from three locations namely Mirzapur, Chunar, and Chhanbey block was studied to represent and cover the entire stretch of young alluvial soil. The physical properties of young alluvial soil are presented in Table 1 and 2 and the profile photograph of the three profiles presented in figure1 to 3.

There was no variation in soil colour observed at all soil depths in the Chhanbey soil profile. But the Chunar soil profile exhibited a red colour in the plough layer and become yellow-red at lower soil depths. Red, yellow and brown colour is related to the extent of oxidation, hydration and diffusion of iron oxides in the soil. Yellow, red and brown colours are mostly due to the presence of goethite, hematite and magnetite respectively. Organic matter in soil tends to impart dark brown to black colour. So far as texture is concerned, coarse-textured soils are usually light in colour whereas well-textured soil appears darker in colour (Phogat *et al.*, 2015).

Also, Bulk density is the weight of soil in a given volume and is an indicator of ease of root growth. Bulk densities higher than 1.6 g cm^{-3} generally cause restrict root growth. Bulk density increases with compaction and trends to increase with depth. Sandy soil was prone to more bulk density. The bulk density of Mirzapur profile varied between 1.49 and 1.94 Mg m^{-3} and is considered to be quite compact. Higher bulk densities are observed at lower soil depths.

On the other hand, Porosity is generally lower in coarse-textured soil than in fine-textured soil. The porosity of profile soil samples of Mirzapur varied from 26.7 to 43.7

(average 34.9 %) and the corresponding value for Chunar and Chhanbey varied from 30.5 to 56.7 % (mean 42.06 %) and 30.4 – 43.6 % (mean 34.7%) respectively. Thus the overall porosity of young alluvial soils varied from 26.7 – 56.6 % (mean 38.4) which is representative of normal soils. The porosity was always higher near the surface soil layers and decreased with increased soil depth.

The water holding capacity did not vary much among profiles of young alluvial soils. The overall variation in the young alluvial soil profiles of Mirzapur district ranged from 40.5 – 44.4 % (average 42.70%).

It was found that the content of clay varied from 29-43% in Mirzapur profile, 27-45% in profile soils of Chunar and 32-42% in profile soil of Chhanbey with an overall range between 27-45% in young alluvial soils. The silt content varies from 29-45% in Mirzapur, 30-43 % in Chunar and 20-48% in Chhanbey; the overall silt content varying between 29 and 48%. The overall sand content varied from 12-42 % in these soils. The textural class varied between loam to clay loam in Mirzapur, clay to clay loam in Chunar and silty clay loam to clay loam in Chhanbey.

3.2 Physical properties of old alluvial soils

The physical properties of old alluvial soil have been presented in (Table 3 and 4) and a profile photograph of the two profiles presented in figure 4 and 5. The old alluvial soils were located in the northeastern part of the district of Mirzapur. Soil profile from two locations namely Jamalpur and Narayanpur were studied to represent old alluvial soil.

The Jamalpur soil profile (Figure 4) varied in colour between 5Y8/2 and 2.5Y 8/2, Yellow colour indicates the presences of ferric iron oxide. The surface soil at Jamalpur was different in colour from the rest of the profile, which had a lighter colour. In Narayanpur, the soil in the plough layer (0-15 cm) was uniform in colour (5Y8/1) and the soils at lower depth (15-60 cm) were a shade darker in chroma than the surface soils. The profile photograph of Narayanpur has been depicted in figure 6 which clearly show accumulation of organic matter in soil profile.

The bulk density of Jamalpur profile varied between 1.52 and 1.70 Mg m⁻³ and is considered to be quite compact. Higher bulk densities were observed at lower soil depths. Thus the overall porosity of old alluvial soil varied from 34.32 – 42.51 % (mean 37.63) which is representative of normal soils. The water holding capacity did not vary much among profiles of old alluvial soils and varied from 42.70 to 45.50 % (mean 44.16 %).

It was found that the content of clay varied from 17 to 27 % in Jamalpur profile and 16 to 22 % in profile soils of Narayanpur with an overall range between 16-27 % in old

alluvial soils. The silt content varied from 40 to 56% in Jamalpur and 36 to 42 % in Narayanpur; the overall silt content varied between 36 to 56 %. The overall sand content varied from 17 to 43 % in the old alluvial soil profiles. The texture was loam in both Jamalpur and Narayanpur.

3.3 Physico-chemical properties of young alluvial soils.

Three soil profiles were studied to represent young alluvial soils of Mirzapur district, namely Mirzapur, Chunar and Chhanbey (Table 1 and 2) The overall trend in pH of young alluvial soils showed slightly acidic (pH 6.4) to alkaline (pH 9.1) soil reaction. The overall EC in the young alluvial soils of Mirzapur ranged from 0-12 to 0.55 dS m⁻¹, which is non-saline. Hence there is no warning on the growth of all types of crops in these soils.

The organic carbon status of the surface soil layer (0-5cm) was also medium in the category and decreased to the low category with the increase in soil depth. Thus, the young alluvial soils are sufficient in the surface soil layer, but are deficient in the lower soil depths. The decrease in organic carbon with depth has previously been reported by different workers. Bhatnagaret *al.* (2003) reported higher amounts of organic carbon in surface than in subsurface soils as a result of its recycling, over the years by plants and subsequent organic matter accumulation. The exchangeable Ca content in Mirzapur profile varied between 3.25 and 4.8 Cmol (+) kg⁻¹, whereas, the Mg content varied between 3.45 to 7.3 Cmol(+) kg⁻¹. There was enough calcium and magnesium content in the soil to support plant growth.

3.4 Physico-chemical properties of old alluvial soils.

Two soil profiles were studied to represent old alluvial soils of Mirzapur district, namely Jamalpur, and Narayanpur (Table 3 and 4). The overall trend in pH of old alluvial soils was that they were very slightly acidic (pH 6.2) to alkaline (pH 8.6) in reaction. The overall EC in the old alluvial soils of Mirzapur ranged from 0.04 to 0.44 dS m⁻¹, which is non-saline. Hence there is no warning on the growth of all types of crops in these soils. The profile samples of Jamalpur had low soil organic carbon content. The values were 0.33 % in the surface soils and decreased to 0.1% at 60-100 cm depth. Exchangeable calcium and magnesium were extracted using neutral normal ammonium acetate and measured by EDTA titration. The exchangeable Ca content in Jamalpur profile varied between 5.25 to 7.50 C mol (+) kg⁻¹, whereas, the Mg content varied between 3.00 to 3.45 C mol(+) kg⁻¹. There was enough calcium and magnesium content in the soil to support plant growth. The exchangeable Ca content in Narayanpur varied between 2.75 to 5.00 C mol (+) kg⁻¹, whereas the magnesium content varied between 3.90 to 6.85 C mol (+) kg⁻¹.

3.5 Available nutrient status of young alluvial soils.

The available N content in Mirzapur profile (Table 1 and 2) varied from 147-235 kg ha⁻¹ with the mean value of 204 kg ha⁻¹, which can be categorized as low (<280 kg ha⁻¹). The highest available N content was observed at 0-5cm (235kg ha⁻¹) and lowest at 60-100cm (147 kg ha⁻¹) depth. The available N content in the soil profile of Chunar was similar to Mirzapur and varied between 201 and 267 kg ha⁻¹, with a mean value of 236 kg ha⁻¹, which is categorized in the low category of available N. The available N content in Chhanbey varied between 78 and 191 kg ha⁻¹ (mean 144 kg ha⁻¹), which is even lower. Thus, the new alluvial soils have available N in the range of 78-267 kg ha⁻¹ and all soils had low available N status. The available P content in soil profiles of Mirzapur varied from 8-16 kg ha⁻¹ with the mean value of 11 kg ha⁻¹, which falls under the low to medium category. Medium category of available P was noticed at 0-5 cm depth (16 kg ha⁻¹) and the lowest value at 60-100cm (8 kg ha⁻¹) depth. Available potassium content in soils of Mirzapur profile varied between 138-219 kg ha⁻¹ with mean value 186 kg ha⁻¹. The highest available K was noticed at 0-5 cm (219 kg ha⁻¹) and the lowest value at 60-100 cm (138 kg ha⁻¹) depth of soil profile. Among the cationic micronutrients, Zn content varied between 0-2 ppm in the Mirzapur soil profile, 2-9 ppm in the Chunar soil profile and 1-2 ppm in the Chhanbey soil profile. The copper content in soil profile varied between 0-11 ppm in Mirzapur, 2-9 ppm in Chunar and 0-3 ppm in Chhanbey. Available iron content varied between 12-29 ppm in Mirzapur soil profile, 2-11 ppm in the Chunar soil profile and 13-70 ppm in the Chhanbey soil profile. Soils are generally categorized as having sufficient iron if the DTPA extractable Fe content is > 4.5 ppm. Available Mn content in soil profiles varied between 3-27 ppm in Mirzapur, 9-25 ppm in Chunar and 3-26 ppm in Chhanbey.

3.6 Available nutrient status of old alluvial soils.

The available N content in soil profiles of Jamalpur (Table 4) varied from 34 to 154 kg ha⁻¹ with a mean value of 102 kg ha⁻¹, which can be categorized as low (<280 kg ha⁻¹). The highest available N content was observed at 0-5cm (151kg ha⁻¹) and lowest at 60-100cm (34kg ha⁻¹) depth. The available N content in the Narayanpur soil profile was similar to Jamalpur and varied between 69 to 220 kg ha⁻¹, with a mean value of 157 kg ha⁻¹, which was also low in available N. The available P content in soil profiles of Jamalpur varied from 7-11 kg ha⁻¹ with the mean value of 9 kg ha⁻¹, which falls under the low category. Medium category of available P was noticed at 0-5 cm depth (11 kg ha⁻¹) and the lowest value at 60-100cm (7 kg ha⁻¹) depth. Available potassium content in Jamalpur soil profile varied between 173 -299 kg ha⁻¹ with mean value 244 kg ha⁻¹. The highest value was noticed at 5-15 cm (299 kg ha⁻¹) and the lowest value at 60-100 cm (173 kg ha⁻¹) depth of soil profile. Among the

cationic micronutrients, Zn content varied between 1-3 ppm in the Mirzapur soil profile. Copper content varied between 1-7 ppm in Jamalpur, 1-4 ppm in Narayanpur. The available Cu content is categorized as sufficient if it is > 0.2 ppm. Available iron content varied between 2-40 ppm in Jamalpur soil profile and 8-63 ppm in the Narayanpur soil profile. Available Mn content varied between 2-40 ppm in Jamalpur, 5-18 ppm in Narayanpur. Soils are generally categorized as having enough manganese if the DTPA extractable Mn content is > 0.2 ppm. Hence all soils of Jamalpur and Narayanpur had sufficient available soil manganese (Francis *et. al.*, 2019).

3.7 Correlation by colour matrix

Correlation by colour matrix is presented in figure 7. It consists of the correlation of various soil properties variable at various depth (0-100cm) at a different location. The soil profile depths are ranging from 0-5, 5-15, 15-30, 30-60 and 60-100 cm in a different location. The red colour corresponds to (-) negative interaction and blue colour correspond to (+) positive interaction and white correspond to neutral interaction of different soil parameter. In first (A) colour matrix represents the correlation of 0-5 cm depth at a different location. The clay having a positive correlation with porosity as well as water holding capacity (WHC) also showing a positive correlation. In the case of nutrient whereas, the nitrogen having a positive correlation with organic carbon as well as with pH. Most of the nutrients like N, P, Zn and Cu showing a positive correlation with pH. In Second (B) colour matrix represents the correlation of 5-15 cm depth at a different location. Carbon stocks showing the strong positive correlation with bulk density (B.D.), Clay and Organic carbon (O.C.). The nutrients like N, P and Zn showing a negative correlation with physical properties of soil. In Third (C) colour matrix represents the correlation of 15-30 cm depth at a different location. In this correlation the most of nutrient **have** a positive correlation with physical properties of soil. Cu, Zn and Mn **show** strong positive correlation with physico-chemical properties. Major nutrient does not imply any positive or negative relation. In Fourth (D) colour matrix represents the correlation of 30-60 cm depth at a different location. It does not indicate any positive relationships in the broader way at some point its showing negative relation but a few points showing the positive relation like N and Mn having positive relation with chemical properties (pH Soil reaction and Organic carbon). In Fifth (E) colour matrix represents the correlation of 60-100 cm depth at a different location. It clearly shows that few parameters having a positive relation like N and Zn having a positive relationship with physico-chemical properties (Clay, pH, EC and OC). Besides the others nutrients these having a negative correlation with the physical properties of soil.

Conclusion

The Overall Alluvial soils of Mirzapur were mostly soil was dominantly yellow in colour and had high bulk densities, low to medium water holding capacity and porosity. The pH of the soil was acidic to neutral. Whereas, most other soils it was neutral to alkaline. Electrical conductivity was low in all soils and organic carbon and carbon stocks were very poor in most soils. The soils did not have any problem associated with calcium and magnesium nutrition. The available N status was poor and available P status in the range of low to medium and potassium status in the medium range in most soils. All the cationic micronutrients were sufficiently supplied in the soil. Hence specific management options must be exerted for agriculture to be profitable in these soils.

Reference

- Adamo, P., Denaix, L., Terribile, F. and Zampella, M., 2003. Characterization of heavy metals in contaminated volcanic soils of the Solofrana river valley (southern Italy). *Geoderma*, 117(3-4), pp.347-366. [https://doi.org/10.1016/S0016-7061\(03\)00133-2](https://doi.org/10.1016/S0016-7061(03)00133-2)
- Akri, M., Chafik, T., Granger, P., Ayrault, P. and Batiot-Dupeyrat, C., 2016. Novel nickel promoted illite clay based catalyst for autothermal dry reforming of methane. *Fuel*, 178, pp.139-147. <https://doi.org/10.1155/2019/4546350>
- Anlauf, R., Reichel, A., 2014. Effect of aging on the physical properties of landfill cover layers. *Eurasian J. Soil Sci.* 3(3): 212 - 219. DOI: 10.18393/ejss.30326
- Bown, T.M. and Kraus, M.J., 1981. Lower Eocene alluvial paleosols (Willwood Formation, northwest Wyoming, USA) and their significance for paleoecology, paleoclimatology, and basin analysis. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 34, pp.1-30. [https://doi.org/10.1016/0031-0182\(81\)90056-0](https://doi.org/10.1016/0031-0182(81)90056-0)
- Braudeau, E., Frangi, J.P. and Mohtar, R.H., 2004. Characterizing nonrigid aggregated soil water medium using its shrinkage curve. *Soil Science Society of America Journal*, 68(2), pp.359-370. <https://doi.org/10.2136/sssaj2004.3590>
- Chekol, W., Mnalku, A., 2012. Selected Physical and Chemical Characteristics of Soils of the Middle Awash Irrigated Farm Lands, Ethiopia. *Ethiopian Journal of Agricultural Sciences* 22(1): 127-142.
- Daugherty, L.A. and Arnold, R.W., 1982. Mineralogy and Iron Characterization of Plinthitic Soils on Alluvial Landforms in Venezuela 1. *Soil Science Society of America Journal*, 46(6), pp.1244-1252.
- Fasina, A.S., Raji, A., Oluwatosin, G.A., Omoju, O.J. and Oluwadare, D.A., 2015. Properties, Genesis, Classification, Capability and Sustainable Management of Soils from South-Western Nigeria. *International Journal of Soil Science*, 10(3), pp.142-152. <https://doi.org/10.3923/ijss.2015.142.152>

- Feakes, C.R. and Retallack, G.J., 1988. Recognition and chemical characterization of fossil soils developed on alluvium: a Late Ordovician example. *Paleosols and weathering through geologic time: Principles and applications: Geological Society of America Special Paper*, 216, pp.35-48.<https://doi.org/10.1130/SPE216-p35>.
- Fitzsimmons, K.E., Magee, J.W. and Amos, K.J., 2009. Characterisation of aeolian sediment from the Strzelecki and Tirari Deserts, Australia: implications for reconstructing palaeoenvironmental conditions. *Sedimentary Geology*, 218(1-4), pp.61-73.<https://doi.org/10.1016/j.sedgeo.2009.04.004>
- Francis, R., Wuddivira, M.N., Darsan, J. and Wilson, M., 2019. Soil slaking sensitivity as influenced by soil properties in alluvial and residual humid tropical soils. *Journal of Soils and Sediments*, 19(4), pp.1937-1947.<https://doi.org/10.1007/s11368-018-2189-7>
- Gaonkar, O.D., Nambi, I.M. and Govindarajan, S.K., 2019. Soil organic amendments: impacts on sorption of organophosphate pesticides on an alluvial soil. *Journal of soils and sediments*, 19(2), pp.566-578.<https://doi.org/10.1007/s11368-018-2080-6>
- <https://doi.org/10.2136/sssaj1982.03615995004600060025x>
- Iqbal, J., Thomasson, J.A., Jenkins, J.N., Owens, P.R. and Whisler, F.D., 2005. Spatial variability analysis of soil physical properties of alluvial soils. *Soil Science Society of America Journal*, 69(4), pp.1338-1350.<https://doi.org/10.2136/sssaj2004.0154>
- Jiraskova, Y., Bursik, J., Seidlerova, J., Kutlakova, K.M., Safarik, I., Safarikova, M., Pospiskova, K. and Zivotsky, O., 2018. Microstructural analysis and magnetic characterization of native and magnetically modified montmorillonite and vermiculite. *Journal of Nanomaterials*, 2018.<https://doi.org/10.1155/2018/3738106>
- Kauffman, S., Sombroek, W. and Mantel, S., 1998. Soils of rainforests characterization and major constraints of dominant forest soils in the humid tropics. In *Soils of Tropical Forest Ecosystems* (pp. 9-20). Springer, Berlin, Heidelberg.https://doi.org/10.1007/978-3-662-03649-5_1
- Khangarot, A.S. and Mehra, R.K., 1977. Characterization of alluvial soils of Udaipur Valley. *Journal of the Indian Society of Soil Science*, 25(3), pp.247-252.
- Malique, F., Ke, P., Boettcher, J., Dannenmann, M. and Butterbach-Bahl, K., 2019. Plant and soil effects on denitrification potential in agricultural soils. *Plant and Soil*, pp.1-16.<https://doi.org/10.1007/s11104-019-04038-5>
- Manchanda, M.L., Kudrat, M. and Tiwari, A.K., 2002. Soil survey and mapping using remote sensing. *Tropical ecology*, 43(1), pp.61-74.
- Mojid, M., Hossain, A., Wyseure, G., 2018. Relation of reactive solute-transport parameters to basic soil properties. *Eurasian J. Soil Sci.* 7(4): 326 - 336. DOI: 10.18393/ejss.454512

- Mondal, S.U.R.A.J.I.T., Das, T.K., Thomas, P., Mishra, A.K., Bandyopadhyay, K.K., Aggarwal, P. and Chakraborty, D.E.B.A.S.H.I.S., 2019. Effect of conservation agriculture on soil hydro-physical properties, total and particulate organic carbon and root morphology in wheat (*Triticumaestivum*) under rice (*Oryza sativa*)-wheat system. *Indian Journal of Agricultural Sciences* 89, 1, pp.46-55.
- Nguyen, T.T. and Marschner, P., 2016. Sorption of water-extractable organic carbon in various clay subsoils: Effects of soil properties. *Pedosphere*, 26(1), pp.55-61. [https://doi.org/10.1016/S1002-0160\(15\)60022-4](https://doi.org/10.1016/S1002-0160(15)60022-4)
- Obi, J., Udoh, I., Obi, I., 2020. Modelling soil properties from horizon depth functions and terrain attributes: An example with cation exchange capacity. *Eurasian J. Soil Sci.* 9(1): 10 - 17. DOI: 10.18393/ejss.623325
- Prospero, J.M., Ginoux, P., Torres, O., Nicholson, S.E. and Gill, T.E., 2002. Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product. *Reviews of geophysics*, 40(1), pp.2-1. <https://doi.org/10.1029/2000RG000095>
- Sebag, D., Disnar, J.R., Guillet, B., Di Giovanni, C., Verrecchia, E.P. and Durand, A., 2006. Monitoring organic matter dynamics in soil profiles by 'RockEval pyrolysis': bulk characterization and quantification of degradation. *European Journal of Soil Science*, 57(3), pp.344-355. <https://doi.org/10.1111/j.1365-2389.2005.00745.x>
- Shah, B.A., 2010. Arsenic-contaminated groundwater in Holocene sediments from parts of middle Ganga plain, Uttar Pradesh, India. *Current Science(Bangalore)*, 98(10), pp.1359-1365.
- Singh, I.S. and Agrawal, H.P., 2005. Characterization, genesis and classification of rice soils of eastern region of Varanasi, Uttar Pradesh. *Agropedology*, 15(1), pp.29-38.
- Singh, R.N., Singh, R.N.P. and Diwakar, D.P.S., 2000. Characterization of old alluvial soils of Sone basin. *Journal of the Indian Society of Soil Science*, 48(2), pp.352-357.
- Singh, S.K., Dey, P., Singh, S., Sharma, P.K., Singh, Y.V., Latore, A.M., Singh, C.M., Kumar, D., Kumar, O., Yadav, S.N. and Verma, S.S., 2015. Emergence of boron and sulphur deficiency in soils of Chandauli, Mirzapur, SantRavidas Nagar and Varanasi districts of Eastern Uttar Pradesh. *Journal of the Indian Society of Soil Science*, 63(2), pp.200-208. <https://doi.org/10.5958/0974-0228.2015.00026.2>
- Sitanggang, M., Rao, Y.S., Ahmed, N. and Mahapatra, S.K., 2006. Characterization and classification of soils in watershed area of Shikohpur, Gurgaon district, Haryana. *Journal of the Indian Society of Soil Science*, 54(1), pp.106-110.
- Spell, R.L. and Johnson, B.G., 2019. Anthropogenic alluvial sediments in North Carolina Piedmont gullies indicate swift geomorphic response to 18th century land-use

- Sulieman, M.M. and AlGarni, A.M., 2019. Soil organic carbon mapping and prediction based on depth intervals using kriging technique: A case of study in alluvial soil from Sudan. *Eurasian Journal of Soil Science*, 8(1), pp.44-53.
<https://doi.org/10.18393/ejss.492466>
- Tao, W., Lee, M.H., Wu, J., Kim, N.H. and Lee, S.W., 2011. Isolation and characterization of a family VII esterase derived from alluvial soil metagenomic library. *The Journal of Microbiology*, 49(2), pp.178-185.
<https://doi.org/10.1007/s12275-011-1102-5>
- Tsozué, D., Basga, S., Nzeukou, A., 2020. Spatial variation of soil weathering processes in the tropical high reliefs of Cameroon (Central Africa). *Eurasian J. Soil Sci.* 9(2): 92 - 104. DOI: 10.18393/ejss.659830
- Tufa, M., Melese, A., Tena, W., 2019. Effects of land use types on selected soil physical and chemical properties: The case of Kuyu District, Ethiopia. *Eurasian J. Soil Sci.* 8(2): 94 - 109. DOI: 10.18393/ejss.510744
- Valentin, C., 1991. Surface crusting in two alluvial soils of northern Niger. *Geoderma*, 48(3-4), pp.201-222.
[https://doi.org/10.1016/0016-7061\(91\)90045-U](https://doi.org/10.1016/0016-7061(91)90045-U)
- Wadia, B.H. and Daga, A.K., 1977. Parameters for Assessment of Electronic Opportunities. *IETE Journal of Research*, 23(4), pp.184-189.
<https://doi.org/10.1080/03772063.1977.11451310>
- Wakabayashi, S., 2019. Soil Dressing with Alluvial Soil Materials: "Dorotsuke". In *Anthropogenic Soils in Japan* (pp. 147-153). Springer, Singapore.
https://doi.org/10.1007/978-981-13-1753-8_10
- Walker, P.H. and Coventry, R.J., 1976. Soil profile development in some alluvial deposits of eastern New South Wales. *Soil Research*, 14(3), pp.305-317.
<https://doi.org/10.1071/SR9760305>
- Yehia, H.A., Fayed, R.I. and Rateb, K.A., 2013. Integration of GIS and Modeling to Study Soil Characterization, Evaluation and Sensitivity to Degradation of some Alluvial Deposits, Egypt. *Alexandria Science Exchange Journal*, 34(4), pp.460-475.
<https://doi.org/10.21608/ASEJAIQJSAE.2013.3109>
- Yitbarek, T., Jembere, A., Kerebeh, H., 2018. Characterization and classification of soils of Wolkite University research sites, Ethiopia. *Eurasian J. Soil Sci.* 7(4): 292 - 299. DOI: 10.18393/ejss.436186