

Original Research Article
ELECTRICAL, ECHOCARDIOGRAPHIC AND CORONARY ARTERY
CHARACTERISTICS OF RIGHT VENTRICULAR INFARCTION: A SERIES
OF 120 CASES

Introduction: Coronary heart disease is the main cause of morbidity and mortality worldwide. Right ventricular (RV) infarction is often difficult to diagnose and has a poor prognosis due to rhythmic and hemodynamic complications.

Objective: The study of electrical, ultrasonographic, and coronarographic features of the VD Infarction.

Materials and methods: Retrospective study of patients hospitalized in the Cardiology Department of the Mohammed VI University Hospital in Marrakech over a period of 24 months for MDI extended to the VD.

Results: 120 patients were hospitalized during this period for MI with extension to the VD. Atypical clinical presentation was noted in 10% of cases. Clinical examination on admission revealed signs of right heart failure in 18% of cases, including 6% complicated by cardiogenic shock. Thrombolysis was performed in 10% of the patients, 67% of them successfully. The ECG found an isolated extension to the V3R leads in 76% of the cases and in association with a V4R overshoot in 45% of the cases, conduction disorders were noted in 28% of the cases, presented essentially by first degree auriculoventricular block. Echocardiography showed impaired LV function in 82% of cases, and longitudinal systolic dysfunction of the LV in 65%. Coronary angiography was performed in 91% of the cases, half of which underwent coronary angioplasty. The combination of both DC and IVA damage was found in 40% of the cases, and damage to the middle DC was the most common in almost half of the cases. The most frequent complications were rhythmic and conductive disorders in 38% of cases, and the evolution was fatal in 8% of cases.

Conclusion: Involvement of the VD during MI is characterized by a very critical initial phase, once overcome, the overall prognosis becomes favorable in the long term.

Keywords: Infarction, Right Ventricle, Right Bypass, Right Coronary, Therapeutic Management

INTRODUCTION:

Right ventricular infarction (RV MI) is a non-rare entity, complicating 40-50% of inferior myocardial infarctions, with higher hospital morbidity and mortality related to hemodynamic and rhythmic complications. The culprit lesion is often an occlusion of the right coronary artery. However, the DV is relatively resistant to infarction and recovers even after prolonged occlusion, as well as the diagnosis of ischemic damage to the DV is often based on the presence of right clinical signs and/or electrical signs in the right leads (1).

The aim of this work is to study the electrical, ultrasonographic, and coronary artery characteristics of MDI with DV extension, as well as its complications and therapeutic modalities.

MATERIALS AND METHODS:

This is a retrospective and descriptive mono-centric study spread over a period of 24 months, covering patients hospitalized in the Department of Cardiology and Vascular Diseases at the Mohammed VI Hospital Center of Marrakech between September 2019 and October 2021, for left heart myocardial infarction extended to the right ventricle.

The records of patients hospitalized with the diagnosis of MDI extended to the RV were indexed from the registers and archives of the department.

RESULTS:

120 patients were hospitalized during this period for MI with extension to the VD. The mean age of our patients was 63 years, with extremes ranging from 40 to 80 years. The peak frequency was between 60 and 70 years, with a sex ratio that was equal to 2.33. In our series, more than 75% of the patients had a high cardiovascular risk. Atypical clinical presentation was noted in 10% of cases (Figure 1).




Figure 1: Distribution according to initial symptomatology

Clinical examination on admission found signs of right heart failure in 18% of cases, of which 6% were complicated by cardiogenic shock (Figure 2). Thrombolysis was performed in 10% of the patients, 67% of them successfully.




Figure 2: Distribution according to physical signs

The ECG found an isolated extension to the V3R leads in 76% of the cases and in association with a V4R overshoot in 45% of the cases, conduction disorders were noted in 28% of the cases, presented essentially by first degree auriculoventricular block (table 1).

UNDER PEER REVIEW

ECG abnormalities		Number of cases	Percentage
Repolarization disorder: <ul style="list-style-type: none"> ▪ Sus offset in V3R (80%) ▪ Its offset in V4R (41%) 	Lower	84	70%
	ASA	4	3%
	Extended anterior	16	13%
	V1, V2, V3	6	5%
	AvR	2	2%
	Circumferential	8	7%
Rhythm disorder (10%)	FV	2	2%
	FA	4	3%
	ESV	4	3%
	ESSV	2	2%
Conduction disorder (28%)	BBG	2	2%
	HBAG	4	3%
	Bi-fascicular block	4	3%
	BAV first degree	16	13%
	Full BAV	8	7%

Table 1: Distribution of patients according to ECG abnormalities

Echocardiography showed impaired LV function in 82% of cases, and longitudinal systolic dysfunction of the VD in 65% based on the TAPSE found <17 in 40% of patients and S'VD <9.5 noted in 45% of cases. Other abnormalities were found on TTE to note; LV abnormalities in 22% of cases, dilatation of the IVC (30%), apical thrombus and pericardial effusion (8%).

Coronary angiography was performed in 91% of the cases, half of which required coronary angioplasty; the combination of both DC and IVA damage was found in 40% of the cases, and middle DC damage was the most common in almost half of the cases (Figure 3,4).




Figure 3: Anomalies on coronary angiography.




Figure 4: The distribution of the segments of the right coronary reaches

The most frequent complications were rhythmic and conductive disorders in 38% of cases, and the evolution was fatal in 8% of cases.

DISCUSSION:

DV infarction is a cause of arterial hypotension related to the sudden decrease in right ventricular systolic ejection. Immediate mortality is high in the absence of adequate treatment (filling, deobstruction, etc.). The ST elevation in the right leads considered in several studies (1,2) as the gold standard in the diagnosis of VD MI is indeed a transient sign that can be absent in half of the patients with VD MI 10 to 12 hours after the onset of pain (3). Hence the interest of a careful echocardiographic examination with an accurate assessment of the systolo-diastolic function of the VD.

The study of the systolic function of the VD involves several parameters according to the new recommendations (4), no parameter alone could conclude the stage of the systolic dysfunction of the VD, some parameters are considered as little or not valid as the systolic ejection fraction according to the Simpson method, on the contrary the fraction of shortening of the VD, the analysis of the longitudinal function by the TAPSE or the peak of the S wave velocity in tissue Doppler are the most retained and validated parameters for the determination of the systolic function, it depends in fact on the underlying heart disease to be studied(5).

DV infarction occurs primarily on DC occlusion in the setting of inferior myocardial damage, it may also occur due to CX occlusion in patients with predominantly left circulation, and less frequently, associated LAI involvement in cases of anterior DV MI(6).

The best treatment of right ventricular infarction remains, in addition to volume expansion, the earliest deobstruction of the thrombosed artery by medicinal means (although the rate of re-occlusion is higher) or instrumental means (primary angioplasty), which may suffice to regress, rapidly, all the clinical symptomatology and ensure a spectacular improvement of the prognosis (7).

Early recognition of DV infarction in patients with acute myocardial infarction is paramount, not only for prognostic purposes, but also to guide specific therapy, in order to avoid treatments that would further reduce right ventricular preload, thus compromising the patient's condition (3).

CONCLUSION :

The diagnosis of DV MI can be difficult; the 12-lead ECG remains the main diagnostic tool in the acute setting, but the results may be transient.

The evolution of DV MI is characterized by a very critical initial phase, dominated by rhythmic and hemodynamic consequences. Once overcome, the overall prognosis becomes favorable in the long term. Revascularization remains the cornerstone in the management of DV MI.

REFERENCE:

1. Wang JJ, Pahlm O, Warren JW, Sapp JL, Horáček BM. Criteria for ECG detection of acute myocardial ischemia: Sensitivity versus specificity. *J Electrocardiol.* 2018; 51(6): S12-S17.
2. Manvi R Nagam, David R Vinson, Joel T Levis « ECG Diagnosis: Right Ventricular Myocardial Infarction ». *The Permanente Journal/Perm J* 2017;21:16-105
3. Haddad F, Hunt SA, Rosenthal DN, MurphyDJ. Right ventricular function in cardiovascular disease, part I: anatomy, physiology, aging, and functional assessment of the right ventricle. *Circulation* 2008;117:1436-48.
4. Webner C. ECG Identification of Right Ventricular Myocardial Infarction. *AACN Adv Crit Care.* 15 déc 2019;30(4):425-31.
5. Kuznetsova NS, Rabinovich RM, Mazur VV, Mazur ES. [Diagnostic difficulties of isolated right ventricular myocardial infarction]. *Kardiologija.* 30 sept 2021;61(9):66-70.

6. Tadic M, Cuspidi C, Versaci F, Calcagno S. Right ventricular infarction: can we still use old tricks? *Minerva Cardiol Angiol.* oct 2021;69(5):499-501.

7. J. Sanz, Damián Sánchez-Quintana, E. Bossone, Harm J. Bogaard, R. Naeije, « Anatomy, Function, and Dysfunction of the Right Ventricle » ; *JOURNAL OF THE AMERICAN COLLEGE OF CARDIOLOGY* VOL . 73 , NO. 12 , 2019

UNDER PEER REVIEW