

Review Article

Role of Selenium in Ruminants Health and Reproduction

Abstract

Selenium (Se) which was earlier classified as a toxic element, has now been proved to be an essential mineral required for proper health, immunity, and reproductive functions of animals. Se status of animals and plants varies markedly around the world as a result of different geological conditions and the concentrations of Se in plant material are highly correlated with those in the soil. Se deficiency has been linked to many health problems in young animals such as increased neonatal mortality, decreased sucking reflex, weakness, higher occurrence of infectious diseases and white muscle disease. In recent years, there has been considerable interest in the use of Se in ruminant diets. Different forms of Se routinely used for dietary supplementation are inorganic, organic and nano Se.

Keywords: Selenium, neonatal mortality, energy metabolism, animal health

Introduction

Minerals are inorganic substances that cannot be synthesized in body, but are required in small amount to support different metabolic functions of body [1]. 22 minerals are needed for animals. Out of these minerals, 15 are micro minerals and 7 macro minerals [2]. Trace minerals, though required in small quantities (less than 100 mg/kg dry matter) have critical roles in immune function, oxidative and energy metabolism in ruminants which are directly or indirectly involved in growth, production and reproduction [3]. They include cobalt, copper, iodine, iron, manganese, molybdenum, Se and zinc [4]. Among these, Se is currently acknowledged to be an essential dietary trace element required for various body functions such as growth, reproduction, immune system and protection of tissue integrity [5]. Some minerals such as Nickel, Chromium, arsenic and Vanadium are known to clearly have a role in animal health but the exact biochemical nature is unknown and its essentiality have weak evidences [6] [7] [8] [9].

Status of Selenium

The Se status of animals and plants varies markedly around the world as a result of different geological conditions and the concentrations of Se in plant material are highly correlated with those in the soil. Agricultural production system in developing countries is under pressure to fulfil the requirement of growing population, which has led to indiscriminate use of fertilizers resulting in severe deficiency of micro minerals in soil. The deficiency of Se in soil and crop plants has been reported in many countries like India, China, Turkey and Pakistan [10]. The concentrations of Se in plant material are highly correlated with those in the soil. Fertilization of soil with Se increases Se concentrations in plants [11]. The Se status in animals and plants varies markedly around the world as a result of different geological conditions. High Se

Comment [၁၁1]: Role of Se in reproduction should be added.

Comment [၁၁2]: The information of the important of Se in ruminant should be provided.

Comment [၁၁3]: Status of Se in ruminant would be more interesting topic.

concentrations are associated with some phosphatic rocks, organic rich black shales, coals, and sulphide mineralization, whereas most other rock types contain very low concentrations. Globally Se deficient soils are far more widespread than are seleniferous ones [12]. Animal health is affected by Se deficiency or excess in the diet, the intake of Se being dependent on the amount of Se taken up by plants as bioavailable Se [13].

Chemical Nature

Se is a naturally occurring metalloid element that is essential to human and animal health in trace amounts but is harmful in excess. It was first identified in 1817 by the Swedish chemist, Jons Jakob Berzelius. It has chemical and physical properties intermediate between metals and non-metals and is similar to those of sulphur, arsenic and tellurium, all of which are in Group VI of the periodic chart of the elements [14]. Like Sulphur, Se can exist in different oxidation states as selenide (2-), Se (0), selenite (4+) and selenate (6+), respectively. It behaves antagonistically with Copper and Sulphur in humans and animals inhibiting the uptake and function of these elements [15].

Essentiality of Selenium

Se which was earlier classified as a toxic element, has now been proved to be an essential mineral required for proper health, immunity, and reproductive functions of animals. The essentiality of Se was proved for the first time from the work that liver necrosis in rats [16]. and exudative diathesis in chicks could be prevented by supplements of Se. Se is required to maintain normal physiological functions and provides a significant dietary source of antioxidant defenses [17]. It is present in all cells and tissues and is necessary for maintaining the vital functions of humans and animals. The content of Se in the organism is naturally very low, the majority of Se being bound in tissues and blood in the form of selenoproteins. It is a component of at least 25 selenoproteins with antioxidant, anti-inflammatory and chemoprotective properties [18]. The most important are glutathione peroxidases (GSH-Px-1, GSH-Px-6), thioredoxin reductases (TrxR1–TrxR3), iodothyronine deiodinases (ID1–ID3), selenophosphate synthetase, Selenop, and selenoprotein W. This element acts as a cofactor of the GSH-Px family of enzymes which protect against oxidative stress. Specifically, Se-dependent GSH-Px enzyme recycles glutathione, reducing lipid peroxidation by catalysing the reduction of peroxides, including hydrogen peroxide. In general, all these enzymes in their reduced state catalyse the breakdown of lipid hydroperoxides and hydrogen peroxides in cells [19]. GSH-Px and selenoprotein P are also involved in the regulation of the inflammatory response.

Important role of selenium in health and reproduction

The ability of Se to improve the immune response in farm animals is well documented [20]. Se deficiency has been reported to decrease humoral immune response and neutrophil killing activity in cattle without any clinical signs [21]. Moreover, [22]. reported that dietary Se supplementation increases antibody titre, neutrophils killing activity and reduces morbidity and mortality in beef cattle. Se improves phagocytosis in white blood cell populations. It might have been a boost in the passive immunity by enhancing immunoglobulin G (IgG) absorption in the new born lamb [23]. The improvement of immune cell functioning is likely due to the enhanced antioxidant status [24]. Se performs significant functions in the male reproductive systems which are regulated by selenoproteins, especially GSH-Px4 (Glutathione peroxidase) and SELENOP (Selenoprotein-P) (Qazi et al., 2019).

Comment [๑๑4]: This basic information of Se is not necessary. It has been intensively reviewed.

Comment [๑๑5]: This is the main topic of the manuscript. Presentations of the data of literature review papers in the forms of Table and/or Figure will be useful.

GSH- Px4 is distinctly expressed in testes and has both an antioxidant as well as a structural role; the latter context is evident from a fact that it constitutes over 50% of mitochondrial capsule (as an oxidatively inactivated protein) in midpiece of mature sperm [25]. In an early stage of spermatogenesis, GSH-Px4 is believed to protect the developing sperm from oxidative stress-induced DNA damage, however, in the later phase, through cross linkage with proteins in midpiece region, it provides the integrity to the sperm midpiece by becoming a structural component of mitochondrial sheath circumventing the flagellum, which is an essential component for sperm stability and motility.

SELENOP serves as a transport protein for Se and is also expressed in vesicle like structures in the basal region of the Sertoli cells [26]. Male reproductive tissues are constantly exposed to reactive oxygen species (ROS) generated as products of normal metabolism, and during the fertilization process spermatozoa normally pass through an area of high oxygen level [27]. Mammalian spermatozoa membranes have a high content of polyunsaturated fatty acids and are highly susceptible to lipid peroxidation caused by reactive oxygen species (ROS) overproduction [28]. ROS induced damages decreased both semen quality and sperm function. High ROS concentrations lead to pathologic changes in sperm cells by intensifying lipid peroxidation, which eventually results in their motility and viability loss [29]. Several defence mechanisms including antioxidant help in counteracting ROS detrimental effects and maintain sperm motility and viability [30]. The activity of the cytoplasmic antioxidant enzymes in sperm cells is very low [31], and the small amount of the cytoplasm in their heads and tails makes them susceptible to oxidative stress. Unlike spermatozoa, seminal plasma is a vital source of antioxidants, including glutathione peroxidase (GSH-Px), superoxide dismutase (SOD), uric acid and vitamin E [32]. Activity of Se-dependent phospholipid hydroperoxide GSH-Px (PH-GSH-Px, GSH-Px4) is very pronounced in late (meiotic) spermatogenic cells, where it acts as a structural protein in sperm heads, whereas the nuclear form GSH-Px4 contributes to chromatin condensation [33].

Se also helps in testicular growth and development of seminiferous tubules, spermatogenesis, steroidogenesis in testes, synthesis and secretion of follicular stimulating hormone (FSH) and luteinizing hormone (LH) [34]. Supplementation of Se has been found to improve semen quality by increasing antioxidative defence of seminal plasma in buck [35], boar [36], Boer goats [37], ram [38] and cockerels [39]. Se deficiency has been linked to many health problems in young animals such as increased neonatal mortality, decreased sucking reflex, weakness, higher occurrence of infectious diseases and white muscle disease [40]. It may also result in immune and endocrine disorders, especially thyroid dysfunction [41].

Supplementation sources and its bioavailability

The bioavailability of Se is associated with its forms. Generally, Se is utilized as inorganic, organic and nano forms. Form of Se supplementation has an important bearing on its possible ameliorative effects and/or on general wellbeing and development of organisms [42]. Both form and the total intake of Se are equally important with regards to the potential health-related effects. It has been reported that organic forms of Se have greater bioavailability compared to the inorganic forms [43], because sodium selenite and sodium selenate must first be converted to hydrogen selenide and then to selenophosphate before they can be utilised in selenoprotein synthesis [44]. The most common inorganic Se sources are sodium selenite (SS) and sodium selenate, which are usually provided in mineral premixes or injected. Organic Se sources are seleno-amino acids (e.g. selenomethionine (Se-Met) and selenocysteine), which are found in Se yeast or in feeds grown on Se-rich soils. Recently,

Comment [336]: This topic should be stressed on the effects of Se sources on health and reproduction of ruminant.

elemental nano-Se has attracted a wide spread attention to its high bioavailability and low toxicity[45] [46]. Nano minerals improve the bioavailability due to its novel characteristics such as high surface activity, a lot of surfaces active centers, strong adsorbing ability and high catalytic efficiency [47]. Nano-Se has efficient functions on animal growth, reproduction and immunity systems [48]. In sheep, Nano- Se had improved ruminal fermentation, nutrient digestibility [49]. In addition, some reports on rats and mice demonstrated that Nano-Se had higher efficiency than sodium selenite and other Se sources in up-regulating selenoenzymes, exhibiting lower toxicity [50]. Subsequent studies also pointed out that Nano-Se has more beneficial effects to improve activity of glutathione peroxidase, blood biochemical indices with lower toxicity comparing with organic or inorganic Se sources[51].

Conclusions:

Adequate essential trace mineral intake and absorption is required for a variety of metabolic functions including immune response, reproduction and growth. Se is recognized as an essential trace element, and its deficiency has been associated with impaired growth, fertility, and health in farm animals. Diets for ruminant animals are often of plant origin and the Se concentration within plants can be extremely variable. Consequently, concentration of dietary Se can be deficient, and Se supplementation may be required. Supplementation of trace elements in animal diets has long been practiced to ensure optimum growth production and improve immune response. In recent years, there has been considerable interest in the use of Se in ruminant diets. Different forms of Se routinely used for dietary supplementation are inorganic, organic and nano Se. There are reports of improved growth, better production performance, reproduction and health in ruminants fed inorganic, organic and nano Se.

REFERENCES:

1. Eruvbetine D. 2003. Canine Nutrition and Health. A paper presented at the seminar organized by Kensington Pharmaceuticals Nig. Ltd, Lagos.
2. Underwood EJ, Suttle NF. 1999. The mineral nutrition of livestock. CABI Publication, New York, USA.
3. Bhalakiya N, Haque N, Patel P, Joshi P. 2019. Role of Trace Minerals in Animal Production and Reproduction. International Journal of Livestock Research, 9(9):1-12
4. Soetan KO, Olaiya CO, Oyewole OE. 2010. The importance of mineral elements for humans, domestic animals and plants-A review. African journal of food science, 4(5):200-222.
5. Pilarczyk B, Tomza-Marciniak A, Dobrzański Z, Szewczuk M, Stankiewicz T, Gaczarzewicz D, Lachowski W. 2013. The effect of selenized yeast supplementation on some performance parameters in sheep. Turkish Journal of Veterinary and Animal Sciences, 37(1):61-67.
6. Keshri A, Roy D, Kumar V, Kumar M, Kushwaha R, Vaswani S. Impact of different source chromium sources on physiological response, blood biochemicals and endocrine status of heat stress in dairy calves. Biological Rhythm Research. 2022; 53(1):58-69.
7. Singh A, Kumar M, Kumar V, Roy D, Kushwaha R, Vaswani S. Effect of Nickle supplementation on antioxidant status, immune characteristics and energy and lipid metabolism in growing cattle. Biological Trace element research, 2019; 190(1):65-75.

Comment [337]: Conclusion should contain the information of the role of Se (or Se sources) on health and production of ruminant.

8. Vaswani S, Mani V, Kewalramani N, Kaur H. 2010. Mitigation of adverse effects of arsenic by supplementing vitamin E in crossbred kids maintained at low protein diet. *Indian journal of Animal Nutrition*. 27(4): 346-352.
9. Praveen Kumar Gupta, Shalini Vaswani, Vinod Kumar, Debashis Roy, Muneendra Kumar, Raju Kushwaha, Avinash Kumar, Amit Shukla. Investigations on Modulating Effect of Vanadium Supplementation on Growth and Metabolism Through Improved Immune Response, Antioxidative Profile and Endocrine Variables in Haryana heifers. *Biological trace element research*. 194: 379-389.
10. Gupta PK and Vaswani S. 2020. Basic information about vanadium 'ultra-trace element or occasionally beneficial element' and its various functions in animals: A review article. *J. Entomol. Zool. Stud.*, 8:645-53
11. Rashid A, Ryan J. 2008. Micronutrient constraint to crop production in the near east potential significance and management strategies. *Micronutrient deficiencies in global crop production*, DOI:10.1007/978-1-4020-6860-7_6
12. National Research Council. 2001. *Nutrient requirements of dairy cattle*. National Academies Press. Washington, DC, USA.
13. Khanal DR, Knight AP. 2010. Selenium: Its Role in Livestock Health and Productivity. *Journal of Agriculture and Environment*, 11:101-106.
14. Fordyce F. 2005. Selenium deficiency and toxicity in the environment. In 'Essentials of medical geology—Impacts of the natural environment on public health' (Eds O Selinus B Alloway JA Centeno RB Finkelman R Fuge U Lindh P Smedley) pp. 373-415.
15. Verma AK, Kumar A, Rahal A, Kumar V, Roy D. 2012. Inorganic versus organic selenium supplementation: a review. *Pakistan journal of biological sciences*, 15(9):418-425.
16. Netto AS, Zanetti MA, Correa LB, Del Claro GR, Salles MSV, Vilela FG. 2014. Effects of dietary selenium sulphur and copper levels on selenium concentration in the serum and liver of lamb. *Asian-Australasian journal of animal sciences*, 27(8):1082
17. Patterson EL, Milstrey R, Stokstad ELR. 1957. Effect of selenium in preventing exudative diathesis in chicks. *Proceedings of the society for experimental Biology and Medicine*, 95(4):617-620.
18. Sordillo LM. 2013. *Selenium-dependent regulation of oxidative stress and immunity in periparturient dairy cattle*. Hindawi Publishing Corporation. *Veterinary Medicine International*.
19. Pappas AC, Zoidis E, Surai PF, Zervas G. 2008. Selenoproteins and maternal nutrition. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 151(4):361-372.
20. Navarro-Alarcon M, Cabrera-Vique, C. 2008. Selenium in food and the human body: a review. *Science of the total environment*, 400(1-3):115-141.
21. Chauhan SS, Celi P, Ponnampalam EN, Leury BJ, Liu F, Dunshea FR. 2014. Antioxidant dynamics in the live animal and implications for ruminant health and product (meat/milk) quality: role of vitamin E and selenium. *Animal Production Science*, 54(10):1525-1536.
22. Hogan JS, Smith KL, Weiss WP, Todhunter DA, Schockley WL. 1990. Relationships among vitamin E, selenium, and bovine blood neutrophils. *Journal of Dairy Science*, 73(9):2372-2378

23. Hall JA, Bobe G, Vorachek WR, Gorman ME, Mosher WD, Pirelli GJ. 2013. Effects of feeding selenium-enriched alfalfa hay on immunity and health of weaned beef calves. *Biological Trace Element Research*, 156(1-3):96-110.
24. Rock MJ, Kincaid RL, Carstens GE. 2001. Effects of prenatal source and level of dietary selenium on passive immunity and thermo metabolism of newborn lambs. *Small Ruminant Research*, 40(2):129-138.
25. Rossi CS, Compiani R, Baldi G, Muraro M, Marden JP, Rossi R, Pastorelli G, Corino C, Dell'Orto V. 2017. Organic selenium supplementation improves growth parameters, immune and antioxidant status of newly received beef cattle. *Journal of Animal and Feed Sciences*, 26(2):100-108.
26. Foresta C, Flohe L, Garolla A, Roveri A, Ursini F, Maiorino M. 2002. Male fertility is linked to the selenoprotein phospholipid hydroperoxide glutathione peroxidase. *Biology of reproduction*, 67(3):967-971.
27. Olson GE, Winfrey VP, Nagdas SK, Hill KE, Burk RF. 2007. Apolipoprotein E receptor-2 (ApoER2) mediates selenium uptake from selenoprotein P by the mouse testis. *Journal of Biological Chemistry*, 282(16):12290-12297.
28. Fujii J, Iuchi Y, Matsuki S, Ishii T. 2003. Cooperative function of antioxidant and redox systems against oxidative stress in male reproductive tissues. *Asian Journal of Andrology*, 5(3):231-242.
29. Aitken RJ, Buckingham D, Harkiss D. 1993. Use of a xanthine oxidase free radical generating system to investigate the cytotoxic effects of reactive oxygen species on human spermatozoa. *Reproduction*, 97(2):441-450.
30. Makker K, Agarwal A, Sharma R. 2009. Oxidative stress and male infertility. *Indian Journal of Medicine Research*, 129(4):357-67.
31. Aitken RJ, Baker MA. 2004. Oxidative stress and male reproductive biology. *Reproduction, Fertility and development*, 16(5):581-588.
32. Bilodeau JF, Chatterjee S, Sirard MA, Gagnon C. 2000. Levels of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Molecular Reproduction and Development: Incorporating Gamete Research*, 55(3):282-288.
33. Raijmakers MT, Roelofs HM, Steegers EA, Mulder TP, Knapen MF, Wong WY, Peters WH. 2003. Glutathione and glutathione S-transferases A1-1 and P1-1 in seminal plasma may play a role in protecting against oxidative damage to spermatozoa. *Fertility and sterility*, 79(1):169-172.
34. Flohe L. 2007. Selenium in mammalian spermiogenesis. *Biological chemistry*, 388(10):987-995.
35. Bedwal RS, Bahuguna A. 1994. Zinc copper and selenium in reproduction. *Experientia*, 50(7):626-640.
36. Shi L, Zhang C, Yue W, Shi L, Zhu X, Lei F. 2010. Short-term effect of dietary selenium-enriched yeast on semen parameters, antioxidant status and Se concentration in goat seminal plasma. *Animal feed science and technology*, 157(1-2):104-108.
37. Marin-Guzman J, Mahan DC, Pate JL. 2000. Effect of dietary selenium and vitamin E on spermatogenic development in boar. *Journal of animal science*, 78(6):1537-1543.
38. Li-Guang S, Ru-Jiea Y, Wen-Bina Y, Wen-Juana X, Chun-Xianga Z, You-Shea R, Lei AS, Fu-Linb L. 2010. Effect of elemental nano-selenium on semen quality, glutathione peroxidase activity, and testis ultrastructure in male Boer ram. *Animal Reproduction Science*, 118:248-254.
39. Kendall N, McMullen S, Green A, Rodway R. 2000. The effect of a zinc cobalt and selenium soluble glass bolus on trace element status and semen quality of ram lambs. *Animal Reproduction Science*, 62(4):277-283.

40. Ebeid TA. 2012. Vitamin E and organic selenium enhances the antioxidative status and quality of chicken cockerel semen under high ambient temperature. *British Poultry Science*, 53(5):708-714.
41. Enjalbert F, Lebreton P, Salat O. 2006. Effects of copper, zinc and selenium status on performance and health in commercial dairy and beef herds:retrospective study. *Journal of Animal Physiology and Animal Nutrition*, 90(11-12):459- 466.
42. Kohrle J, Jakob F., Contempre B, Dumont JE. 2005. Selenium, the thyroid, and the endocrine system. *Endocrine reviews*, 26(7):944-984.
43. Pappas AC, Zoidis E, Surai PF, Zervas G. 2008. Selenoproteins and maternal nutrition. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 151(4):361-372.
44. Burk RF, Hill KE. 2015. Regulation of selenium metabolism and transport. *Annual review of nutrition*, 35:109-134.
45. Kachuee R, Moeini MM, Souri M. 2013. The effect of dietary organic and inorganic selenium supplementation on serum Se Cu Fe and Zn status during the late pregnancy in Merghoz goats and their kids. *Small Ruminant Research*, 110(1):20-2
46. Xu DX, Shen HM, Zhu QX, Chua L, Wang QN, Chia SE, Ong CN. 2003. The associations among semen quality, oxidative DNA damage in human spermatozoa and concentrations of cadmium, lead and selenium in seminal plasma. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 534(1-2):155-163.
47. Zhang J, Wang X, Xu T. 2008. Elemental selenium at nano size (Nano-Se) as a potential chemopreventive agent with reduced risk of selenium toxicity: comparison with se-methyl selenocysteine in mice. *Toxicological sciences*, 101(1):22-31.
48. Wang Y, Yan X, Fu L. 2013. Effect of selenium nanoparticles with different sizes in primary cultured intestinal epithelial cells of crucian carp, *Carassius auratus gibelio*. *International Journal of Nanomedicine*, 8:4007.
49. Shi L, Xun W, Yue W, Zhang C, Ren Y, Liu Q, Wang Q, Shi L. 2011. Effect of elemental nano-selenium on feed digestibility, rumen fermentation, and purine derivatives in sheep. *Animal Feed Science and Technology*, 163 (2-4):136-142.
50. Zhang JS, Gao XY, Zhang LD, Bao YP. 2001. Biological effects of a nano red elemental selenium. *Biofactors*, 15(1):27-38.
51. Yaghmaie PA, Ramin S, Asri-Rezaei A, Zamani. 2017. Evaluation of glutathione peroxidase activity trace minerals and weight gain following administration of selenium compounds in lambs. *Veterinary Research Forum*, 8(2):133 – 137.