

GENETICS AND GENOMICS OF AROMA OF RICE AND ITS QUALITY ENHANCEMENT PARAMETERS: A REVIEW

ABSTRACT

Rice (~~*Oryza sativa*~~RYZA SATIVA L.) is a crucial cereal crop because it provides essential vitamins to more than 50% of the world's population. Almost 90% of the world's rice output is produced in Asia, particularly South Asia, where its cultivation is predominantly focused. People from different regions of the world have varying tastes in rice. The scent is influenced by more than 500 volatile chemical components. The most crucial for rice flavors among them has been found as 2 acetyl-1-pyrroline (2AP). It is observed that the fragrance characteristic is correlated with three quantitative trait loci (QTLs). The most important QTL for fragrance is called Qar8.1, and it is located on chromosome 8. The identification of QTLs in the genome uses a variety of markers, including molecular markers and DNA markers.

Formatted: Font: Italic

Keywords: aromatic rice, 2 acetyl-1-pyrroline (2AP), quantitative trait loci (QTLs), volatile organic compounds(VOCs).

1. INTRODUCTION

Rice, which comes from the plant genus *Oryza sativa* L. and is more generally known as rice, is an essential source of vitamins and proteins that contribute to proper nutrition for more than half of the world's population. This multipurpose crop has a high concentration of a wide variety of minerals, including thiamine, riboflavin, niacin, vitamin E, zinc, potassium, and iron.[1]. In East African marketplaces, the predominant aspect that motivates purchasers is aroma, despite the fact that consumers' preferences about the quality of the grain are a matter of personal taste. Hence, rice farmers all over the globe who plant aromatic rice varieties may gain large profits from the unique scent of their product due to the fact that aromatic rice types have a higher demand.[2]. In 2018, India was the top exporter of basmati rice. Genetic factors influence rice fragrance, but environmental factors also play a part. Asia, especially South Asia, is the region where rice is grown for over 90% of the world's rice production. Both biotic and abiotic factors can affect rice production, although drought stress is the main one that affects rainfed areas.[3] Additionally, due to drought, climate change has a negative association with rice output. Consequently, it is essential to cultivate drought-tolerant cultivars. In the majority of Asian nations, rice serves as a significant source of protein. Six of the 27 species in the *Oryza* genus have diploid genomes (n = 12: AA, BB, CC, EE, FF, and GG), while five have polyploid genomes (n = 24: BBCC, CCDD, HHJJ, HHKK, and KKLL). More over half of the world's population eats rice as a staple diet, making it the most significant cereal crop[4]. Rice is divided into aromatic and non-aromatic varieties based on whether an aroma is present[5]. Non-aromatic rice varieties are typically developed in all nations that cultivate rice and are high yielding, show strong agronomic performance, and are highly adaptable to the environmental conditions. Contrarily, the majority of types of aromatic rice genotypes are produced in a small number of nations, have low yields, poor agronomic performance, and are very susceptible to environmental conditions [6]. The fragrant rice is widely praised for its outstanding scent and superior grain quality despite its subpar performance[7]. According to phylogenetic study, the aroma genes are derived from the wild cousins (*Oryza nivara* and *O. rufipogon*), with the Indian sub-continent serving as their primary region of origin before from where they spreading to other areas of the world[8]. Natural selection played a major role in the evolution of aromatic rice varieties over thousands of years. Local farmers then continued this process by selecting varieties that suited their preferred cultivation methods. Finally, plant breeders continued this process over time by making improvements[9]. As a result, it is possible to view the scent of rice as a natural occurrence that has been supported and maintained by the selection process.

Comment [JY1]: This statement is out of place. The paragraph is about the importance of rice and not exportation of rice.

Comment [JY2]: This should be the beginning of another paragraph

Comment [JY3]: Relocate this to the nutritional benefits of rice in the first paragraph

Comment [JY4]: This should be the beginning of another paragraph

Comment [JY5]: Relocate to the first paragraph on the importance of rice

Formatted: Font: Italic

Formatted: Font: Italic

1.1 PREFERENCES OF RICE CONSUMERS AND AROMA OF RICE

FRAGRANCE

1.1.1.2 Preferences of Rice Consumers Preferences

People from many regions of the world have varying tastes in rice. Regardless of the physical and chemical properties of rice, low-income individuals prefer inexpensive rice to Basmati in general[8]. However, Muhammad's research in Indonesia shows that wealthy people prefer rice with a nice flavor[10]. Due of to the fragrance, rice prices are impacted. The Ugandan population prefers rice that is white in colour, whole grained, and aromatic. Rwandan urban residents, however, favour prefer long-grained, fragrant rice. Tanzanian consumers like eating the nation's flavorful indigenous rice[8]. Additionally, rural residents and employers of police and military personnel's employers consume rice of lowest possible quality [11]. Consumers in Western nations favour preferred rice with a fluffy texture and a softer scent, but those in Asian countries like preferred rice with a sticky texture and a strong perfume.

Comment [JY6]: What do you mean by this word?

Overall, research indicates that a range of elements, such as flavour, texture, scent, and cooking method, affect customer preferences for rice. Producers and marketers of rice may better match-meet the demands and expectations of their target customers by being aware of these preferences.[12]

Comment [JY7]: You mean cooking time?

Comment [JY8]:

4-21.3 AROMA OF RICE

Several civilizations throughout the world place a great emphasis on rice's scent as an essential quality. When rice is cooked, a variety of volatile organic compounds (VOCs) are emitted, giving rice its distinctive scent. Aldehydes, ketones, alcohols, esters, and other substances are among these VOCs. The type of rice, the way it is cooked, and the place where it is farmed can all affect the distinctive scent of the grain. For instance, basmati rice has a nutty and fragrant scent, whilst jasmine rice is renowned for its sweet and flowery perfume. The scent of rice affects its perceived quality and nutritional worth in addition to its sensory appeal. According to studies, individuals frequently believe that rice with a strong scent is of a higher calibre and has more nutrients. Overall, many people all around the world appreciate and like rice's scent, which is a significant component of its total sensory experience appeal. Rice is priced based on its scent in both domestic and foreign markets. More than 500 volatile chemical molecules, for example, impact the scent of rice, among other things[13]. The most crucial for rice flavouring among them has been discovered as 2 acetyl-1-pyrroline (2AP)[14, 15]. 2AP is an aromatic chemical that some researchers have also called 2-AcPy or 2-ACP. 2-AP is known by its IUPAC designation, 5-acetyl-3, 4-dihydro-2H-pyrrole. The content of 2AP in rice may be separated out and quantified using a variety of techniques, but the most practical method is gas chromatography combined with mass spectrometry, which is also the most expensive and time-consuming. Although both short and long grains include GC-MS 2AP and other aromatic chemicals. As a result, there are several ways to assess rice's scent. Generally speaking, scent may be measured by either boiling the kernel or vegetative component and then sniffing the cooked kernel for aroma[10, 16] or by testing the flavour by biting between the teeth[15][8][17]. The rice breeders employ a quicker method that Sood and Siddiq devised for assessing rice smell[18]. This states that any segment of the rice plant, except roots, can be treated with 0.1 M potassium hydroxide for 10–12 minutes before a panel of sensory specialists, trained in rice fragrance assessment, evaluates the emitted scent. Because each person's sensitivity to rice smell differs, a panel of specialists is required. This is currently accepted as a fundamental method for sensory evaluation of scent on a global scale. Different factors, such as variations in rice varieties and environmental factors, might affect 2AP content [19].

Comment [JY9]: This statement is incomplete.

2. GENETIC BASIS OF AROMATIC RICE

The scent of rice is thought to be controlled by a single recessive gene at first, but further research by other scientists showed that this attribute may also be controlled by a dominant gene or additional genes[20][21][22]. On chromosomes 4, 8, and 12, several rice aroma QTLs have been found, whereas in Pusa 1121, at least three QTLs have been found on chromosomes 3, 4, and 8 [23]. It has been shown that the fragrance characteristic is correlated with three quantitative trait loci (QTLs). The most important QTL for fragrance is Qar08.1, which is situated on chromosome 8 [24]. This characteristic has been linked to another important QTL on chromosome 3 called qar03-1 [25]. Additionally, several alleles of the badh1 gene, which is a part of the QTL qar04.1 on chromosome 4, have been linked to fragrance in rice cultivars [8, 10]. It is interesting to note that non-aromatic genotypes of rice have homozygous forms of the main gene responsible for scent. According to research, the main factor influencing rice's scent is a non-functional variation of the badh2 gene, which produces the betaine aldehyde dehydrogenase enzyme. Certain rice genotypes lack smell due to the existence of a functioning allele of the badh2 gene. There are currently 19 alleles connected to the badh2 gene that have been identified as influencing rice cultivar aroma. The badh2-E7 variation of these alleles was discovered to be the one that was present in the

highest number in the Ugandan cultivars of aromatic rice.[26]. However, compared to the badh2-E7 gene, the badh2-p allele was shown to be more common in fragrant rice types in India. Both indica and japonica forms of cannabis included this gene, badh2-p. 8 bp deletion in exon 7 has been described as a functional signature of fragrant rice and is present in aromatic rice cultivars worldwide. [27]. Exon 13 in certain aromatic lines, however, exhibited a 3 bp gain rather than an 8 bp loss. Most of the described variants for the badh2 gene contain the deletions and additions[28]. According to Bindusree et al., Seeragasamba, a short-grain aromatic rice variety similar to indica, has fragrance thanks to an 8 bp insertion in the badh2 gene's promoter region. In addition, the 8 bp deletion "GATTAGGC" is followed by two significant mutation events (A to T and T to A) and is linked to scent in rice. Twenty of the 23 aromatic accessions in the research by Akwero et al. in Uganda [28] carried the badh2-E7 allele with an 8 bp deletion and three SNPs [29].

Comment [JY10]: Maintain consistency in the in-text citations

Comment [JY11]: Maintain consistency in the citations

3. MOLECULAR MARKERS FOR RICE AROMA GENES

Numerous markers, including DNA and molecular markers are used to identify QTLs in the genome. Each sort of molecular marker has advantages and disadvantages, and they are used for a variety of purposes. [30]. Using molecular markers, we may rapidly and precisely identify the required trait controlled by a QTL in a single cultivar. [23]"To investigate genetic variation within and among populations, as well as to identify individuals or species and trace evolutionary history, researchers commonly employ markers such as restriction fragment length polymorphism (RFLP), random amplified polymorphic DNA (RAPD), amplified fragment length polymorphism (AFLP), inter simple sequence repeats (ISSR), simple sequence repeats (SSR), and single nucleotide polymorphisms (SNPs). However, the selection of the appropriate marker for a given study depends on the research question and the organism's characteristics, as each marker has its own advantages and disadvantages.[31].

4. STANDARD OF BASMATI RICE

The most crucial aspect for consumers is the flavor and scent of the rice. The demand for high-quality rice is rising in tandem with population growth [32]. A sizable portion of India's economy continues to be devoted to basmati rice. After China, India ranks as the second-largest producer of basmati rice. Basmati rice was also produced in Pakistan[33].

5. GRAIN QUALITY IMPROVEMENT

5.1 Estimation of protein content

With the use of a hand dehusker, seeds from 118 genotypes were dried to produce brown rice, which was then crushed to a powder for the measurement of protein content. This procedure ~~is was being~~ carried out using the micro Kedah approach. ~~Afterward, afterwhich~~ the crude protein content was determined. From the chosen genotypes, 10 are high in protein, and 11 are not basmati[32]. The results are shown ~~below in Fig 1~~ after a frequency distribution table is produced and the protein content is checked.'

Comment [JY12]: Check spelling

Comment [JY13]: Check this statement again




Fig. 1. Screening of genotypes based on protein content

5.2 Amount of protein.

Between basmati, non-basmati, and wild rice species, there were significant variances in the protein content. Comparing wild accessions to basmati and non-basmati, they have higher protein content[34].

6. GRAIN CHARACTERISTICS

7.1 Milling quality

Brown rice recovery, milled rice recovery, and head rice recovery are the variables used to evaluate milling quality and milling of rice efficiency. Whole grain was defined as kernels with a length more than or equal to three-quarters[35]. The primary determinant of rice market value and one of the most crucial milled rice specifications is head rice recovery[36].

Comment [JY14]: Describe how these variables are used evaluating the milling quality.

Comment [JY15]: Three quarters of what?

Comment [JY16]: Say something about this and relate it to milling quality.

7.2 Qualitative Appearance

Appearance also plays an important role in affecting its rice market value. The grain appearance is correlated with size shape, chalkiness, and translucency [37]. Phenotype of rice is described with the help of grain length, width, and thickness. Chalkiness also affects the market value. For instance, greater chalkiness lowers the market value[38].

7. COOKING AND EATING STANDARDS

The cooking and eating quality of rice determines a number of characteristics, including how easily it may be prepared, how firm it remains, and how sticky it stays while being consumed. The ease with which certain physicochemical characteristics, such as apparent amylose content (AAC), gel consistency, gelatinization temperature (GT), and pasting viscosity, may be assessed is directly proportional to the degree to which they correlate with the cooking and eating quality of rice [39]. The starch that constitutes up to 90% of milled rice has qualities that are connected to all of these variables. The two types of

molecules that make up starch are branching amylopectin and linear and helical amylose. I2-KI solution is used in a streamlined process to quantify amylose content.[40]Rice's amylose content, as determined by the I2-KI solution technique, is more frequently referred to as the apparent amylose content (AAC). This is due to the fact that long chains of amylopectin may also bind to I2, which is why AAC is the more accurate term. Waxy milled rice (1-2%), very low milled rice (5-12%), low milled rice (12-20%), moderate milled rice (20-25%), and high milled rice (>25%) are the five various categories that milled rice may be placed into based on its AAC value. [41].

Comment [JY17]: This selected portion should be relocated to the Milling quality above (section 7.1)

8. NUTRITIONAL QUALITY

The nutritional value of most significant staple foods s nutritional value is intimately connected to people's health. Protein is the second-most prevalent component of milled rice after carbohydrates. Additionally, lysine influences the nutritional value of food[42]. In an area with low development where a lack of micronutrients is evident R, the nutritional value of rice's nutritional value may be improved by breeding in additional micronutrients. In comparison to milled rice, brown rice offers higher minerals, vitamins, dietary fiber, and phenolic[43].

9. GENETICS AND GENOMICS OF RICE GRAIN QUALITY

The most common quality traits used to evaluate rice grain quality are milling quality, attractiveness, nutritional content, and cooking quality. Numerous genes governing quality attributes have been cloned in rice utilizing cutting-edge technologies and genomics. Numerous significant quantitative trait loci (QTLs) including qBRR-10, which improves brown rice recovery, are correlated with rice milling quality [44]. Rice grain size and shape are influenced by the genes GS3, GW5, GLW7, GW8, GS2, and GS9[45]. Starch is crucial for maintaining the integrity of rice grains, especially those used for cooking and eating[45][46].

CONCLUSION AND FUTURE PERSPECTIVE.

After all review, it's concluded that aroma and grain quality alternate by the environmental effect as well as genetic effect. Aroma is a significant trait that controls the prices of variety. The gene badh2 controls the proportion of aroma. Moreover, we which can be improved aromawithusing this gene. To further understand the variations in rice aroma's kind, strength, and stability as well as the minor genes driving it, sophisticated genomic and metabolomic methods must be used. Also, there is a need to enhance the marketing and manufacturing of short-grain aromatic rice types.

Comment [JY18]: Explain this method

REFERENCES

1. Mathure, S.V., et al., *Comparative quantitative analysis of headspace volatiles and their association with BADH2 marker in non-basmati scented, basmati and non-scented rice (Oryza sativa L.) cultivars of India*. Food Chemistry, 2014. **142**: p. 383-391.
2. Ishfaq, M., et al., *Growth, yield and water productivity of dry direct seeded rice and transplanted aromatic rice under different irrigation management regimes*. Journal of Integrative Agriculture, 2020. **19**(11): p. 2656-2673.
3. Mottaleb, K.A., et al., *Benefits of the development and dissemination of climate-smart rice: ex-ante impact assessment of drought-tolerant rice in South Asia*. Mitigation and Adaptation Strategies for Global Change, 2017. **22**: p. 879-901.
4. Wakte, K., et al., *Thirty-three years of 2-acetyl-1-pyrroline, a principal basmati aroma compound in scented rice (Oryza sativa L.): a status review*. Journal of the Science of Food and Agriculture, 2017. **97**(2): p. 384-395.
5. Singh, R., et al., *Genetics and biotechnology of quality traits in aromatic rice*. Aromatic rices, 2000: p. 47-70.
6. Lina, G. and Z. Min, *Formation and release of cooked rice aroma*. Journal of Cereal Science, 2022: p. 103523.
7. Adak, T., et al., *Development and validation of HS-SPME-GCMS/MS method for quantification of 2-acetyl-1-pyrroline in rice cultivars*. Journal of Food Science and Technology, 2023: p. 1-10.
8. Ndikuryayo, C., et al., *Breeding for rice aroma and drought tolerance: a review*. Agronomy, 2022. **12**(7): p. 1726.
9. Zhang, B., et al., *Introgression lines: valuable resources for functional genomics research and breeding in rice (Oryza sativa L.)*. Frontiers in Plant Science, 2022. **13**: p. 863789.
10. Ndikuryayo, C., et al., *Breeding for Rice Aroma and Drought Tolerance: A Review*. Agronomy 2022, 12, 1726. 2022, s Note: MDPI stays neutral with regard to jurisdictional claims in published
11. Lu, S., et al., *Quantifying supply chain food loss in China with primary data: A large-scale, field-survey based analysis for staple food, vegetables, and fruits*. Resources, Conservation and Recycling, 2022. **177**: p. 106006.
12. Pandey, V., M. Pant, and V. Snasel, *Blockchain technology in food supply chains: Review and bibliometric analysis*. Technology in Society, 2022: p. 101954.
13. Verma, D.K. and P.P. Srivastav, *Extraction, identification and quantification methods of rice aroma compounds with emphasis on 2-acetyl-1-pyrroline (2-AP) and its relationship with rice quality: A comprehensive review*. Food Reviews International, 2022. **38**(2): p. 111-162.
14. Fan, Z., et al., *A multi-omics framework reveals strawberry flavor genes and their regulatory elements*. New Phytologist, 2022. **236**(3): p. 1089-1107.
15. Renuka, N., et al., *Co-functioning of 2AP precursor amino acids enhances 2-acetyl-1-pyrroline under salt stress in aromatic rice (Oryza sativa L.) cultivars*. Scientific Reports, 2022. **12**(1): p. 3911.
16. Beatriz Vieira, M., et al., *DNA-Based Tools to Certify Authenticity of Rice Varieties—An Overview*. 2022.
17. Zaghum, M.J., K. Ali, and S. Teng, *Integrated genetic and omics approaches for the regulation of nutritional activities in rice (Oryza sativa L.)*. Agriculture, 2022. **12**(11): p. 1757.
18. Khasa, R., S. Kumar, and R.K. Jain, *Marker assisted selection of Aerobic \times Basmati segregating lines for physio-morphological and aroma trait*. 2022.

19. Dwiningsih, Y. and J. Alkahtani, *Potential of Pigmented Rice Variety Cempo Ireng in Rice Breeding Program for Improving Food Sustainability*. 2023.
20. Sakran, R.M., et al., *Molecular genetic diversity and combining ability for some physiological and agronomic traits in rice under well-watered and water-deficit conditions*. *Plants*, 2022. **11**(5): p. 702.
21. Tian, Y., et al., *Creation of Two-Line Fragrant Glutinous Hybrid Rice by Editing the Wx and OsBADH2 Genes via the CRISPR/Cas9 System*. *International Journal of Molecular Sciences*, 2023. **24**(1): p. 849.
22. Negoro, H. and H. Ishida, *Development of sake yeast breeding and analysis of genes related to its various phenotypes*. *FEMS Yeast Research*, 2022. **22**(1).
23. Ullah, I., et al., *Pyramiding of Four Broad Spectrum Bacterial Blight Resistance Genes in Cross Breeds of Basmati Rice*. *Plants*, 2023. **12**(1): p. 46.
24. Yuan, H., et al., *Fine mapping and candidate gene analysis of qGSN5, a novel quantitative trait locus coordinating grain size and grain number in rice*. *Theoretical and Applied Genetics*, 2022: p. 1-14.
25. Li, T., et al., *Genetic dissection of quantitative trait loci for grain size and weight by high-resolution genetic mapping in bread wheat (*Triticum aestivum* L.)*. *Theoretical and Applied Genetics*, 2022: p. 1-15.
26. Imran, M., S. Shafiq, and X. Tang, *CRISPR-Cas9-mediated editing of BADH2 gene triggered fragrance revolution in rice*. *Physiol Plant*, 2023. **175**(1): p. e13871.
27. Cheng, J., et al., *Rice grain quality: Where we are and where to go?* *Advances in Agronomy*, 2022. **172**: p. 211-252.
28. Sreenivasulu, N., et al., *Post-genomics revolution in designing premium quality rice in the high yielding background to address the demands of consumers from the 21st Century*. *Plant Communications*, 2022.
29. Singh, G., et al., *Molecular profiling of BADH2 locus reveals distinct functional allelic polymorphism associated with fragrance variation in Indian aromatic rice germplasm*. *Physiology and Molecular Biology of Plants*, 2022. **28**(5): p. 1013-1027.
30. Tao, C., et al., *Development of New Low Glutelin Content japonica Rice Lines with Good Eating Quality and Fragrance by Molecular Marker-Assisted Selection*. *Chinese Journal OF Rice Science*, 2023. **37**(1): p. 55.
31. Ab Razak, S., et al., *Identification of aromatic rice from genetic landrace resource using molecular marker integrated with chemical assessment*. *Journal of Biotech Research*, 2022. **13**: p. 199-206.
32. Khurana, S. and S. Kumar, *Milling, Physicochemical, Cooking and Textural Properties of Some Basmati and Non-Basmati Rice Cultivars*.
33. Pandey, S., et al., *Curcumin loaded core-shell biopolymers colloid and its incorporation in Indian Basmati rice: An enhanced stability, anti-oxidant activity and sensory attributes of fortified rice*. *Food Chemistry*, 2022. **387**: p. 132860.
34. Kaur, R., et al., *Protein profiling in a set of wild rice species and rice cultivars: a stepping stone to protein quality improvement*. *Cereal Research Communications*, 2022: p. 1-15.
35. Deng, Z., et al., *Waxy is an important factor for grain fissure resistance and head rice yield as revealed by a genome-wide association study*. *Journal of Experimental Botany*, 2022. **73**(19): p. 6942-6954.
36. Aboye, A.D. and A.M. Teto, *Pilot Review Study on Current Research Status and Economic Value Limitations of Bread Wheat (*Triticum aestivum* L.) Production in Ethiopia*. *Economics*, 2022. **11**(4): p. 137-166.

37. Fan, P., et al., *Recent research advances in the development of chalkiness and transparency in rice*. Agriculture, 2022. **12**(8): p. 1123.
38. Lu, Y., et al., *Combined effects of SSII-2RNAi and different Wx alleles on rice grain transparency and physicochemical properties*. Carbohydrate Polymers, 2023. **308**: p. 120651.
39. Kheto, A., et al., *Advances in isolation, characterization, modification, and application of Chenopodium starch: A comprehensive review*. International Journal of Biological Macromolecules, 2022.
40. Raj, S.R.G. and K. Nadarajah, *QTL and Candidate Genes: Techniques and Advancement in Abiotic Stress Resistance Breeding of Major Cereals*. International Journal of Molecular Sciences, 2022. **24**(1): p. 6.
41. Shivakoti, R., *Validation of NINA-DISH food frequency questionnaire with multiple 24-hour dietary recalls among pregnant women in Pune, India*.
42. Eldridge, A.L. and E.A. Offord, *Global landscape of nutrient inadequacies in toddlers and young children*. Building Future Health and Well-Being of Thriving Toddlers and Young Children, 2020. **95**: p. 12-22.
43. Kamaluddin, P.S., et al., *Marker-Assisted Selection for Value Addition in Crop Plants*. Technologies in Plant Biotechnology and Breeding of Field Crops, 2022: p. 23.
44. Zhou, H., D. Xia, and Y. He, *Rice grain quality—traditional traits for high quality rice and health-plus substances*. Molecular Breeding, 2020. **40**: p. 1-17.
45. Mbanjo, E.G.N., et al., *The genetic basis and nutritional benefits of pigmented rice grain*. Frontiers in genetics, 2020. **11**: p. 229.
46. Wei, X., et al., *A quantitative genomics map of rice provides genetic insights and guides breeding*. Nature Genetics, 2021. **53**(2): p. 243-253.