

RELATIONSHIP BETWEEN AGE AND LUMBOSACRAL ANGLE (LSA) OF ADULT NIGERIANS: A CROSS-SECTIONAL STUDY.

ABSTRACT

Background: The study evaluated the relationship between age and lumbosacral angle (LSA) of adult Nigerians. **Methods:** Two hundred and twenty-six (226) radiographs (113 males to 113 females) were obtained from Radiology departments of Federal Medical Centre (FMC), Owerri. Inclusion criteria were, lateral radiographs of lumbosacral region marked above 18 years of ages and only clear X-ray films were used. Using the Ferguson's approach, the LSA angle was measured. All data obtained were expressed both in descriptive and inferential statistics. **Results:** Both male and female categories showed a progressive increase in LSA across age groups, but later decreased in the age group of 60 years and above. The mean LSA of male subjects was 35.74° while that of females was 41.46° . There was a significant difference ($t = 3.16$, $p = 0.002$) between the LSA of male and female categories. However, there was a weak, positive correlation between age and LSA of males ($r = 0.187$) and females ($r = 0.392$). **Conclusion:** The males had a lower mean LSA compared to the females. Age was significantly correlated with LSA in both sexes.

Keywords: Lumbosacral angle, Age, Nigerians.

INTRODUCTION

The vertebral column, also called the backbone, is a series of approximately 33 bones called vertebrae, which are separated by intervertebral discs that perform protective, supportive, and axial and movement functions. There are about 23 intervertebral discs in the vertebral column which account for one-quarter of the total length of the backbone and vary from one individual to another. The thoracic and pelvic curvatures are called primary curvatures whereas cervical and lumbar curves are termed secondary curvatures [1].

The most important region in the vertebral column in terms of mobility and weight bearing is the lumbosacral region. It is the key weight-bearing region of the vertebral column [2,3]. Abnormality and pathologies of the lumbar vertebral region contributes to the development of a collection of pathological symptoms such as low back pain [4,5].

The lumbar lordotic curve is the anterior curvature of the lumbar spine in the mid-sagittal plane [6]. One of the major postural analyses of an individual is lumbar lordosis. Lumbar lordosis is defined to be the inwardly concave curvature which extends from the first lumbar vertebrae to the fifth lumbar vertebrae, formed majorly by the wedging of the intervertebral discs of each of these vertebrae and counters the forward inclination of the sacrum and thus restores the upward inclination of the spinal column [7].

Of all these methods, radiography remains the gold standard and lordotic measurement can accurately be measured in a supine lateral lumbosacral spine radiograph [8]. The lumbosacral angle (LSA), one of the clinically important radiographic angles related to this curve, and frequently measured by the Ferguson's technique is important in the management of patients with low back disorders because it may be affected by disorders such as inflammation, and degeneration [9,10].

Various study data on lumbosacral angle (LSA) available suggest differences with respect to sexual dimorphism, racial or ethnic populations [11,12,13,14]. Few studies have examined the effects of body type and size on LSA [15,16]. However, there is no consensus of opinion on the significance of either a decreased or an increased LSA with age. Madufuro *et al* [17] concluded that lumbosacral angle does not increase significantly after the age 36–40 years. There is need to update current knowledge on the relationship between age and LSA of adult Nigerians, hence this study.

MATERIALS AND METHODS

This study was retrospective and non-interventional. The information was gathered from lumbosacral radiographs as well as a sample of anthropometric and demographic information from x-ray films in the Radiology department of Federal Medical Centre (FMC), Owerri. The clinical research and ethics committees of this hospital granted approval for the study's execution. As part of the study inclusion criteria, only lateral radiographs of lumbosacral region marked above 18 years of ages and clear x-ray films were used. Exclusion criteria included x-ray films with any macro trauma to the low back region, x-ray films with a clinically or radiographically detectable scoliosis and/or kyphosis of the lumbar spine and x-rays films with any spinal surgery or instrumentation.

In order to measure the Lumbosacral Angle/Ferguson's angle, the Ferguson's approach was employed in this study since it is the most often used technique for calculating this angle. It is calculated from the lumbosacral spine's lateral projection using the following procedure;

- Draw a line parallel to the sacral base, plane of the lumbosacral joint (AB)
- Draw another plane horizontal to the sacral base (AC) as part of Ferguson's approach for calculating the lumbosacral angle
- Measure the angle created between the horizontal plane and the plane parallel to the sacral base.




Figure 1: Ferguson's Method of measuring Lumbosacral Angle [18].

Data were analyzed using Statistical Package for social Sciences (SPSS) version 23.0. Both descriptive and inferential statistics were applied. The means of LSA for both male and female categories in line with their respective age groups were expressed in form of tables. Significant differences between male and female LSA were done using Independent Samples t-test, and correlation between age and LSA for both sexes were performed using Pearson's correlation. Level of significance was set at 0.05.

RESULTS

The data of two hundred and twenty-six (226) radiographs comprising of one hundred and thirteen (113) radiographs each for males and females, which met the inclusion criteria were

analysed. For the male category as shown in table 1, the mean LSA for age group (<31 years) was 23.09±11.21°, mean LSA for age group (31 – 40 years) was 34.29±9.02°, mean LSA for age group (41 – 50 years) was 36.88±11.70°, mean LSA for age group (51 – 60 years) was 41.61±9.37°, and mean LSA for age group (> 60years) was 36.80±12.34°. For the female category, the mean LSA for age group (<31 years) was 25.42±7.26°, mean LSA for age group (31 – 40 years) was 34.21±12.81°, mean LSA for age group (41 – 50 years) was 39.50±12.28°, mean LSA for age group (51 – 60 years) was 47.61±11.71°, and mean LSA for age group (> 60years) was 46.09±11.67°.

There was a significant difference ($t = 3.16$, $p = 0.002$) between the LSA of male and female categories. However, there was a weak, positive correlation between age and LSA of males ($r = 0.187$) and females ($r = 0.392$).

Table 1: Age-based descriptive statistics for LSA by sex

Age group (years)	Male LSA (°)			Female LSA (°)		
	Mean±SD	SEM	CL (95.0%)	Mean±SD	SEM	CL (95.0%)
< 31	23.09±11.21	2.89	6.21	25.42±7.26	15.00	9.01
31 - 40	34.29±9.02	2.07	4.35	34.21±12.81	2.56	5.29
41 -50	36.88±11.70	2.62	5.48	39.50±12.28	2.51	5.19
51 -60	41.61±9.37	2.42	5.19	47.61±11.71	2.44	5.06
> 60	36.80±12.34	1.86	3.75	46.09±11.67	1.94	3.95
All ages	35.74±12.67	1.19	2.36	41.46±13.27	1.25	2.47

LSA = Lumbosacral Angle, SEM = Standard Error of Mean, SD = Standard Deviation, CL = Confidence Level




Figure 2: Mean values of Lumbosacral angles (LSA) for males and females by age

Table 2: Sexual differences between age and LSA of Nigerians

Parameters	Mean±SD	N	SEM	df	t	p < 0.05	Inference
Age	Female	53.43±15.30	113	1.44	112	0.82	0.412
	Male	51.73±16.42	113	1.55			
LSA	Female	41.46±13.27	113	1.25	112	3.16	0.002
	Male	35.74±12.67	113	1.19			

LSA = Lumbosacral Angle, SEM = Standard Error of Mean, SD = Standard Deviation, df = degree of freedom

Table 3: Pearson's correlation between Age and LSA of Nigerians

Parameters	Male		Female		
	Age	LSA	Age	LSA	
Age	Pearson Correlation				
	1	.187*	1	.392**	
LSA	Pearson Correlation				
	.187*	1	.392**	1	
		Sig. (2-tailed)			
		0.047		<.001	

LSA = Lumbosacral Angle

DISCUSSIONS

The present study was done to evaluate the relationship between age and lumbosacral angle (LSA) of adult Nigerians. The mean LSA for age group (<31 years) was 23.09±11.21°, mean

LSA for age group (31 – 40 years) was $34.29 \pm 9.02^\circ$, mean LSA for age group (41 – 50 years) was $36.88 \pm 11.70^\circ$, mean LSA for age group (51 – 60 years) was $41.61 \pm 9.37^\circ$, and mean LSA for age group (> 60years) was $36.80 \pm 12.34^\circ$. For the female category, the mean LSA for age group (<31 years) was $25.42 \pm 35.50^\circ$, mean LSA for age group (31 – 40 years) was $34.21 \pm 12.81^\circ$, mean LSA for age group (41 – 50 years) was $39.50 \pm 12.28^\circ$, mean LSA for age group (51 – 60 years) was $47.61 \pm 11.71^\circ$, and mean LSA for age group (> 60years) was $46.09 \pm 11.67^\circ$.

In line with a similar study by Maduforo *et al.* [17], there was a relative increase in LSA with age up to a maximum age group 36-40years. Okpala [12] also reported from his study that as the age group increased in study population, there was an increase in the LSA until it began to decrease when the age group was above 65 years of age. Also, an earlier research done using a Caucasian population showed that their LSA values varied between 15° and 25° in an erect position and that females have higher values than males [19]. In concordance with this current study, Kalichman *et al.* [20] concluded there was no difference in the lumbosacral angle between males and females, however, the study found that there was a significant variation between them – which contradicts this current study.

In this study, the mean LSA of male subjects was 35.74° while that of females was 41.46° . This differences were significant and they were in concordance with a study done by Mukherjee *et al* [9] which reported that the mean LSA in females were larger compared to that of the study males. In addendum, Agichani *et al.* [21] found in their study that the mean LSA was $38.27^\circ (\pm 8.66)$ in males and $40.57^\circ (\pm 7.18)$ in females. As shown in table 4, similar studies showed varying LSA values suggesting ethnic and racial differences among the study populations.

Table 4: Comparison of mean LSA of the present study with related literature

Author(s)	Year	Gender	Mean LSA
Okpala [12]	2014	Both	43.38° (males), 45.54° (females)
Agichani <i>et al.</i> [21]	2017	Both	38.27° (males), 40.57° (females)
Naqvi <i>et al.</i> [14]	2020	Both	36° (males), 33° (females)
Mukherjee <i>et al.</i> [9]	2021	Both	37° (males), 45° (females)

Present study	2023	Both	35.74° (males)
			41.46° (females)

LSA = Lumbosacral angle

In this present study, there were significant weak, positive correlations between age and LSA ($p < 0.05$) in both male and female subjects. This agrees with a related study done by Wagh *et al.* [22] which reported significant, weak correlations between age and LSA in both males ($p = 0.0177$) and females ($p = 0.0151$). In another study, a significant relationship was also noticed between age and LSA [23].

CONCLUSION

The study concluded that the LSA progressively increased with age across age groups and later decreased at the age group of 60 years and above. The males had a lower mean LSA compared to that of the females. Age was significantly correlated with LSA in both sexes.

REFERENCES

1. Bazira, P. J. (2021). Clinically applied anatomy of the vertebral column. *Surgery (Oxford)*, 39(6), 315-323.
2. Loughenbury, P. R., & Tsirikos, A. I. (2020). Indications and principles of conservative treatment in injuries affecting the vertebral column. *Orthopaedics and Trauma*, 34(5), 291-297.
3. Sluis, I. F., Bartholdy, B. P., Hoogland, M. L., & Schrader, S. A. (2022). Age estimation using vertebral bone spurs; Testing the efficacy of three methods on a European population. *Forensic Science International: Reports*, 6, 100301.
4. Li, W., Gong, Y., Liu, J., Guo, Y., Tang, H., Qin, S., & Chen, B. (2021). Peripheral and central pathological mechanisms of chronic low back pain: a narrative review. *Journal of Pain Research*, 1483-1494.
5. Noonan, A. M., & Brown, S. H. (2021). Paraspinal muscle pathophysiology associated with low back pain and spine degenerative disorders. *JOR spine*, 4(3), e1171.
6. Bogduk, N. Clinical anatomy of the lumbar spine and sacrum. 4th ed., Churchill Livingstone, pp.51-54, 2005.
7. Been, E., Barash, A., Pessah, H. & Peleg, S. (2010), A new look at the geometry of the lumbar spine, *Spine*, 35:1014–1017.

8. Vrtovec T, Pernus F, Likar B. A review of methods for quantitative evaluation of spinal curvature. *Eur Spine J* 2009; 18: 593-607.
9. Mukherjee, U., Kundu, S., Gulati, R., & Mandal, P. (2021). Lumbosacral angle variations in middle aged patients with chronic low back pain-A retrospective study. *Pain*, 1(2), 3.
10. Napitupulu, P., Nareswari, G., Rahmansyah, M., & Tanaji, T. C. (2022). Radiographic lumbosacral angle has a low yield for diagnosing low back pain in workers 20-70 years of age. *Universa Medicina*, 41(3), 263-270.
11. Oyakhire, M.O., Didia, B.C. & Bob-Evans, Y.E.(2013). Radiographic evaluation of the Lumbar Lordotic angle of the Spine in a population of Nigerians. *Asian Journal of Medical Science*, 4(3): 69-75.
12. Okpala FO. Measurement of lumbosacral angle in normal radiographs: A retrospective study in Southeast Nigeria. *Ann Med Health Sci Res* 2014;4:757-62.
13. Khanal UP. Measurement of Lumbosacral Angle in Normal Nepalese Population by Computed Tomography. *J Clin Res Radiol* 2018;1(2):1-7.
14. Naqvi SZG, Ali A, Siddiqui A, Ali SD, Qureshi M, Aliuddin AM. Measurement of Lumbosacral Angle in Normal Radiographs: A Cross-Sectional Study. *J Liaq Uni Med Health Sci [Internet]*. 2020 Dec. 30 [cited 2023 Apr. 12];19(04):238-41.
15. Araújo, F. A., Simões, D., Silva, P., Alegrete, N., & Lucas, R. (2019). Sagittal standing posture and relationships with anthropometrics and body composition during childhood. *Gait & posture*, 73, 45-51.
16. Tang, L., Hu, Z., Lin, Y. S., & Hu, J. (2022). A statistical lumbar spine geometry model accounting for variations by Age, Sex, Stature, and body mass index. *Journal of biomechanics*, 130, 110821.
17. Maduforo C, West O, Nwankwo N, Onwuchekwa R, Etawo US, Ogbulu D. (2021). Study of the Lumbosacral Angles of Males in Port Harcourt, South- South, Nigeria. *The Nigerian Health Journal*, Vol. 12(1): 22-24\
18. Hellems H, Keats T. (1971). Measurement of the normal lumbosacral angle. *Am J Roentgenol*. 113:642-5.
19. Bryan GJ. Lumbar vertebrae. In: *Diagnostic Radiography, A Concise Practical Manual*. 4th ed. London: Churchill Livingstone; 1987. p. 152.

20. Kalichman L, Li L, Hunter DJ, Been E. Association between computed tomography-evaluated lumbar lordosis and features of spinal degeneration, evaluated in supine position. *Spine J* 2011;11:308-15.
21. Agichani S, Joshi SD, Joshi SS. (2017). Evaluation of lumbosacral angle amongst central Indians. *Journal of Evolution of Medical and Dental Sciences*. 6(83): 5797-5801.
22. Wagh D, Nikam V, Takale S. (2021). Geometric Study of Lumbosacral Spine Radiographic Parameters in Western Maharashtra Population. *International Journal of All Research Education and Scientific Methods*. 9(8): 1291-1298
23. Tao Y, Galbusera F, Niemeyer F, Samartzis D, Vogele D, Wilke HJ. (2021). Radiographic cervical spine degenerative findings: a study on a large population from age 18 to 97 years. *Eur Spine J*. 30(2):431-443.