

UNDIFFERENTIATED CONNECTIVE TISSUE DISORDER PRESENTING AS TUBERCULAR MONOARTHRITIS AND MEMBRANOUS NEPHROPATHY: A CASE REPORT

ABSTRACT

Background:

The Term undifferentiated connective tissue disease is used to define conditions characterised by the presence of signs and symptoms suggestive of a systemic autoimmune disease that does not satisfy the classification criteria for defined connective tissue diseases (CTD) such as systemic lupus erythematosus (SLE), Sjögren's syndrome (SS), rheumatoid arthritis (RA) and others[1].

The most characteristic symptoms of UCTD are represented by arthritis and arthralgias, Raynaud's phenomenon and leukopenia, while neurological and renal involvements are rare. Only one-third of the UCTDs evolve into full-blown CTD within the initial years (3-5). Here we are presenting a case of a 36-year female presenting with mono arthritis, proteinuria and severe anaemia that was diagnosed to have undifferentiated connective tissue disorder with tubercular mono arthritis of knee joint and membranous nephropathy. The patient was given anti-tubercular drugs , corticosteroids, hydroxychloroquine and PRBC transfusions. The patient was discharged and showed significant improvement on subsequent follow-ups. This case report provides insight into the various presentations of connective tissue disorder and also its multi-system involvement. Also sometimes these disorders cannot be classified by a criterion clinically or based on laboratory investigations into one single connective tissue disease.

1. AIMS

This case report provides insight into the various presentations of connective tissue disorder and also its multi-system involvement.

2. INTRODUCTOION

Undifferentiated connective tissue disease (UCTD) is a systemic heterogeneous group of autoimmune diseases which do not fulfill the criteria of connective tissue diseases.^[2,3]In 1980, LeRoy et al proposed the concept of undifferentiated connective-tissue syndromes (UCTS) to characterize mixed or overlapping syndromes. ^[4] Since that time, innumerable case reports, case series, and clinical studies have used variable criteria to define UCTD. Such variation in the definition of UCTD results in ambiguity of study results and difficulty with interpreting the findings and conclusions. In 1999, Mosca et al proposed preliminary classification criteria for UCTD that have become increasingly accepted and used.^[5] The authors suggested defining cases of UCTD as those in which signs and symptoms are consistent with a CTD but that do not fulfill the classification or diagnostic criteria for any one of the defined CTDs (ie, RA, SLE, SSc, PM/DM, MCTD, SS). In order to fulfill the criteria for UCTD, antinuclear antibodies must be present, along with a disease duration of at least 3 years. The kidney is frequently involved in the clinical course of connective tissue diseases in which a wide variety of glomerular, tubulointerstitial and vascular lesions are observed. In undifferentiated connective tissue disease (UCTD) there is evidence that renal involvement is mild and there are limited detailed histopathological studies.

3. CASE PRESENTATION

A 36-year female patient not a known case of any co morbidities, a housewife presented with complaints of weight loss for 2 months and right-sided knee joint swelling for one month. The patient was relatively asymptomatic before 2 months then the patient developed a complaint of weight loss. It was gradual in onset and progression. It was associated with easy fatigability and generalized weakness. The patient was unable to carry out daily activities. The patient developed a complaint of swelling over the right knee joint. It was gradual in onset and progressive. The patient was unable to bear weight on that limb. It was associated with swelling and tenderness and reduced range of motion of that joint. There was no history of fever, chest pain, breathlessness, multiple joint pains, joint stiffness or swelling of any other joints. The patient had no significant past history. The patient was 2 Para with Both full-term normal vaginal deliveries. The patient had a complaint of oligomenorrhea. LMP was 24/5/2021. No significant personal history. On admission, the patient was conscious, cooperative and well oriented to time place and person. Her temperature was afebrile. Pulse was 110 beats per minute , blood pressure was 100/60 mm of hg. On general examination pallor was present and mild grade I pedal oedema was present . Following investigations were done

Hemoglobin	5.2 mg/dl
Total counts	10900 cells per cu mm
Platelets	2.71 lac per cu mm
ESR	21 mm/hour
S. Creatinine	1.5 mg/dl

total bilirubin	0.3
Serum sodium	139 mmol/L
Serum potassium	3.9 mmol/L
Serum Total protein	7 g/dl
Albumin	2

Based on the findings in above investigations, patient was further investigated for anaemia and renal impairment. The patient was found to have microcytic hypochromic anaemia but iron profile was within normal limits. Further USG abdomen was done wherein the kidney size was normal with preserved cortico- medullary differentiation. Also in view of proteinuria 24 hour, urinary protein was sent. Also suspecting an autoimmune condition Antinuclear Antibody (ANA) profile was sent. The 24-hour urinary protein was 1700 mg. ANA was positive at 1:40 dilution with the nuclear pattern . Based on above findings further ANA profile and complement 3 and complement 4 were sent (C3 , C4) .C3,C4 were within normal ranges so were cholesterol and triglycerides. All of these findings indicated Nephritic syndrome. ANA profile was positive for anti- Sjögren's syndrome-related antigen A(SSA/ anti-Ro antibodies), anti- Sjögren's syndrome-related antigen B (SSB). Ultrasonography of right knee indicated an echoic collection with few internal septa and no internal vascularity noted in the suprapatellar pouch approx 7*3.4 in size extending from the knee suggestive of moderate joint effusion in the right knee. After orthopaedic consultation, joint fluid was drained and sent for examination. The fluid was turbid pale yellow in colour with ph of 7, sugar 50mg/dl, protein 1.0/dl, Adenosine deaminase (ADA) 29u/L and Lactate dehydrogenase (LDH) 569u/L . Total cell count of synovial fluid were 13000 out of which 95% were polymorphs ,5% lymphocytes . No organism was seen on gram , ZN and KOH stains nor any organism was detected in culture. Fluid was sent for Cartridge based nucleic acid amplification test (CBNAAT) under National tuberculosis elimination programme (NTEP) to rule out Koch.

During the course of treatment , seven pints of packed cell volume was transfused. After attaining Hb of 10.6, a renal biopsy was done to find out the definite pathology in kidneys and to rule out lupus nephritis . The renal biopsy was suggestive of membranous nephropathy. Meanwhile the CBNAAT for synovial fluid came positive for mycobacterium tuberculosis, sensitive to rifampicin and isoniazid . The patient was given anti-tubercular drugs according to the NTEP regime and along with that she was given corticosteroids and hydroxychloroquine . After few months of treatment the patient didn't achieve remission and hence was started on azathioprine. Patient is currently on azathioprine .

4. DISCUSSION

A considerable proportion of individuals have clinical and serological features that are suggestive of a systemic autoimmune disorder, but cannot be diagnosed as having a defined connective tissue disease. These diseases have been variably defined as incomplete systemic lupus erythematosus, latent lupus, early undifferentiated connective tissue diseases, and undifferentiated connective tissue diseases (UCTD). It is the fact that the majority of patients

(80–99%) are female, with a mean age at disease onset ranging from 32 to 44 years [7]. Literature data show clearly that 30% of patients with an undifferentiated onset will develop a defined CTD over the follow-up. Most commonly patients develop systemic lupus erythematosus (SLE), although an evolution to other CTDs such as systemic sclerosis (SSc), Sjogren's syndrome (SS), mixed connective tissue disease (MCTD), systemic vasculitis, polydermatomyositis (PM/DM), rheumatoid arthritis (RA). [9]

As patient didn't have any rash / oral ulcers / pericarditis / pleuritis / psychosis and nor anti sm and anti ds DNA were present hence SLE was less likely. Also, the patient didn't have any complaints of dry eye or dry mouth nor any positive oral or ocular signs, so the possibility of sjogrens was less likely. As the disorder didn't fit into a single criterion , it was classified as early undifferentiated connective tissue disorder.

Diagnosis of SLE is based on clinical findings and laboratory findings after all other differential diagnoses are excluded. The diagnosis can be refined with a detailed medical history, physical examination, SLE-specific laboratory tests, and regular follow-up. The American College of Rheumatology (ACR) defined in 1997 or the Systemic Lupus International Collaborating Clinics (SLICC) classification criteria described in 2012 are often used by clinicians due to the lack of specific diagnostic criteria. [5,6] As these criteria are insufficient in some cases, some other diagnostic criteria have been proposed in the literature. According to these criteria, if the patient has non-specific organ involvement for SLE (e.g., optic neuritis), and fulfils two or three criteria of ACR or SLICC (four criteria for the definite diagnosis should be positive), the patient is diagnosed with probable SLE. [7] Although this case did not fully meet the SLE criteria, she fulfilled three criteria according to SLICC: non-scarring alopecia, peripheral sensorial neuropathy, and ANA positivity. These findings were compatible with the probable SLE described in the literature. In addition, the literature data suggest that a patient having signs and symptoms suggestive of a connective tissue disease (CTD) without fulfilling the criteria of any defined CTD and having ANA positivity can be defined as having a UCTD. If the disease duration is less than three years, patients can be defined as having an early UCTD. A longer follow-up would allow the correct classification of at least some false-negative patients.

Classification criteria of the SS have been updated for years and the most recent 2016 ACR and European League Against Rheumatism (EULAR) criteria have been established. Primary SS is not associated with other diseases, while secondary SS may be associated with other rheumatic diseases. In this case, although there was no supportive finding in the salivary gland biopsy, Schirmer's test results was negative. As the patient could not fulfil the primary SS classification criteria fully, she was diagnosed with secondary SS.

5.CONCLUSION

Inflammatory rheumatic diseases can be seen together, and diseases may overlap with each other. This case highlights the importance of detailed medical history and ultrasonographic assessment in addition to laboratory testing in rheumatic diseases. Based on our experience, we suggest that methotrexate may be useful when UCTD and axSpA are seen together.

6. REFERENCES:

1. Mosca M, Tani C, Neri C, Baldini C, Bombardieri S. Undifferentiated connective tissue diseases (UCTD). *Autoimmun Rev.* 2006 Nov;6(1):1-4. doi: 10.1016/j.autrev.2006.03.004. Epub 2006 Apr 19. PMID: 17110308.
2. Cervera R, Khamashta MA, Hughes GR. 'Overlap' syndromes. *Ann Rheum Dis.* 1990 Nov;49(11):947-8. doi: 10.1136/ard.49.11.947. PMID: 2256747; PMCID: PMC1004273.
3. Kelly A, Panush RS. Diagnostic uncertainty and epistemologic humility. *Clin Rheumatol.* 2017 Jun;36(6):1211-1214. doi: 10.1007/s10067-017-3631-8. Epub 2017 Apr 22. PMID: 28432522.
4. LeRoy EC, Maricq HR, Kahaleh MB. Undifferentiated connective tissue syndromes. *Arthritis Rheum.* 1980 Mar;23(3):341-3. doi: 10.1002/art.1780230312. PMID: 7362686..
5. Mosca M, Neri R, Bombardieri S. Undifferentiated connective tissue diseases (UCTD): a review of the literature and a proposal for preliminary classification criteria. *Clin Exp Rheumatol.* 1999 Sep-Oct;17(5):615-20. PMID: 10544849..
6. Sciascia S, Roccatello D, Radin M, Parodis I, Yazdany J, Pons-Estel G, Mosca M. Differentiating between UCTD and early-stage SLE: from definitions to clinical approach. *Nat Rev Rheumatol.* 2022 Jan;18(1):9-21. doi: 10.1038/s41584-021-00710-2. Epub 2021 Nov 11. PMID: 34764455.
7. Cavazzana I, Franceschini F, Belfiore N, Quinzanini M, Caporali R, Calzavara-Pinton P, Bettoni L, Brucato A, Cattaneo R, Montecucco C. Undifferentiated connective tissue disease with antibodies to Ro/SSA: clinical features and follow-up of 148 patients. *Clin Exp Rheumatol.* 2001 Jul-Aug;19(4):403-9. PMID: 11491495.
8. Mosca M, Neri R, Bencivelli W, Tavoni A, Bombardieri S. Undifferentiated connective tissue disease: analysis of 83 patients with a minimum followup of 5 years. *J Rheumatol.* 2002 Nov;29(11):2345-9. PMID: 12415590.