

# Original Research Article

## Soil Health: Indicators of Soil Management for Vegetable Cultivation under the Clove plant

---

### ABSTRACT

Low soil health indicates poor productivity soil optimal for supporting agricultural businesses because it will reduce the production of agricultural commodities for vegetable crop farmers in Ternate City which are carried out under plantation crops. This study aims to determine the level of soil health of vegetable farms under clove plantations. Field observation methods and laboratory analysis were carried out to observe the characteristics of the land (soil and plants) which became the research variables. The drill and soil profile approach is used in observing and taking soil samples. Research data were analyzed quantitatively and qualitatively by using a scale (scoring value) and then determined the level of soil health of agricultural land in the study location. Soil variables observed were physical properties (including soil morphology), chemical properties soil, and soil biological properties. The results showed that the level of soil health on vegetable farms under clove stands in Tobololo Village was in the medium (53-58%) and high (63-67%) classes. Soil health indicators which are variables that affect the decline in the value of soil health on vegetable farms under clove stands are C-organic, base saturation, total nitrogen, and available potassium. The provision of organic matter, nitrogen, and potassium fertilizers will increase base saturation as well as soil fertility and health.

*Keywords: soil health indicators, soil characteristics, soil health, vegetable crops*

### 1. INTRODUCTION (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

The increasing population of Ternate city has an impact on the economic needs of farmers and consumers, especially on the need for food and vegetable consumption. The demand for vegetable production is increasing, on the one hand, clove and nutmeg farmers also need additional income to meet their daily needs, while harvesting clove flowers, seeds, and nutmeg mace takes 6 months to 1 year[1]. In addition to the need for food, the need for shelter has also increased resulting in land conversion, where productive agricultural land is used for settlements[2];[3]. As a result, small and narrow lands under clove and nutmeg plantations or between gardens in clove and nutmeg plantations bordering the forest are processed into fields for cultivating food crops and vegetables. The small land area causes the number of commodities cultivated by farmers to be small but many types on one business land which are planted simultaneously or mixed in the form of *multiple cropping* or planting after harvest which is known as *intercropping (relay cropping)*[4]. Types of horticultural crops and vegetables that are cultivated and have business opportunities (R/C ratio and B/C ratio > 1) to be developed by farmers on Ternate Island include tomatoes (*Solanum lycopersicum*), chillies (*Capsicum frutescens*), cucumbers (*Cucumis sativus*), long beans (*Vigna unguiculata* sp), eggplant (*Solanum melongena*), kale (*Ipomoea aquatica*), spinach (*Amaranthus* sp), vegetable terubuk/candles (*Saccharum edule*), and pumpkin[5].

The island of Ternate is physiographically an archipelago, a small island with a small lowland area, and is dominated by mountains. This resulted in agricultural business land being

generally located in hilly areas with a slope of > 8%, thus limiting farmers in planting the types of plants and cultivation techniques used. The hilly shape of the land, the small land area, and the types of plants and cultivation technology used are natural conditions and problems that exist for farmers on the slopes of Mount Gamalama, but because the land is the result of a volcanic eruption, the soil is fertile for planting various types of plants. Such land characteristics become important natural assets for the livelihoods of the people of Ternate Island[6]. Continuous land management will reduce soil health thereby affecting the productivity of vegetable crops cultivated under clove and nutmeg stands.

Soil health is an indicator of land quality and productivity[7]; [8]. The concept of soil health is the integration and optimization of soil properties that are useful for increasing land productivity, especially the main components of land, namely soil, plants, and the environment[9];[10]. Healthy soil means fertile and productive soil to be used as agricultural land for the cultivation of crops, both horticultural (vegetable) crops, food crops, and plantation crops as well as agroforestry [11]. Based on these problems, this research was conducted to know the level of soil health and land productivity as well as management measures for sustainable agriculture.

## 2. MATERIAL AND METHODS

Ternate Island as the research location is divided into 12 land units, based on soil type, landform, and land use for vegetable crops under clove and nutmeg stands. Field and laboratory research in each land unit to collect data based on soil health indicators. Soil samples for analysis of soil properties in the laboratory are composite samples at a depth of 0-30 cm [12]. Soil health indicators are research variables that include soil properties, among others; soil color, soil texture, soil structure, water content, slope class, soil density, soil erosion, earthworms, soil organic matter, soil pH, salinity, nitrogen (N), Phosphorus (P<sub>2</sub>O<sub>5</sub>), potassium (K) available cation exchange capacity (CEC), base saturation (BS), and aluminum saturation. The properties of plants, among others; are ground cover and plant appearance[13];[14]. Evaluation criteria for each indicator refer to the value of each element as an indicator of soil health[12]. The value of each indicator is used as a percentage score for soil health class based on soil performance as follows: I. 81-100% Very High (Very Fertile); II. 61-80% High (Lush); III. 41-60% Moderate (Moderately Fertile); IV. 21-40% Low (Less Fertile); V. 0-20% Very Low (Infertile) [12].

**Table 1. Assessment criteria and soil health indicator scores**

Number	Land Property	(score 1)	(score 2)	(score 3)	(score 4)	(score 5)
1	Soil color*&****	red, gray, greenish gray	yellow, reddish yellow, yellowish red	Yellowish brown, reddish brown	dark brown to very dark brown	black
2	Water content *&****	<10%, >80%	10-20%, 70-80%	20-30%, 60-70%	30-40%, 50-60%	40-50%
3	Slope*	>30%	15-30%	8-15%	3-8%	0-3%
4	Soil texture*&****	S, C	LS, SC, SiC	SL, SCL	Si, SiCL, CL	L, SiL
5	Soil structure*&****	Details	plate	Pole, prism	cubes/clumps	Crumbs, granules
6	Organic matter (C-organic)**	none (<1)	a little (1-2)	enough (2-3)	many (3-5)	very much (>5)

7	pH (H <sub>2</sub> O)**	<4,5	4.5-5.5	7.6-8.5	5,5-6	6-7,5
8	CEC (cmol (+)kg <sup>-1</sup> )**	<5	5,1-6,9	7.0-24	25-40	>40
9	Basic saturation (KB)**	<20	20-40	41-60	61-80	>80
10	Al saturation (%)* *	<5	5.0-10	10,1-20	21-40	>40
11	Total nitrogen (%)* *	<0.1	0.1-0.2	0.2-0.5	0.5-0.75	>0.75
12	Phosphorus/P <sub>2</sub> O <sub>5</sub> (ppm)**	<4	5.0-7	8.0-10	11.0-15	>15
13	Available potassium (cmol (+)kg <sup>-1</sup> )	<0.1	0.1-0.3	0.4-0.5	0.6-1.0	>1.0
14	Salinity (dS /m)**	<1	1-2.0	2,1-3	3,1-4	>4
15	Earthworms***	none (0)	little, wormhole shit (1-2)	enough, wormhole shit (3-5)	Lots of wormhole shit (6-9)	Abundant wormhole excrement (≥10)
16	Groundcover plants***	<45%	45-64%	65-74%	75-99%	100%
17	Avalanche***	big moat	small ditch	channel	sheet	No erosion
18	Soil density***	hard, dense, very poor root penetration	hard, solid	Firm, limited root penetration	leave	Free root penetration
19	Plant appearance***	white leaves, stunted, elemental stress	dwarf, elemental stress	moderate growth, slight elemental stress	green leaf stress-free element	Green leaves, normal growth, free from elemental pressure

Source: \*[14]; \*\*[12]; \*\*\*[13]; \*\*\*\* Research modification

### 3. RESULTS AND DISCUSSION

#### 3.1 Climatic Condition

The climate classification according to Schmidt and Ferguson (1951) is classified as a wet climate type (B) with an average dry month (SW < 60 mm) of 2.1 months and an average wet month (BB > 100 mm) of 9.1 months. with a Q index of 23.1 %. The wet climate type (B) is usually covered with tropical rainforest vegetation. Agro-climatic zone according to Oldemanetal.(1980) classified it as zone C2with an average dry month (BK <100 mm) of 3 months and an average wet month (BK > 200 mm) of 6 months. According to[15], the monthly average rainfall on Ternate Island ranges from 115.6 mm to 263.5 mm with an annual rainfall of 2,233 mm/year. Rainy days range from 12 to 20 days with an average of 16 days. The rainiest days in May and December are 20 days. The maximum monthly air temperature ranges from 28 .8 °C to 31.9 °C with an annual average of 31.1 °C.



12	Phosphorus/P <sub>2</sub> O <sub>5</sub> (ppm)	5	5	4	4	4	4	4	4	3	3
13	Available potassium (cmol (+)kg <sup>-1</sup> )	2	2	2	2	2	2	2	2	2	2
14	Salinity (dS/m)	1	1	1	1	1	1	1	1	1	1
15	Earthworm population	2	2	2	2	1	2	2	1	3	3
16	Ground cover crops (%)	4	4	3	3	3	3	3	3	5	5
17	Avalanche	4	4	3	3	2	3	3	2	5	5
18	soil compaction	5	5	5	5	4	5	5	4	5	5
19	plant appearance	4	4	4	4	3	4	4	3	5	5
<b>Total score</b>		<b>65</b>	<b>65</b>	<b>58</b>	<b>59</b>	<b>55</b>	<b>58</b>	<b>59</b>	<b>55</b>	<b>64</b>	<b>64</b>

Based on the results of the analysis of soil health indicators in Table 1, the soil health level at the study site was classified as moderate at SST 3, 4, 5, 6, 7, and 8 according to the criteria for soil health and fertility by [12]. The moderate level of soil health on vegetable farming land in Tobololo Village is generally influenced by the low availability of total N, and the percentage of base saturation in the soil. As the results of research by [21], total N and base saturation affect soil quality in coconut plantations. The availability of C-organic and available K in the soil also determines the level of soil health value. Organic matter sourced from C-organic as a source of soil nutrients will be the main variable for assessing the level of soil health on agricultural land. This is because it affects the development of soil microorganisms. In the opinion of [22]; [23], the addition of organic matter to the soil in addition to increasing soil fertility and improving the physical structure of the soil, also functions to restore the microbiological balance in the soil.

### **3.2.2 High Grade**

Soil health levels with high value (64-65%) are found in SPL 1, 2, 9, and 10. High soil health values are because the soil surface is 100% covered with vegetation so that soil erosion does not occur and soil texture and structure are good, i.e. clay soil texture and crumb structure with loose consistency and the activity of soil organisms (earthworms) are sufficiently available. This is because there are 1-3 earthworms in each SST. Based on the research results of [24] that the level of earthworm density in vegetable fields in Tobololo Village is moderate. The level of soil health is high but in some soil chemical properties, it is still low, especially the availability of potassium and the percentage of Base Saturation (KB). The high and low values of the health indicators that affect the health level of the soil for each SST in the study location are shown in Table 2. In general, the results of this study indicate that both high and moderate health levels have problems with soil chemical indicators, namely the availability of potassium and the level of base saturation (BS).

## **4. CONCLUSIONS**

Soil health level in vegetable fields under clove stands was moderate (55-59%) and high (64-65%). The results of this study indicate that both moderate and high health levels have problems with soil chemical indicators, namely the availability of potassium and nitrogen and the level of base saturation (KB). Potassium (K) and cow manure contain lots of nitrogen (N).

## **REFERENCES**

1. Suparman, Bahtiar, Nurhasanah, & DAS, SS (2018). Diversity and Mapping Clove Varieties (*Syzygium aromaticum*) on Hiri Island, In Ternate Municipality. *Journal of Agriculture and Veterinary Science*, 11 (8), 11–16.
2. Harini, R., Yunus, HS, Kasto, & Hartono, S. (2013). Agricultural Land Conversion: Determinants and Impacts for Food Sufficiency in Sleman Regency. *Indonesian Journal of Geography*, 44 (2), 120–133.
3. Hossaimah, H., & Subari, S. (2017). Acceleration of the Transfer of Functions (Conversion) of Agricultural Land to Non-Agriculture in the District of Galis, Pamekasan Regency. *AGRISAINTIKA: Journal of Agricultural Sciences*, 1 (2), 97.
4. Paudel, MN (2016). Multiple Cropping for Raising Productivity and Farm Income of Small Farmers. *Journal of Nepal Agricultural Research Council*, 2 (December), 37–45. <https://doi.org/DOI>:
5. Umasugi, B., Teapon, A., & Sudjud, S. (2022). Food in the Garden, Island Farmer Farming Model: Analysis of land characteristics and income of farmers on Ternate Island. *Agro Bali: Agricultural Journal*, 5 (2), 342–348.
6. Kurniawan, A., Nurdin, AS, Fatrawana, A., Tamnge, F., Ashari, R., Utami, NN, Hadun, R., Marasabessy, MH, & Partnerships, E. (2023). Strategy for Sustainable Livelihood of Communities Around Forest Area on Ternate and Hiri Island, North Maluku. *International Journal for Research Trends and Innovation*, 8 (3), 505–512.
7. Doran, JW, & Zeiss, MR (2000). Soil health and sustainability: managing the biotic component of soil quality. *Applied Soil Ecology*, 15 (June).
8. Knight, A., Cheng, Z., Grewal, SS, Islam, KR, Kleinhenz, MD, & Grewal, PS (2013). Soil health as a predictor of lettuce productivity and quality: A case study of urban vacant lots. *Urban Ecosystems*, 16 (3), 637–656.
9. Moebius-Clune, BN, Es, HM, Idowu, OJ, Schindelbeck, RR, Moebius-Clune, DJ, Wolfe, DW, Abawi, GS, Thies, JE, Gugino, BK, & Lucey, R. (2008). Long-Term Effects of Harvesting Maize Stover and Tillage on Soil Quality. *Soil Science Society of America Journal*, 72 (4), 960–969.
10. Guo, M. (2021). Soil health assessment and management: Recent development in science and practices. *Soil Systems*, 5 (4).
11. Mulyono, A., Suriadikusumah, A., Harriyanto, R., & Djuwansah, MR (2019). Soil quality under agroforestry tree pattern in upper Citarum watershed, Indonesia. *Journal of Ecological Engineering*, 20 (1), 203–213.
12. Balitan, BPT (2005). Chemical Analysis of Soil, Plants, Water, and Fertilizers. In *Agricultural Research and Development Agency Ministry of Agriculture*.
13. Sundermeier, A., Educator, E., & County, W. (2004). *Ohio Soil Health Card: Sustainable Agriculture Fact Sheet Series*. Ohio State University Extension. [nrcs.usda.gov/wps/portal/nrcs/site/soils/home/](https://nrcs.usda.gov/wps/portal/nrcs/site/soils/home/); [southcenters.osu.edu/soil-and-bioenergy/](https://southcenters.osu.edu/soil-and-bioenergy/); [ohioline.osu.edu](https://ohioline.osu.edu).
14. Riwandi, ., & Handajaningsih, M. (2011). Relationship between Soil Health Assessment and the Growth of Lettuce. *Journal of Tropical Soils*, 16 (1), 25–32.
15. Umasugi, B., Prijono, S., Soemarno, & Ariffin. (2021). Soil Moisture Analysis and Rainwater Management at Clove Plantation to Meet the Water Requirements of Clove Plants During the Dry Season. *International Journal on Advanced Science, Engineering and Information Technology*, 11 (6), 2197–2203.
16. Soil Survey Staff, U. [United SD of A. (2015). *Soil Taxonomy Keys*. Research Center and Center for Research and Development of Agricultural Land Resources, Agency for Agricultural Research and Development, Ministry of Agriculture.
17. Umasugi, B., & Kamaluddin, AK (2015). Effect of Gamalama Volcanic Ash on Agricultural Plants in Ternate City. *Proceedings of the National Seminar on Outstanding Higher Education Research Results*, 1, 289–291.

18. Hadun, R., Rayes, M.L., Munir, M., & Prijono, S. (2016). Characterization of Land Resources in the Clove Plantation Area in Ternate Island, North Maluku, Indonesia. *IOSR Journal of Agriculture and Veterinary Science (IOSR-JAVS)*, 9 (2), 1–7.
19. Sudjud, S., & Hadun, R. (2018). The Potential Areas for Crop Development in Morotai Island Regency, Indonesia. *International Journal on Advanced Science, Engineering and Information Technology*, 8 (6), 2374–2379.
20. Shadikin Nurdin, A., Kurniawan, A., & Hadun, R. (2020). Assessment of land evaluation on erosion reduction with Agroforestry approach around Lake Ngade area. *IOP Conference Series: Earth and Environmental Science*, 499 (1).
21. Irawan, S., Antriandarti, E., & Laia, D.H. (2022). A Study of the Relationship of Soil Quality with Land Suitability for Coconut (*Cocos nucifera* L.) Development in Kebonagung Sub-district, Pacitan District. *IOP Conference Series: Earth and Environmental Science*, 1111 (1).
22. Yulianti, T. (2010). Organic Matter: Its Role in Soil Health Management and Control of Soil-Transmitted Pathogens Towards Organic Tobacco Farming. *Bulletin of Tobacco Plants, Industrial Fibers & Oils*, 2 (1), 26–32.
23. Mann, C., Lynch, D., Fillmore, S., & Mills, A. (2019). Relationships between field management, soil health, and microbial community composition. *Applied Soil Ecology*, 144 (December 2018), 12–21.
24. Umasugi, B., Teapon, A., & Ishak, L. (2022). Earthworms; Soil Health Indicators in Soil Management for Vegetable Cultivation. *National Agribusiness Seminar*, 2 (1), 147–151.