

## Original Research Article

# Acute Toxicity and Anti-Anxiety Activity Screening of Hydroalcoholic Leaf Extracts of *Bryophyllum pinnatum*, *Terminalia catappa* and *Tapinanthus dodoneifolius* Growing on *Terminalia catappa* Tree

### ABSTRACT

**Aims:** Hydroalcoholic *T. dodoneifolius*, *B. pinnatum* and *T. catappa* leaf extracts were investigated for their acute toxicity and anxiolytic activities. **Study design:** Acute toxicity (LD50) was determined using the limit dose acute and Anxiolytic activities were assessed by open-field behavioural testing in mice. **Place and duration of study:** The study took place in the Neurobehavioural room of Department of Pharmacology & Therapeutics, Faculty of Pharmacy, Ahmadu Bello University, Zaria, in late December 2022. **Methodology:** Groups of mice (n=6; equal sexes) were each exposed to the open-field paradigm (OFT) following 1 hour of single oral administration of distilled water, extracts (125, 250, and 500 mg/kg) and diazepam (0.5 mg/kg) using the behavioural indices of mean centre zone time (M% CZT), centre zone re-entry (MCZR) and defecation/urination frequency (MD/UF). **Results:** Acute toxicity testing shows that the three extracts are safe with LD50 of 5g/kg. Behavioural results indicate, compared to the distilled water treatment (M% CZT, 6.39±1.53; MCZR, 4.67±1.15, & MU/UF, 4.83±0.75), single acute *T. dodoneifolius* (M% CZT, 7.50±1.73, 13.72±2.43, & 20.94±3.91\*; MCZR, 7.33±0.88, 9.50±0.76, & 11.50±1.6\*; MD/UF, 4.17±0.48, 3.17±0.54, & 1.83±0.60\*), and *B. pinnatum* (M% CZT, 9.72±1.91, 9.78±1.32, & 19.67±2.01\*; MCZR, 6.00±1.15, 10.50±0.76, & 12.50±1.72\*; MD/UF, 3.17±0.79, 2.33±0.42, & 1.33±0.49\*) extract treatments demonstrated consistent dose-dependency in their anxiolytic activity across the three parameters with *T. catappa* extract (M% CZT, 6.95±1.27, 12.17±2.01, & 16.84±1.49\*; MCZR, 8.17±1.60, 10.17±1.67, & 10.83±1.96; MD/UF, 2.83±0.60, 2.00±0.73, & 3.67±0.42) exhibiting dose-dependent anxiolytic effect only on the M% CZT. However, at the highest dose level of 500 mg/kg, all plant extract treatments caused anxiolytic effects that are comparable with those of 0.5 mg/kg diazepam treatments (M% CZT, 22.61±1.31\*; MCZR, 12.17±1.66\*; & MD/UF, 2.62±0.21\*). **Conclusion:** These findings justify the traditional use of these medicinal plant extracts in the remediation of nervous and related disorders.

**Keywords:** Acute, Anxiolytic, Open-field test, Centre zone time, Centre zone re-entry, Mice, Defecation, Urination

## 1. INTRODUCTION

Anxiety disorders are chronic psychobehavioural states that are characterized by inappropriate, inexplicable and/or excessive fearfulness, tension, and irritability – all of which may be disabling to the victims [1; 2; 3]. The various forms of these mood disorders – generalized anxiety disorder (GAD), obsessive-compulsive disorder (OCD), panic anxiety disorder (PD), social anxiety disorder (SAD) and post-traumatic stress disorder (PTSD) – collectively constitute a high disease prevalence and socio-economic burden [4; 5; 6]. This scenario is heightened by the paucity, the toxicity and effectiveness concerns of the currently available anti-anxiety therapeutic agents [7; 8]. ~~To~~ Therefore, to both meet the increasing demand ~~of for~~ these disorders and mitigate the afore-mentioned anti-anxiety therapeutic and clinical liabilities, there is a need to discover new agents - preferably with novel anxiolytic mechanism(s) to ~~complement~~ complement and/or substitute the traditional benzodiazepine- and serotonin- related drugs [9; 10; 11].

The plant kingdom has been a limitless source of new pharmacophores for diverse disease spectra [12; 13]. Anti-anxiety agents are not an exception to this – with anxiety-relieving activity and the presence of phytochemicals reported for some medicinal plants, including *Hydrocotyle umbellata* L., *Ginkgo biloba*, *Piper methysticum*, *Camellia sinensis*, *Leonurus cardiac*, *Valeriana officinalis*, *Withania somnifera*, *Nymphaea alba*, *Passiflora incarnata*, *Commelina benghalensis*, *Turnera aphrodisiaca* and *Crataegus oxyacantha* in both animal and human studies [14; 15; 16]. There is a need, therefore, to search the plant kingdom further to discover new anxiolytic agents. Thus, based on the afore-mentioned ethno-medicinal grounds, this study set out to assess the anti-anxiety potential activity of leaf extracts of *B. pinnatum*, *T. catappa* and *T. dodneifolius* growing on *T. tree*.

*B. pinnatum* (Lam.) Pers. (syns: *Bryophyllum calcicola*, *Kelanchoe pinnata*, *Bryophyllum calcinum*, *Cotyledon calcina*, *Bryophyllum germinans*, etc.) commonly known as miracle leaf, air plant, life plant, Goethe plant and cathedral bells, and of the family Crassulaceae, is a plant that is now naturalized to the tropical and subtropical regions of the world [17]. This perennial plant, ~~that which~~ is increasingly becoming an ornamental item in Nigerian households, has a fleshy cylindrical stem that can grow up to 1 meter tall with succulent, fleshy, elliptical-shaped, curved and margin-serrated deep green leaves [18].

Formatted: Justified



Plate 1: *Bryophyllum pinnatum*

Reports from Nigeria and other parts of the world show *B. pinnatum* and its congeners are used ethno-medicinally for the treatment of several illnesses including gastric ulcer, hypertension, smallpox, cancers, migraines, diabetes mellitus, psychosis, colds, dysmenorrhea, hepatitis, diarrhea and dysentery, ear infection, kidney stones, conjunctivitis, retained placenta, chicken pox, rheumatoid arthritis, pulmonary infection, heart failure and palpitation [19; 20; 21; 22; 23; 24].

Reported pharmacological activities of extracts of *B. pinnatum* and its congeners include anti-hypertensive, hypoglycemic, nephroprotective,

hepatoprotective, antibacterial, immunosuppressive, analgesic, anti-inflammatory, insecticidal and anti-carcinogenic effects [24; 25; 26; 27; 28; 29; 30; 31; 32]. These biological activities are viewed to be due to the presence of several important phytochemicals (4-hydroxy-3-methoxy-cinnamic acid, bufadienolide, protocatechuic acid, epigallocatechin-3-o-syringate, quercetin, 4',5-dihydroxy-3',8-dimethoxy flavone-7-O- $\beta$ -D-glucopyranoside,  $\beta$ -amyrin, 22-dihydrobrassicasterol, bryophyllin A, oxalic acid, malic acid), vitamins (riboflavin, thiamine, pyridoxine, ascorbic acid, niacin), minerals (calcium, magnesium, zinc, manganese, copper, potassium, iron) and amino acids (glutamic acid, glycine, cysteine, tyrosine, casein hydrolysate, phenylalanine, protein hydrolysate, methionine) [33; 34; 35; 36] in these extracts.

Previous studies have shown extracts of *B. pinnatum* and its congeners to exert dose-dependent anticonvulsant activity, antidiabetic, anxiolytic, hypnotic and muscle relaxant effects in rodents [37; 38], and antidiabetic and anxiolytic effects in young Zebra fish [40]. However, scientific reports on the anxiolytic activity of *k. pinnata* are scanty, and this study—~~therefore, therefore,~~ aims to investigate its acute anxiolytic effect in mice in the open field test.

*Terminalia catappa* L. (syns: *Phytolacca javanica* (Osbeck); *Terminalia mauritiana* (Blanco); *Terminalia moluccana* (Lamk.); and *Terminalia procera* (Roxb)) is a medium-large sized Tropical tree plant, commonly known as Tropical almond, belonging to the Combretum family. The plant at full maturity has a thick stem with dark brown/~~gray-grey~~ fissured bark; stem branches bearing spirally arranged simple, broadly obovate, round and blunt tipped leaves; stalkless, ovoid to ovate thin-skinned fleshy fruits, each containing a seed enclosed within a hard shell; and smallish five-lobed, axillary-spiked mildly unpleasant smelling white or cream coloured male-dominated flowers [41].

In Nigeria, the fruit of this plant is consumed orally as snacks. Ethnomedicinally, different parts of *Terminalia catappa* have been widely reported to be helpful in the treatment of leprosy, headache, dermatitis, abdominal colic, fever, gastropeptic dyspepsia, hepatitis, urinary tract infection, diarrhea, heart failure and poor sexual drive [42; 43; 44; 45; 46]. ~~Extracts—In addition,~~ extracts from different parts of this plant have been reported to exhibit antidiabetic, anti-lipidemic, antineoplastic, hepatoprotective, aphrodisiac, anti-inflammatory, antibacterial, antioxidant, analgesic and anti-ageing effects [44; 45; 47; 48; 49; 50; 51; 52]. But scientific literature has scanty reports on the anxiolytic activity of extracts from *T. catappa*. Thus, this study aims to screen the aqueous methanol leaf extract of this plant for its acute anxiolytic activity employing the mouse open field test.

The third medicinal plant whose leaf extract ~~is to will~~ be investigated in this study is *Tapinanthus dodoneifolius* (DC.) DENSER (syns: *Agelanthus dodoneifolius*; *Dentimetuladodoneifolius*; *Loranthus chevalieri*; *Loranthus dodoneifolius*; *Loranthus knoblercheri*; *Loranthus uelensis* and *Scurrula dodoneifolia*) [54].



Plate 2: Terminalia catappa

*Terminalia catappa* (Almond) tree hosting *Tapinanthus dodoneifolius* epiphytes

This plant is a green leafy shrub found in most parts of Nigeria and the Tropics growing as a parasitic plant on host trees such as *Theobroma cacao*, *Tamarindus indica*, *Acacia nilotica*, *Vitellera paradoxa* (Shea butter), Kolanut, *Parkia biglobosa*, *Terminalia catappa*, *Citrus sinensis* and *Azardirachta indica*. It is known as *Viscum album* in Europe, mistletoe in English, *afomo onisana* in Yoruba, and *Kauchii* in Hausa [55; 56]. Its extracts have been reported for efficacy in diverse diseases, including hypertension, menstrual pain, diabetes mellitus, cancers, abdominal colic, gonorrhoea, malaria, diarrhoea, dysentery and peptic ulcer [57; 58; 59;

60; 61]. ~~Several~~ In addition, ~~several~~ important pharmacological activities which have been associated with extracts obtained from *Tapinanthus dodonaeifolius* and its synonyms include antimicrobial, anti-ulcerogenic, antiplasmodial, antihypertensive, anticancerous, anticonvulsant and hepatoprotective [55; 57; 58; 59; 60; 61; 62; 63]. These ~~biologic~~ ~~biological~~ activities may be due to the presence of numerous important phytochemical compounds such as anthraquinones, saponins, tannins, alkaloids, phlobatannins, ~~carbohydrate~~ ~~carbohydrates~~, flavonoids, anthracene, cardiac glycosides, steroid and triterpenes, and a dihydropyranone Dododeine [58; 59; 60; 61; 67]. Previous studies indicate a bark extract of *Tapinanthus dodonaeifolius* to have exhibited anxiolytic, antidepressant and spatial memory-enhancing ~~effect~~ ~~effects~~ and a hydromethanol leaf extract and its fractions of *Tapinanthus globiferus* epiphytic on *Azadirachta indica* tree to have demonstrated an anxiolytic activity [64; 65; 66]. But scientific reports on the anxiolytic activity of the leaf extract of *Tapinanthus dodonaeifolius* are scarce. ~~This and this~~ study is to investigate the acute anxiolytic activity of *Tapinanthus dodonaeifolius* in mice using the mouse open field test.

## 2. MATERIAL AND METHODS

### 2.1 Drugs, Open-field apparatuses and Plant materials

Diazepam tablets (Roche, 5 mg) were obtained from a pharmaceutical outfit near the Ahmadu Bello University campus, Zaria. Units of 40 x 40 x 40 cm transparent plexiglass open-field apparatus and distilled water were graciously supplied by ~~the~~ Department of Pharmacology laboratory, Faculty of Pharmaceutical Sciences, Ahmadu Bello University, Zaria. *B. pinnatum* fresh leaves were harvested in ~~the~~ Gwagwalada area of the Federal Capital Territory, Abuja. Freshly fallen leaves of *T. catappa* and fresh leaves of *T. dodonaeifolius* growing on *T. catappa* were harvested on the main campus of Ahmadu Bello University, Zaria, Nigeria, all in December 2022. Following collection, all leaves were each separately briskly washed to remove dirt and contaminants. ~~Then,~~ *T. dodonaeifolius* and *T. catappa* leaves were separately air-dried until complete dryness while, due to the difficulty of getting it to dryness through air-drying on account of ~~its~~ ~~their~~ succulent nature, *B. pinnatum* leaves were first pulverized to fine texture using an electrical blender, following which it was thinly spread on a clean flat inert white surface and air-dried by the aid of an electrical fan until ~~the~~ constant dry weight was achieved. All dry leaves were ground to fine powders and kept in plastic containers at room temperature till use. Twenty grammes (21 g) of dry *B. pinnatum* leaves, ~~and~~ 40 g each of dry *T. dodonaeifolius* ~~dodonaeifolius~~ and *T. catappa* leaves were each soaked in 0.5 L of equal volumes of distilled water and ethanol for 24 hours, following which the macerates were separately Whatman paper-filtered. ~~The and the~~ resulting filtrates were separately air dried aided by electric fanning.

### 2.2 Animals

Healthy adult white albino Swiss mice of both sexes were sourced from the animal house of the Faculty of Pharmaceutical Sciences, Ahmadu Bello University, Zaria, Kaduna State, North-west Nigeria. They were kept in clean plastic battery cages for at least 7 days under room temperature under ~~a~~ 12-hour light/dark cycle

Formatted: Justified

with access to feeds and water *ad libitum*. Prior to and throughout this study, the mice were humanely handled under good laboratory practices and ethics.

### 2.3 Experimental design

#### 2.3.1 Acute toxicity testing

Twelve female adult mice ( $22.0 \pm 0.5$  g) were used for the acute toxicity testing. Based on the long/widespread use of and previous high acute safety reports on the selected medicinal plants [25; 27; 45; 46; 58; 60; 68], and to minimize the number of animals used, mice (n=3) were each administered an oral limit dose of an extract at 5000 g/kg/10 ml distilled water and 10 ml/kg distilled water for the control group. They were all observed acutely for toxicological signs for 24 hours, and thereafter, daily for 14 days.

#### 2.3.2 Behavioural studies

The screening for the acute anxiolytic activities of the selected leaf extracts was carried out in randomized groups of mice (n=6; an equal number of sexes) weighing  $21.0 \pm 0.8$  g in units of 40 cm square plexiglass open-field apparatus whose floors were each partitioned with gridlines into 16 equal squares. The behavioural experimentation took place in a quiet, fairly large room having about 10-lux illumination. Activity screening was done by placing a mouse orally administered 1 hour earlier with 125, 250, and 500 mg/kg extracts, 0.5 mg/kg diazepam or 10 ml/kg distilled water, in the center of the open-field and allowed to freely explore the test arena for 5 minutes according to the procedures as described in [69] – with minor modifications. The innermost central squares were taken as the centre zone, and the rest of the squares as the periphery of the test arena. Three important mouse anxiety parameters, such as percentage centre zone time, which is the percentage of the test spent by the mouse with all four paws within the central zone, centre zone re-entries and frequency of defecation/urination by the animals were all observed and recorded

## 3. RESULTS AND DISCUSSION

### 3.1 Plant extraction yields

Table I indicates the extract yields of 21 g of *B. pinnatum* and 40 g each of *T. dodoneifolius* and *T. catappa* dry leaf powders.

Table I: Percent plant extract yields

Extract	Yields (g)	% yield (g/100 g)
<i>Tapinnatus dodoneifolius</i>	4.40 (bright green)	20.10
<i>Bryophyllum pinnatum</i>	7.80 (deep green)	19.50
<i>Terminalia catappa</i>	13.40 (rich brown)	31.50

### 3.2 Acute toxicity testing results

The three plants appear to be well tolerated as the experimental animals exhibited no toxicological signs during the 24 hours following extract administration and throughout the 14-day test.

### 3.3 Behavioural experiment results

Formatted: Justified

Formatted: Justified

Formatted: Justified

Formatted: Justified

Formatted: Justified

3.3.1 Effect of single acute doses of hydroalcoholic *B. pinnatum*, *T. dodoneifolius* and *T. catappa* leaf extracts on mean centre zone times (M% CZT) of mice (n+6) exposed to the OFT.

In the open field test (Table II), compared to distilled water treatment, all the extracts caused a dose-dependent increase in the mean %CZT spent by the mice at the centre zones of the open field maze with a significant increase ( $P < 0.05$ ) in this rodent parameter at the highest doses of the extracts. The mean %CZT of  $20.94 \pm 3.91$  recorded by the experimental group treated with the highest dose of *T. dodoneifolius* was comparable to  $22.61 \pm 1.31$  recorded for the 0.5 mg/kg diazepam treated-mice.

Table II: Effect of plant extracts on mean percent centre zone times

Treatments (mg/kg)	Mean percent centre zone times (M% CZTs)		
Plant extracts	125.00	250.00	500.00
D/water (10 ml/kg)	$6.39 \pm 1.53$	$6.39 \pm 1.53$	$6.39 \pm 1.53$
<i>T. dodoneifolius</i>	$7.50 \pm 1.73$	$13.72 \pm 2.43$	$20.94 \pm 3.91^*$
<i>B. pinnatum</i>	$9.72 \pm 1.91$	$9.78 \pm 1.32$	$19.67 \pm 2.01^*$
<i>T. catappa</i>	$6.95 \pm 1.27$	$12.17 \pm 2.01$	$16.84 \pm 1.49^*$
Diazepam (0.5)	$22.61 \pm 1.31^*$	$22.61 \pm 1.31^*$	$22.61 \pm 1.31^*$

All results were expressed as mean  $\pm$  S.E.M of mice (n=6). Analysis was done using the One-way ANOVA. Significance set at  $P$ -values  $\leq 0.05$ .

3.3.2 Effect of single acute doses of hydroalcoholic *B. pinnatum*, *T. dodoneifolius* and *T. catappa* leaf extracts on the mean centre zone re-entries (MCZR) of mice (n+6) exposed to the OFT.

All the plant extracts (Table III) caused dose-dependent increases in the mean frequencies of CZR, but only the highest doses of *T. dodoneifolius* and *B. pinnatum* extracts caused significant ( $P$ -values  $\leq 0.05$ ) increases in the mean CZR that are comparable to that of 0.5 mg/kg diazepam treatment.

Formatted: Justified

Formatted: Justified

Formatted: Justified

Formatted: Justified

Formatted: Justified

Formatted: Justified

Formatted: Justified

Table III: Effect of plant extracts on mean centre zone re-entries

Treatments (mg/kg)	Mean centre zone re-entries (MCZR)		
	125.00	250.00	500.00
Plant extracts			
D/water (10 ml/kg)	4.67±1.15	4.67±1.15	4.67±1.15
<i>T. dodoneifolius</i>	7.33±0.88	9.50±0.76	11.50±1.6*
<i>B. pinnatum</i>	6.00±1.15	10.50±0.76	12.50±1.72*
<i>T. catappa</i>	8.17±1.60	10.17±1.67	10.83±1.96
Diazepam (0.5)	12.17±1.66*	12.17±1.66*	12.17±1.66*

All results were expressed as mean ± S.E.M of mice (n=6). Analysis was done using the One-way ANOVA. Significance set at  $P$ -values  $\leq 0.05$ .

### 3.3.3 Effect of single acute doses of hydroalcoholic *B. pinnatum*, *T. dodoneifolius* and *T. catappa* leaf extracts on mean defecation/urination frequency (MD/UF) of mice (n=6) exposed to the OFT

Compared to the distilled water-treated mice, dose-dependent and significant ( $p \leq 0.05$ ) mean D/UF reduction was only observed with *T. dodoneifolius* and *B. pinnatum* extracts-treated mice – with the two extracts at their highest doses of 500 mg/kg exhibiting significant ( $P \leq 0.05$ ) mean D/UF reductions that even surpassed that of 0.5 mg diazepam treatment. On the other hand, *T. catappa*-treated mice exhibited dose-independent and insignificant ( $P > 0.5$ ) reduction in this anxiety parameter.

Table IV: Effect of plant extracts on mean defecation/urination frequencies

Treatments (mg/kg)	Mean defecation/urination frequencies (MD/UFs)		
	125.00	250.00	500.00
Plant extracts			
Dist. water (10 ml/kg)	4.83±0.75	4.83±0.75	4.83±0.75
<i>T. dodoneifolius</i>	4.17±0.48	3.17±0.54	1.83±0.60*
<i>B. pinnatum</i>	3.17±0.79	2.33±0.42	1.33±0.49*

<i>T. catappa</i>	2.83±0.60	2.00±0.73	3.67±0.42
Diazepam (0.5)	2.62±0.21*	2.62±0.21*	2.62±0.21*

All results were expressed as mean ± S.E.M of mice (n=6). Analysis was done using the One-way ANOVA. Significance set at  $P$ -values  $\leq 0.05$ .

Extract yields of dry leaf powers of these medicinal plants in this study are relatively high (Table I). *T. globiferus*, *B. pinnatum*, and *T. catappa* extracts have been associated with efficacy for several diseases in diverse traditional cultures [19; 42; 56], and the observed high extract yields of these medicinal plants may be an indication of the rich deposits of diverse bioactive phytoconstituents that have been reported for extracts obtained from these plants in previous studies [17; 20; 48; 50; 58; 60].

Despite previous reports of a rich repository of phytochemicals in the extracts of these plants, the relatively high safety exhibited by these extracts are in agreement with some earlier reports of high tolerability for *T. dodoneifolius* DC. (Loranthaceae) [58; 61], *T. catappa* [67; 68] and *B. pinnatum* [70; 71] extracts. Their individual apparent wide margin of tolerability is well supported by the long and widespread ethnomedicinal uses of the three plants for the traditional alleviation of diverse diseases, including infections, hypertension, diabetes mellitus, convulsions, cough, and cancers with scarce or no reports of significant toxicity [18; 19; 42; 52; 57].

OFT has remained one of the most relevant preclinical psychotropic drug discovery tools since its invention close to a century ago by Carl S. Hall to assess rodents' general locomotory and anxiety-related emotional behaviours [72; 73]. This test is based on the conflict created in the animal between its natural tendency to explore and aversion to exposure or open space when placed in the open-field maze [76; 77; 78]. ~~In previous studies, this test has been deployed in the anxiolytic and sedative-anxiolytic activity screening of several medicinal plants. This test has been deployed in the anxiolytic and sedative-anxiolytic activity screening of several medicinal plants in previous studies~~ [80; 81; 82; 83]. Its attraction is ~~derived from its operational simplicity, cost-effectiveness and relative high throughput when compared to other standard tests used in characterizing ethological patterns in animals and screening~~ viewed to derive from its operational simplicity, cost-effectiveness and relative high throughput when compared to other standard tests used in characterizing ethological patterns in animals and in the screening of anxiolytic or anxiogenic activity in both putative and established anxiety-modulatory agents [74; 75]. Its other strengths include ~~the~~ capacity to generate multiple anxiety-related parameters, ~~and~~ the affordability of temporal assessment of these parameters. ~~high~~ ~~High~~ inter-parameter correlation and good predictive validity for full benzodiazepine and 5HT1A agonists [76; 77; 78]. The plethora of rodent anxiety-related parameters that can be harvested on the OFT include percent centre zone time (proportion of test time spent in the central zone of field: the greater the percentage, the less anxiogenic the animal is); centre zone entries (frequency of central zone entry and re-entries: higher values indicates anxiolysis); latency to centre zone entry

Formatted: Justified

Formatted: Justified

Formatted: Justified

(time delay before the animal enters the central zone of the maze, short latency indicates anxiolytic effect); ~~defecation~~defecation/urination frequency (higher ~~faecal~~faecal/ urine droppings indicate anxiogenicity) and a number of rears (a rodent event during which the animal stands erect on its hindlimbs with its hindlimb either completely in the air (unsupported rear) or rests ~~them—them~~against the walls (supported rear) of the maze – this parameter is viewed to mirror either anxiogenesis or anxiolysis depending if the rear is supported or not). Other parameters that can be obtained in the OFT are grooming (a rodent behavioural event in which the animal engages in self-cleaning, licking and brushing of face and body; it is said to indicate high anxiety level); stretch attend postures (the animal while lowering and keeping its trunk close to the floor of the test apparatus, alternately elongates and retracts it in order to assess the risks in its immediate environment – this parameter indicates anxiety); a number of squares crossed (this is a measure of the locomotory and exploratory activities which is indicative of anxiolysis) and freezing (the animal pulls itself together and remains immobile usually close to the walls of the test apparatus – it indicates high anxiety levels) [76; 77; 78; 79].

The anxiolytic activity screening of hydroalcoholic *T. globiferus*, *B. pinnatum*, and *T. catappa* extracts in the OFT was assessed using the percent centre zone time, centre zone entries and ~~defecation~~defecation/urination frequency. On the mean percent centre zone time, all ~~the~~three plant extracts demonstrated dose-dependent anxiolytic activity that was significant at their highest doses and with the *T. dodoneifolius* extract's anxiolytic activity comparable to the standard anxiolytic drug. However, on the mean centre zone re-entries and mean defecation/urination frequencies, only *T. dodoneifolius* and *B. pinnatum* extracts exhibited dose-dependent anxiolytic effects that were significant and comparable to those of 0.5 mg/kg of diazepam - the standard anxiolytic drug. This study is one of the few reports on the anxiolytic activity of these plants despite their widespread traditional use for the alleviation of a broad spectrum of diseases [27; 32; 46; 47; 57; 63; 67].

These ~~finding~~findings are in agreement with previous reports of similar anxiolytic, antidepressant and cognition-enhancing activities of *T. dodoneifolius* extracts [64] and anxiolytic activity of its congener *T. globiferus* extracts [65; 66] in mice activity. They are also similar to previous ~~reported~~reports of anxiolytic activity of *B. pinnatum* and its congener *K. pinnata* extracts in young zebrafish and rodents [40; 84]. Anxiolytic activity reports on *T. catappa* were scanty, but it has been reported to exhibit significant antidepressant effect in rodents subjected to chronic unpredictable stress [85].

Hydroalcoholic *T. dodoneifolius* and *B. pinnatum* extracts have demonstrated consistent anxiolysis across the three rodent anxiety parameters, and *T. catappa* has less consistency in this study. Use of single acute as opposed to repeated acute doses of the extracts coupled with the use of only the OFT (reportedly less sensitive to the anxiolytic activity of selective serotonin-reuptake inhibitors [86] and anxiety-related behaviours in some rodent genetic strains [87]) as a sole paradigm in this study, may have reduced the size and consistency of our findings.

#### 4. CONCLUSION

~~This study has shown hydroalcoholic T. dodoneifolius, B.pinnatum and T. catappa extracts possess anxiolytic properties. On the whole, this study has shown hydroalcoholic T. dodoneifolius, B.pinnatum and T. catappa extracts possess anxiolytic property.~~ This finding justifies their traditional use in the alleviation of nervous and related disorders. However, further studies are recommended to finetune the present findings.

Formatted: Justified

#### CONSENT

Consent not applicable

Formatted: Justified

#### ETHICAL APPROVAL

Ethical approval for the conduct of this study was sought and obtained from the Ethical committee of the Ahmadu Bello University, Zaria.

#### REFERENCES

- [1] Adwas A. A., Jbireal J. M., & Azab A. E. (2019). Anxiety: Insights into signs, symptoms, etiology, pathophysiology, and treatment. *East African Scholars Journal of Medical Sciences*, 2(10), 580-591.
- [2] Noyes Jr R. & Hoehn-Saric R. (1998). *The anxiety disorders*. New York: Cambridge University Press. Pp. 343.
- [3] Zinbarg, R. E., & Barlow, D. H. (1996). Structure of anxiety and the anxiety disorders: A hierarchical model. *Journal of Abnormal Psychology*, 105 (2): 181–193.
- [4] Bandelow B. & Michaelis S. (2015). Epidemiology of anxiety disorders in the 21st century. *Dialogues in clinical neuroscience*, 17(3), 327–335.
- [5] Stein D. J., Scott K. M., de Jonge P., Kessler R. C. (2017). Epidemiology of anxiety disorders: from surveys to nosology and back. *Dialogues Clin Neurosci*.19 (2):127-136.
- [6] Clark D. M. (1999). Anxiety disorders: why they persist and how to treat them. *Behav Res Ther*. 37 Suppl 1: S5-27.
- [7] Farach F. J., Pruitt L. D., Jun J. J., Jerud A. B., Zoellner L. A., & Roy-Byrne P. P. (2012). Pharmacological treatment of anxiety disorders: current treatments and future directions. *Journal of anxiety disorders*, 26 (8), 833–843.
- [8] Melaragno A. J. (2021). Pharmacotherapy for Anxiety Disorders: From First-Line Options to Treatment Resistance. *Focus (Am Psychiatr Publ)*.19 (2): 145-160.
- [9] Sanabria E., Cuenca R. E., Estes M. Á., & Maldonado M. (2021). Benzodiazepines: Their Use either as Essential Medicines or as Toxic Substances. *Toxics*, 9 (2): 25.
- [10] Argyropoulos S. V., Sandford J. J., Nutt D. J. (200). The psychobiology of anxiolytic drug. Part 2: Pharmacological treatments of anxiety. *Pharmacol Ther*. 88 (3): 213-27.
- [11] Cryan, J. F., & Sweeney, F. F. (2011). The age of anxiety: role of animal models of anxiolytic action in drug discovery. *British journal of pharmacology*, 164 (4): 1129–1161.
- [12] Katiyar C., Gupta A., Kanjilal S., & Katiyar S. (2012). Drug discovery from plant sources: An integrated approach. *Ayu*, 33 (1): 10–19.
- [13] Newman D. J. and Cragg M. G. (2020). *Journal of Natural products*. 83 (3): 770-803.
- [14] Gabriel de Oliveira M., Kelle da Silva Moreira L., Turones L. C., de Souza Almeida D., Martins A. N., Silva Oliveira T. L., Barreto da Silva V., Borges L. L., Costa E. A., Realino de Paula J. (2021).

Formatted: Justified

Mechanism of action involved in the anxiolytic-like effects of Hibalactone isolated from *Hydrocotyle umbellata* L. J Tradit Complement Med.12 (4): 318-329.

[15] Khan A., Akram M., Thiruvengadam M., Daniyal M., Zakki S. A., Munir N., Zainab R., Heydari M., Mosavat S. H., Rebezov M. & Shariati M. A. (2022). Anti-anxiety Properties of Selected Medicinal Plants. *Curr Pharm Biotechnol.* 23 (8):1041-1060.

[16] Thakur P. & Rana A. C. (2013). Anxiolytic potential of medicinal plants. *Int J Nutr Pharmacol Neurol Dis.* 3 (4): 325-31.

[17] The Plant list (2022). World Flora Online, id: wfo-0000357656.

[18] Al-Snafi A.E. (2013). The Chemical Constituents and Pharmacological Effects of *Bryophyllum calycinum*. A review. *International Journal of Pharma Sciences and Research (IJPSR)*, vol. 4, n o 12. Pp. 171-176.

[19] Rastogi R. and Mehrotra B.(1990). "Compendium of Indian Medicinal Plants," Central Drug Research Institute, Lucknow, Vol. 1, pp. 388-389.

[20] Pattewar S. V. (2012). *Bryophyllum pinnatum*: Phytochemical and Pharmacological Profile. *International Journal of Phytopharmacy.* Vol. 2 (1), pp.1-8.

[21] Recknagel, R. O. (1967). Carbon tetrachloride hepatotoxicity. *Pharmacological Reviews*, 19 (2): 145-208.

[22] Quazi M., Sayyed N., Sheikh S., Gomase P. & Choudhari A. (2011). Phytochemical analysis of chloroform extract of roots of *Kalanchoe pinnata* by HPLC and GCMS. *Int J Pharm Sci Res.* 14 (4): 1693-9.

[23] Plangger N., Rist L., Zimmermann R. & von Mandach U. (2006). Intravenous tocolysis with *Bryophyllum pinnatum* is better tolerated than beta-agonist application. *Eur J Obstet Gynecol Reprod Biol.* 124 (2):168-72.

[24] Yamagishi T, Haruna M., Yan X. Z., Chang J.J., and Lee K.H. (1989). Antitumor agents, 110, Bryophyllin B, A Novel Potent cytotoxic Bufadienolide from *Bryophyllum Pinnatum*. *J. Nat. Prod.* 1989, 52 (5): 1071--1079.

[25] Yadav N. P. and Dixit V. K. (2003). Hepatoprotective activity of leaves of *Kalanchoe Pinnata* Pers.. *Journal of Ethnopharmacology.* 86 (2-3): 197-202.

[26] <http://www.rain-tree.com/coirama.html>.

[27] Salahdeen H.M. and Yemitan O.K. (2006). Neuropharmacological Effects of Aqueous Leaf Extract of *Bryophyllum Pinnatum* in Mice. *African Journal of Biomedical Research.* 9 (2): 101-107.

[28] Obaseki-Ebor E.E. (1985). Preliminary report on the invitro antibacterial activity of *Bryophyllum pinnatum* leaf juice. *Afr J Med Med Sci.* 14 (3-4): 199-202.

[29] Supratman U., Fujita T., Akiyama K., Hayashi H., Murakami A., Sakai H., Koshimizu k. & Ohigashi, H. (2001). Anti-tumor Promoting Activity of Bufadienolides from *Kalanchoe pinnata* and *K. daigremontiana*× *butiflora*. *Bioscience, biotechnology, and biochemistry*, 65 (4): 947-949.

[30] [http://en.wikipedia.org/wiki/Kalanchoe\\_pinnata](http://en.wikipedia.org/wiki/Kalanchoe_pinnata).

[31] Rossi-Bergmann B., Costa S. S., Borges M. B. S., da Silva S. A., Noletto G. R., Souza M. L. M., and Moraes V. L.G. (1994). Immunosuppressive effect of the aqueous extract of *Kalanchoe Pinnata* in mice. *Phytothera. Res.* 8 (7): 399-402.

[32] Siddharta P. and Chaudhuri. A.K.N. (1992). Further studies on the Anti-inflammatory profile of the Methanolic Fraction of the fresh leaf extract of *Bryophyllum Pinnatum*. *Fitoterapia.* 63 (5): 451-459.

[33] Supratman U., Fujita T., Akiyama K., and Hayashi H. (2000). New insecticidal bufadienolide, Bryophyllin C from *Kalanchoe pinnata*. *Biosci Biotechnol Biochem.* 64 (6): 1310-1312.

[34] Akinpelu D.A. (2000). Antimicrobial activity of *Bryophyllum Pinnatum* leaves *Fitoterapia.* 71 (2): 193-194.

[35] Okwu D.E. and Josiah C. (2006). Evaluation of the chemical composition of two Nigerian medicinal plants. *African Journal of Biotechnology.* 5 (4): 357--361.

[36] Toshihiro A., Koke W. C. M. C., Toshitake T. and Taro M (1991). Sterols of *Kalanchoe pinnata*. First report of the isolation of both C-24 epimers of 24-Alkyl-A25-sterol from higher plants. *Lipids.* 26 (8): 660-665.

- [37] Mora-Pérez A. & Hernández-Medel Mdel R. (2016). Anticonvulsant activity of methanolic extract from *Kalanchoe pinnata* Lam. stems and roots in mice: A comparison to diazepam. *Neurologia*. 31(3): 161-8.
- [38] Yemitan O. K. & Salahdeen H. M. (2005). Neurosedative and muscle relaxant activities of aqueous leaf extract of *Bryophyllum pinnatum*. *Fitoterapia*. 76 (2): 187-93.
- [39] Tiwari V., Singh A. & Tiwari A. (2017). Evaluation of antidiabetic and antianxiety potential of *Kalanchoe pinnata* root standardized extracts. *Egypt Pharmaceut J*.16 (1): 24-36.
- [40] Martins Fernandes Pereira K., Calheiros de Carvalho A., André Moura Veiga T., Melgoza A., Bonne Hernández R., Dos Santos Grecco S., Uchiyama Nakamura M., & Guo S. (2022). The psychoactive effects of *Bryophyllum pinnatum* (Lam.) Oken leaves in young zebrafish. *PLoS one*, 17 (3), e0264987.
- [41] Quartey, G.A. (2022) Mechanical Properties of *Terminalia catappa* from Ghana. *Materials Sciences and Applications*, 13 (5), 334-341.
- [42] Chen P. S., Li J. H., Liu T. Y., & Lin T. C. (2000). Folk medicine *Terminalia catappa* and its major tannin component, punicalagin, are effective against bleomycin-induced genotoxicity in Chinese hamster ovary cells. *Cancer letters*, 152(2), 115-122.
- [43] Anand A. V., Divya N. & Kotti P. P. (2015). An updated review of *Terminalia catappa*. *Pharmacogn Rev*. 9 (18): 93-8.
- [44] Ratnasooriya W. D. & Dharmasiri M. G. (2000). Effects of *Terminalia catappa* seeds on sexual behavior and fertility of male rats. *Asian J. Androl*. 2 (3): 213-219.
- [45] Ahmed S. M., Swamy B. M. V., Dhanapal P. G. R. & Chandrashekar V. M. (2005). Anti-diabetic activity of *Terminalia catappa* Linn. leaf extracts in alloxan-induced diabetic rats. *Iran. J. Pharmacol. Therapeut*. 4 (1): 36-39.
- [46] Lim T. K. (2012). *Edible Medicinal and Non-Medicinal Plants*, Vol. 2. Springer, New York, 143-157.
- [47] Akinsanya O. B., Ayodele P. F., Onifade O. F. & Salimom M. O. (2021). EXTENUATING EFFECTS OF *Terminalia catappa* LEAVES AND *Persea americana* SEED EXTRACTS ON STREPTOZOTOCIN-INDUCED LIPIDS PERTURBATION AND PANCREATIC DAMAGE IN DIABETIC RATS. *Open Journal of Bioscience Research (OJBR)*. Volume: 2; Issue: 1, Pages: 01-09.
- [48] Fan YM, Xu LZ, Gao J, Wang Y, Tang XH, Zhao XN, et al. Phytochemical and anti-inflammatory studies on *Terminalia catappa*. *Fitoterapia*. 75 (3-4): 253-60.
- [49] Gao J., Tang X., Dou H., Fan Y., Zhao X. & Xu Q. (2004). Hepatoprotective activity of *Terminalia catappa* L. leaves and its two triterpenoids. *J Pharm Pharmacol*. 56 (11): 1449-55.
- [50] Taganna J. C., Quánico J. P., Perono R. M., Amor E. C. & Rivera W. L. (2011). Tannin-rich fraction from *Terminalia catappa* inhibits quorum sensing (QS) in *Chromobacterium violaceum* and the QS-controlled biofilm maturation and Las A staphylolytic activity in *Pseudomonas aeruginosa*. *J Ethnopharmacol*. 134 (3): 865-71.
- [51] Lin C. C., Hsu Y. F. & Lin T. C. (2001). Antioxidant and free radical scavenging effects of the tannins of *Terminalia catappa* L. *Anticancer Research*. 21 (1A): 237-243. PMID: 11299741.
- [52] Pandya N. B., Tigari P., Dupadahalli K., Kamurthy H. & Nadendla R. R. (2013). Antitumor and antioxidant status of *Terminalia catappa* against Ehrlich ascites carcinoma in Swiss albino mice. *Indian journal of pharmacology*, 45 (5): 464-469.
- [53] Wen K. C., Shih I. C., Hu J. C., Liao S. T., Su T. W. & Chiang H. M. (2010). Inhibitory Effects of *Terminalia catappa* on UVB-Induced Photodamage in Fibroblast Cell Line. *Evid Based Complement Alternat Med*. 2011:904532.
- [54] WFO (2023): *Tapinanthus dodoneifolius* (DC.) Danser. Published on the Internet; <http://www.worldfloraonline.org/taxon/wfo-0000413258>.
- [55] Ayorinde B. T., Akanji M. A. and Yakubu M. T. (2008). Alterations in some marker enzymes of liver and kidney damage following chronic administration of aqueous extract of *Tapinanthus globiferus* leaves to rats., *Pharmacognosy Magazine*, 4(15): S9- S14.
- [56] Raji I. A., Chaskda, A. A., Manu S. A., & Downs C. T. (2021). Bird species use of *Tapinanthus dodoneifolius* mistletoes parasitising African locust bean trees *Parkia biglobosa* in Amurum Forest Reserve, Nigeria. *Journal of Ornithology*, 162 (4), 1129 - 1140.

- [57] Burkill H. M. (1995). The useful plants of west tropical Africa. Vol. 1-3. No. 2 Edition.
- [58] Baso A. A. & Mudi S. Y. (2017). Evaluation of Antiulcer and Phytochemical Activities of Leaf Extract of *Tapinanthus dodoneifolius* DC. (Loranthaceae) Grown on *Tamarindus indica* Tree. *Bayero Journal of Pure and Applied Sciences*, 10 (1): 392-396.
- [59] Mohammed M., Idris A., Gandu I., Tanko U. M., Muhammad A. & A Adeiza A. (2019). Phytochemical and Antimicrobial Study on the Leaf Extract of *Tapinanthus dodoneifolius* (Van Tiegh) Loranthaceae. *Journal of Advances in Medicine and Medical Research*. **Volume 29, Issue 5, Page 1-9.**
- [60] Deeni Y. Y. & Sadiq N. M. (2002). Antimicrobial properties and phytochemical constituents of the leaves of African mistletoe (*Tapinanthus dodoneifolius* (DC) Danser) (Loranthaceae): an ethnomedicinal plant of Hausaland, Northern Nigeria. *J Ethnopharmacol*. 83 (3): 235-40.
- [61] Abdullahi Z., Anuka J. A., Salawu A. O., & Hussaini I. M. (2015). In-vivo antiplasmodial activity of methanol whole plant extracts of *Tapinanthus dodoneifolius* (DC) Danser in mice. *African Journal of Pharmacy and Pharmacology*, 9 (37), 936-942.
- [62] Ofem O. E., Eno A. E., Imoru J., Nkanu E., Uneh F., and Ibu J. O. (2007). "Effect of crude aqueous leaf extract of *Viscum album* (mistletoe) in hypertensive rats," *Indian Journal of Pharmacology*, vol. 39, no. 1, pp. 15–19.
- [63] Ouédraogo S., Aristide T. N., Soméa M. L., Pierre I. G., Christa S., Bernard B. and Ramarosan, A. (2005). Cardiovascular Properties of Aqueous Extract from *Tapinanthus Dodoneifolius* DC. Danser. *Afr. J. Traditional, Complementary and Alternative Medicines*, 2 (1): 25 – 30.
- [64] Harquin Simplice F., David Emery T. & Hervé Hervé N. A. (2014). Enhancing Spatial Memory: Anxiolytic and Antidepressant Effects of *Tapinanthus dodoneifolius* (DC) Danser in Mice. *Neurol Res Int*. 2014:974308.
- [65] Umarudeen A. M. & Magaji G. M. (2019). Comparative in-vivo anxiolytic efficacy of aqueous and methanol *Tapinanthus globiferus* leaf extracts. *Int Arch Med Health Res*.1(3): 89-93.
- [66] Umarudeen A. M. & Aminu C. (2020). Acute Toxicological and In-vivo Anxiolytic Activity Screening of Aqueous and Chloroform Fractions of Hydroalcoholic *Tapinanthus globiferus* Leaf Extracts. *World Journal of Innovative Research (WJIR)* ISSN: 2454-8236, Volume-8, Issue-
- [67] Akogu O. (2022). Study to Determine Toxicity of methanolic *Terminalia Catappa* Linn. (Almond Fruit) Leaf Extract on Adult Wistar Mice Using Dietrich Lorke Method. *Journal of Research in Agriculture and Animal Science*. Volume 9, Issue 3, pp: 12-14.
- [68] Iheagwam F. N., Okeke C. O., De Campos O. C., Adegboye B. E., Ogunlana O. O., & Chinedu S. N. (2021). Toxicopathological, proinflammatory and stress response evaluation of *Terminalia catappa* extract in male Wistar rats. *Toxicology reports*, 8 (18), 1769–1776.
- [69] Hall C. & Ballachey L. E. (1932). A study of the rat's behavior in a field. A contribution to method in comparative psychology. University of California Publications in Psychology 6: 1–12.
- [70] Yemitan, O. K., Akinsuyi, A. O., Jewo, P. I., Oguntola, J. A., & Olayemi, S. O. (2020). Toxicological and Reversibility Studies of *Bryophyllum pinnatum* Leaf Extract on Biochemical, Peroxidation and Histopathologic Parameters in Rodents. *Journal of Complementary Medicine Research*, 11(1), 95-95.
- [71] Ozolua, R.I., Idogun, S.E., & Tafamel, G.E. (2010). Acute and Sub-Acute Toxicological Assessment of Aqueous Leaf Extract of *Bryophyllum Pinnatum* (Lam.) in Sprague-Dawley Rats. *American Journal of Pharmacology and Toxicology*, 5 (3), 145-151.
- [72] Hall C. S. & Ballachey E. L. (1932). "A study of the rat's behavior in a field: a contribution to method in comparative psychology". University of California Publications in Psychology. 6 (1): 1–12.
- [73] Hall C. S. (1934). Drive and emotionality: factors associated with adjustment in the rat. *Journal of Comparative Psychology*, 17 (1), 89–108.
- [74] Seibenhener M. L. & Wooten M. C. (2015). Use of the Open Field Maze to measure locomotor and anxiety-like behavior in mice. *J Vis Exp*. (96): e52434.
- [75] Rex A., Voigt J. P., Voits M. & Fink H. (1998). Pharmacological evaluation of a modified open-field test sensitive to anxiolytic drugs. *Pharmacol Biochem Behav*. 59 (3): 677-83.
- [76] Lister R. G. (1990). Ethologically based animal models of anxiety disorders. *Pharmacol Ther*. 46 (3): 321-40.

- [77] Walsh R. N. & Cummins R. A. (1976). The open-field test: a critical review. *Psychol Bull.* 83 (3): 482-504.
- [78] Choleris E., Thomas A. W., Kavaliers M. & Prato F. S. (2001). A detailed ethological analysis of the mouse open field test: effects of diazepam, chlordiazepoxide, and an extremely low frequency pulsed magnetic field. *Neurosci Biobehav Rev.* 25 (3): 235-60.
- [79] Haque M. & Abubakar A. R. (2019). Anxiolytic Testing of Medicinal Plants in Nigeria: Frequently Used Experimental Models. *Medeniyet Med J.* 34 (1): 83-98.
- [80] Moniruzzaman M, Sharoti Bhattacharjee P, Rahman Pretty M, Sarwar Hossain M. (2016). Sedative and Anxiolytic-Like Actions of Ethanol Extract of Leaves of *Glinus oppositifolius* (Linn.) Aug. DC. *Evid Based Complement Alternat Med.* Volume 2016: 8541017.
- [81] Shahed-Al-Mahmud M., Lina S. M. M. (2017). Evaluation of sedative and anxiolytic activities of methanol extract of leaves of *Persicaria hydropiper* in mice. *Clin Phytosci* 3 (1), 20, 12 pages.
- [82] Foyet H. S., Tsala D. E., Bouba A. A. & Hritcu L. (2012). Anxiolytic and Antidepressant-Like Effects of the Aqueous Extract of *Alafia multiflora* Stem Barks in Rodents. *Adv Pharmacol Sci.* 2012:912041.
- [83] Barua C. C., Talukdar A., Begum S. A., Borah P. & Lahkar M. (2012). Anxiolytic activity of methanol leaf extract of *Achyranthes aspera* Linn in mice using experimental models of anxiety. *Indian J Pharmacol.* 44 (1): 63-7.
- [84] Habib M. R.; Billah M. M.; Mohammad Mahfuz-e-Alam; Kashfia Nawrin; Hasan, M. R.; Rahman M. M.; Islam, M. A.; Khan I. N. (2015). Evaluation of the analgesic, sedative-anxiolytic, cytotoxic and thrombolytic potentials of the different extracts of *Kalanchoe pinnata* leaves. *Journal of Coastal Life Medicine.* Vol.3 No.12 pp. 982-986.
- [85] Chandrasekhar Y., Ramya E. M., Navya K., Phani Kumar G., Anilakumar K. R. (2017). Antidepressant like effects of hydrolysable tannins of *Terminalia catappa* leaf extract via modulation of hippocampal plasticity and regulation of monoamine neurotransmitters subjected to chronic mild stress (CMS). *Biomed Pharmacother.* 86: 414-425.
- [86] Prut L, Belzung C. (2003). The open field as a paradigm to measure the effects of drugs on anxiety-like behaviors: a review. *Eur J Pharmacol.* 463 (1-3): 3-33.
- [87] Thompson T., Grabowski-Boase L. & Tarantino L. M. (2015). Prototypical anxiolytics do not reduce anxiety-like behavior in the open field in C57BL/6J mice. *Pharmacol Biochem Behav.* 133: 7-17.