

Assessment of Rice Yield Sensitivity to Changing Weather Conditions in Prayagraj Using DSSAT V4.7.5 Crop Simulation Model

ABSTRACT

Climate change will be one of the deciding factors for future food production, which needs sustainable intensification of agriculture. Field experiments were conducted during 2012-19 to determine the effect of changing weather such as (T_{max}, T_{min}, T_{avg}, Solar radiation & CO₂ concentration) on grain yield, LAI, Anthesis days and maturity days of four rice cultivars i.e. (Swarna sub 1, Sarjoo 52, Pant Dhan 4 and NDR 359) at the college of forestry farm, SHUATS Prayagraj. The DSSAT-CERES rice model was calibrated and validated for the cultivars in Prayagraj conditions and it was observed that the values i.e. Percent error, RMSE, nRMSE and Pearson correlation coefficient (r) were good in agreement and within permissible limits. Among all the four varieties NDR 359 yields more followed by Pant Dhan 4, Swarna sub 1 and Sarjoo-52. The results revealed that by increasing temperature (T_{max}, T_{min}, T_{avg}) for all the varieties and phenophases the yield got reduced but increased in Solar radiation and CO₂ concentration conditions the yield got increased. In case of LAI the same result was observed but during the phenophase of flowering to maturity stage of the crop there was no effect found. During the interaction between changing weather with anthesis days & maturity days it was found that the anthesis days and maturity days got increased with increased T_{max} and T_{avg}. Other weather parameters had no effect on it. The interaction of weather parameters with yield, LAI, anthesis days & maturity days were found significant at 5% or 1% level for all the four varieties & phenophases. The research outcome indicates that, the future farming will be challengeable due to climate change, we must prepare with suitable varieties and crop management plan to tackle the situation.

Keywords: DSSAT CERES-rice model, Climate change, Yield, Sensitivity analysis, Temperature, Solar Radiation, CO₂ concentration

Introduction

Rice is the most important food for half of the world's population especially in Asia (Schiller *et al.*, 2006). In India Rice occupies about 23.3% of gross cropped area of the country.

Among the rice growing countries in the world, India stands first in rice area (44 million ha) and second in rice production (112.91 million metric tons) after China. (FAO, 2017). It contributes 21.5 per cent of global rice production. Uttar Pradesh is the fourth largest state in country. The state has almost 5.86-million-hectare land under rice cultivation producing about 12.5 million tonnes of rice. The state ranks 3rd in the country in production of rice. In Prayagraj district, rice is produced in an area of 171824 hectare with a production of 370562 MT and productivity of 21.57 q/ha. (Dwivedi, 2018).

As climate change deals with future issues, the use of Crop Simulation Models proves a more scientific approach to study the impact of climate change on agricultural production. The decision support system for agrotechnology transfer (DSSAT) was originally developed by an international network of scientists, cooperating in the International Benchmark Sites Network for Agrotechnology Transfer project (IBSNAT, 1993; Tsuji, 1998; Uehara, 1998; Jones *et al.*, 1998), to facilitate the application of crop models in a systems approach to agronomic research. DSSAT models simulate growth, development, and yield of crops as a function of soil-plant-atmosphere-management dynamics. The CERES-Rice model of the DSSAT modeling system is an advanced physiologically based rice crop growth simulation model and has been widely applied to understanding the relationship between rice and its environment. (Ritchie *et al.*, 1987) and (Hoogenboom *et al.*, 2003) have provided a detailed description of the model. The model uses a detailed set of crop specific genetic coefficients, which allows the model to respond to diverse weather and management conditions. (Basak *et al.*, 2012).

Weather plays an important role in agricultural production. It has a profound influence on crop growth, development and yields. Weather factors contribute to optimal crop growth, development and yield. Climate is the primary determinant of agricultural productivity. (Adams *et al.*, 1998). Changes in temperature, CO₂, and precipitation under the scenarios of climate change for the next 30 year present a challenge to crop production. Understanding these implications for agricultural crops is critical for developing cropping systems resilient to stresses induced by climate change. (Hatfield *et al.*, 2011). The optimum temperature for rice cultivation is between 25°C and 35°C. Any further increase in mean temperatures during sensitive stages may reduce rice yields drastically. In tropical regions, the temperature increase due to the climate change is probably near or above the optimum temperature range for the physiological activities

of rice (Baker *et al.*, 1992). In temperate regions, rice growth is impressed by limited period that favours its growth (Reyes *et al.*, 2003). Increasing trend of daily maximum temperature may decrease the rice spikelet fertility, which affects for reduction of the yield while the increasing trend of atmospheric CO₂ concentration could increase the rice yield (Dharmarathna *et al.*, 2012). High temperatures would induce sterility and lead to low harvest index and grain yield.

Materials and Methods

Prayagraj is taken as the representative experimental site in Uttar Pradesh. The experiment was carried out at College of Forestry farm, Sam Higginbottom University of Agriculture, Technology and Sciences, Naini, Prayagraj. It is situated at 25.4358° N latitude, 81.8463° E longitude and at an altitude of 98 m (322 ft) above mean sea level.

Weather data:

For the study the daily weather parameters like Tmax, Tmin, Solar radiation and Rainfall were collected from Meteorological unit, Department of Environmental Sciences and NRM, College of Forestry, SHUATS, Naini, Prayagraj from 1st January 2012 – 1st January 2020.

Soil data:

Soil data Layer wise (0 – 120 cm) data of soil physical and chemical properties which includes Bulk density, Hydraulic conductivity, organic carbon content, clay and silt content etc. of Prayagraj district was collected from India Meteorological Department, New Delhi.

Crop management data:

Crop management data which is required for DSSAT input from 2012-2019 taken from College of Forestry, SHUATS, of four rice varieties. The data was used first calibration and then validation of the model. Planting date of the rice crop was 2 June for all the four cultivars.

Cultivars:

In this study there are four cultivars are used such as (Swarna sub 1, Sarjoo-52, Pant dhan 4 & NDR-359). The genetic coefficients of Cultivars were taken from India Meteorological Department, New Delhi, then again, they are calibrated for Prayagraj condition via trial-and-error method, which is presented in the Table below.

Table 1. Genetic coefficient of four varieties used in DSSAT

Cultivar	P1	P2R	P5	P2O	G1	G2	G3	G4
SWARNA SUB 1	750.0	150.0	400.0	11.3	59.0	0.0220	1.00	1.00
SARJOO - 52	450.0	170.0	365.0	12.2	47.0	0.0238	1.00	1.00
PANT DHAN 4	830.0	160.0	300.0	11.4	45.0	0.0300	1.00	1.00
NDR 359	500.0	200.0	450.0	12.5	62.0	0.019	1.00	1.00

For efficiently study the effect of weather parameters on rice crop, the whole duration of rice crop is divided in to four phases as listed below. By applying phase wise in environmental modification, the difference can be seen and analysed.

- Phase 1 : Germination – Transplanting
Phase 2 : Transplanting – Panicle initiation
Phase 3 : Panicle initiation – Flowering (Anthesis duration)
Phase 4 : Flowering – Maturity (Maturity duration)

Calibration & Validation of the model:

Model calibration is the adjustment of parameters, so that simulated values compare well with observed values. Before models can be applied with confidence, they need to be calibrated and validated for the varieties and environment of interest. Validation is determined by statistical analysis. It is used to check the accuracy of the model. Statistical based criteria provide a more objective method for evaluation of the performance of the models (Ducheyne, 2000). Here in this study only crop yield data is used for validation of the model. For the validation of the model Percent Error, RMSE, nRMSE and r was calculated.

Significance test:

The Pearson correlation is the most widely used correlation statistic and linear regression analysis is used to measure the degree of the relationship between linearly related variables which is widely used in climate research, will be employed in this study to find out significance level of 0.05 and 0.01 (indicates 5% and 1% risk respectively) trends with the help of IBM SPSS statistics package.

Sensitivity of field to changing climate:

During the growing season, the mean temperature, the maximum and minimum temperature, rainfall distribution pattern, and diurnal changes, or a combination of these, may be highly correlated with grain yields (Mooman and Vergara, 1965). In this study we will change the weather parameters in DSSAT like maximum, minimum and average temperature, solar radiation and CO₂ concentration to see the effect on yield and yield parameters. (Hardacre and Turnbull, 1986) state that temperature affects the duration of crop growth and consequently the time during which incident radiation can be intercepted and transformed to dry matter. Effects of increase in temperature, solar radiation and CO₂ concentration on rice yield and other parameters assessed by increasing the maximum, minimum & average temperature by +1, +2, +3, +4, +5 °c , solar radiation by +1.0, +1.5, +2.0, +2.5, +3.0 MJ/day and CO₂ concentration by +20, +40, +60, +80, +100 ppm respectively.

Results:

Calibration and validation of model

Calibration of model is done for 2 years 2012 and 2013 and model is validated from 2014 to 2019 for the cultivar to check the accuracy of model in Prayagraj condition. The Actual and Simulated yield data from the year of 2012 – 2019 for all the four variety are given below in the table 2. The percent error, root mean square error, normalized root mean square error and Pearson correlation coefficient for the year (2012-2019) shows that simulated yield values were in good agreement with the observed yield values. Average percent error, RMSE, nRMSE and r value are 3.29, 183.78, 3.62, 0.828 and 9.08, 431.24, 9.75, 0.542 respectively, for Swarna sub 1 & Sarjoo-52 variety. For Pant dhan 4 & NDR 359 rice cultivar average percent error, RMSE, nRMSE and r value are 4.29, 238.14, 4.61, 0.778 and 5.52, 338.96, 5.84, 0.71 respectively, which shows that model is suited well for these variety in Prayagraj condition.

Impact of changing weather on various growth phases

Sensitivity of rice yield to changing weather

The average yield data of 8 years for four rice crop varieties with changing maximum temperature, average temperature, minimum temperature, solar radiation & CO₂ conc. for four phenophases under Prayagraj condition presented in Table 3. The table shows decreasing trend of yield with increasing in temperature for all varieties. In case of increase in solar radiation and

CO₂ conc. the table shows increasing trend of yield for all varieties & phenophases. Maximum yield reduced during germination – transplanting stage followed by transplanting to panicle initiation stage for all the four variety. Among all the cultivars NDR-359 yield was reduced to -29.44% followed by Swarna sub 1 (-19.41%), Pant dhan 4 (-14.95%) and Sarjoo-52 (-13.44%). When minimum temperature increased, the rice yield was decreased to -4.52% in case of Pant dhan 4 followed by Sarjoo 52 (-3.96%), NDR 359 (-3.15%) & Swarna sub 1 (-1.45%). In case of increased in average temperature the yield of NDR 359 cultivar got reduced to -41.38% followed by Sarjoo-52 (-26.55%), Pant dhan 4 (-19.49%) and Swarn sub 1 (-14.37%). Increased in Solar radiation, the crop yield got increased. Among all cultivar Sarjoo-52 it was (15.39%), Pant dhan 4 (11.12%), Swarna sub 1 (11.09%) & NDR 359 (3.05%). Increased in CO₂ concentration, the crop yield of Sarjoo 52 (8.49%), NDR 359 (7.46%), Pant dhan 4 (7.26%) & Swarna sub 1 (7.19%) got increased.

Sensitivity of Leaf area index (LAI) to changing weather

For all the four phases of crop, the Leaf area index of four rice crop varieties with changing maximum temperature, average temperature, minimum temperature, solar radiation & CO₂ under Prayagraj condition presented in Table 4. The table shows decreasing trend of LAI with increasing trend of maximum, average & minimum temperature for all varieties. There was increasing trend of LAI with increasing in Solar radiation & CO₂ for all varieties. For the last phase of crop growth (Flowering - Maturity), there is no increase / decrease of Leaf area index as we change all the weather parameters (T max, T avg, T min, SRAD & CO₂ conc.). Maximum LAI reduced during germination – transplanting stage for all the four variety. Among all the cultivars NDR-359, LAI was reduced to -29.88% followed by Sarjoo-52 (-18.13%), Pant dhan 4 (-14.07%) and Swarna sub 1 (-13.75%). When minimum temperature increased, the rice crop LAI was also decreased i.e., -7.02% in case of Swarna sub 1 followed by Sarjoo 52 (-6.29%), NDR 359 (-2.62%) & Pant dhan 4 (-2.38%). In case of increased in average temperature the LAI of NDR 359 got reduced to -52.49% followed by Sarjoo-52 (-46.07%), Pant dhan 4 (-42.11%) and Swarn sub 1 (-41.15%). Increased in Solar radiation, the LAI got increased. Among all cultivar Sarjoo-52 it was (10.04%), Swarna sub 1 (3.86%), NDR 359 (2.98%) & Pant dhan 4 (2.20%). Increased in CO₂ concentration, the LAI of Sarjoo 52 (4.81%), Swarna sub 1 (1.88%), Pant dhan 4 (1.56%) & NDR 359 (1.55%) got increased.

Sensitivity of Anthesis days to changing weather

Table 5. shows the variation in anthesis days due to effect of increase in all the weather parameters (T max, T avg, T min, SRAD & CO₂ conc.) on four rice varieties (Swarna sub-1, Sarjoo-52, Pant dhan-4, NDR 359). All the varieties are showing increase in Anthesis days as increase in maximum & average temperature during Germination – flowering stage of crop. For the period of Flowering to Maturity, there is no increase / decrease of Anthesis days as we change all the weather parameters (T max, T min, SRAD & CO₂ conc.). As it is the last stage there is no more effect of weather parameters on anthesis duration. There is no effect of (T min, SRAD & CO₂conc.) on anthesis days. In case of increased in maximum temperature the anthesis days of NDR 359 got increased to (45.45%) followed by Swarn sub 1 (19.75%), Pant dhan 4 (16.53%) and Sarjoo-52 (15.88%). Increased in average temperature, the anthesis days also got increased. Among all the cultivar, in case of NDR 359 it was (45.45%), Swarna sub 1 (29.49%), Pant dhan 4 (26.09%) & Sarjoo-52 (18.66%).

Sensitivity of Maturity days to changing weather

The Maturity days of four rice crop varieties with changing maximum temperature & average temperature from (1.0 -5.0°C) under Prayagraj condition presented in Table 6. The table shows increasing trend of Maturity days with increasing in T max & T avg for all varieties and phenophases. There is no effect of changing (T min, SRAD & CO₂conc.) on maturity days. Among all the cultivars maturity days of NDR-359 was increased to 35.54% followed by Pant dhan 4 (14.67%), Swarna sub 1 (13.87%) & Sarjoo-52 (13.55%). In case of average temperature increased, the rice maturity days was also increased i.e 39.87% for NDR 359 followed by Swarna sub 1 (20.12%), Sarjoo-52 (19.08%) & Pant dhan 4 (18.29%).

Table 2. Comparison of cultivars observed value with simulated value of grain yield for the year 2012 – 2019.

Year	Swarna Sub 1			Sarjoo-52			Pant Dhan-4			NDR- 359		
	Actual Yield (Kg/ha)	Simulated Yield (Kg/ha)	Percent Error %	Actual Yield (Kg/ha)	Simulated Yield (Kg/ha)	Percent Error %	Actual Yield (Kg/ha)	Simulated Yield (Kg/ha)	Percent Error %	Actual Yield (Kg/ha)	Simulated Yield (Kg/ha)	Percent Error %
2012	5118	5023	1.85	4796	4688	2.25	5349	5060	5.40	5725	5558	2.91
2013	5371	5209	3.01	4635	5111	10.26	5283	5454	3.23	6238	6530	4.68
2014	5246	5552	5.83	4581	5196	13.42	5472	5636	2.99	6021	6545	8.70
2015	5460	5236	4.10	4093	4644	13.46	5016	5393	7.51	6107	5726	6.23
2016	5127	5310	3.56	4412	4939	11.94	5164	5395	4.47	5514	5847	6.03
2017	4813	4930	2.43	4257	4649	9.20	4837	4902	1.34	5397	5110	5.31
2018	4941	5139	4.00	4329	4584	5.89	5320	5505	3.47	6202	5815	6.23
2019	4432	4501	1.55	4264	4531	6.26	4798	4515	5.89	5156	5364	4.03
Average	5063.5	5112.5	3.29	4420.87	4792.75	9.08	5154.87	5232.5	4.29	5795	5811.87	5.52
RMSE	183.78			431.24			238.14			338.96		
nRMSE	3.62			9.75			4.61			5.84		
r	0.828			0.542			0.778			0.71		

NOTE: RMSE=(Root Mean Square Error), nRMSE =(Normalised Root Mean Square Error) and r =(Pearson correlation coefficient)

Table 3. Effect of changing weather parameters on yield (percentage basis) of four rice cultivars for four phenophases.

	Germination – Transplanting				Transplanting – Panicle initiation				Panicle initiation – Flowering				Flowering – Maturity				
		Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359
CTRL	kg/ha	5112.5	4792.75	5232.5	5811.87	5112.5	4792.75	5232.5	5811.87	5112.5	4792.75	5232.5	5811.87	5112.5	4792.75	5232.5	5811.87
Tmax	+1	-0.23	0.04	-0.38	-5.51	-0.40	0.05	-0.40	-5.69	0.86	0.79	0.91	0.43	0.02	-0.07	-0.48	-1.19
	+2	-1.57	1.65	0.15	-7.85	-0.63	0.96	1.38	-10.01	-0.02	-0.07	0.27	-0.18	-0.53	-0.71	-0.73	-2.55
	+3	-4.91	-1.32	-1.26	-7.03	-4.00	-1.08	-3.24	-11.32	-0.13	-0.65	-0.28	-0.50	-1.04	-1.17	-1.39	-3.00
	+4	-11.98	-6.36	-6.84	-16.43	-8.04	-4.37	-6.63	-14.24	-1.41	-1.14	-1.02	-0.60	-1.33	-1.94	-1.03	-4.51
	+5	-19.41	-13.44	-14.95	-29.44	-11.30	-8.40	-10.65	-23.34	-2.94	-1.38	-1.80	-1.17	-2.42	-2.67	-2.04	-5.63
Tmin	+1	-0.77	-2.22	-2.36	-1.31	-0.52	-2.19	-2.30	-1.38	-0.22	-0.56	-1.08	-1.77	-0.11	-0.07	-0.62	-0.36
	+2	-1.45	-3.95	-4.52	-2.81	-1.07	-3.96	-4.31	-3.15	-0.51	-1.17	-1.87	-1.90	-0.13	-0.19	-0.99	-0.43
	+3	-	-	-	-	-	-	-	-	-0.97	-1.50	-2.39	-3.05	-0.20	-0.23	-1.44	-0.98
Tavg	+1	-2.07	-2.71	-1.38	-6.64	-1.87	-1.73	-1.48	-6.30	-0.87	-0.51	-0.15	-2.66	0.05	-0.82	-0.04	-0.97
	+2	-4.67	-3.51	-2.37	-6.73	-3.64	-1.85	-0.80	-12.27	-1.65	-1.26	-0.30	-5.85	-1.19	-1.26	-0.92	-2.05
	+3	-11.00	-8.09	-7.38	-16.59	-9.49	-3.68	-5.77	-11.87	-1.86	-1.82	-0.80	-7.10	-1.95	-2.38	-2.74	-3.60
	+4	-4.34	-20.60	-12.40	-27.80	-4.94	-12.84	-11.91	-21.90	-4.90	-2.28	-2.12	-8.50	-3.40	-3.81	-3.51	-3.91
	+5	-14.37	-26.55	-18.17	-41.38	-13.79	-24.63	-19.49	-35.32	-8.03	-2.94	-7.36	-10.07	-5.41	-5.81	-5.16	-4.93
SRAD	+1.0	3.33	5.80	3.46	1.83	3.48	5.64	3.86	1.89	1.68	2.39	2.57	0.20	0.43	0.04	0.24	-0.86
	+1.5	5.68	9.19	5.93	1.51	5.15	8.24	5.96	2.77	3.08	3.69	3.84	0.22	0.66	0.24	0.38	-1.21
	+2.0	7.25	10.86	7.57	1.48	7.10	10.92	8.24	3.30	4.19	5.05	5.09	0.32	0.92	0.44	0.53	-1.41
	+2.5	8.72	12.92	9.86	3.33	8.35	12.88	9.92	4.17	5.58	5.84	5.98	0.37	1.20	0.47	0.70	-1.78
	+3.0	11.09	15.39	11.12	3.05	9.50	14.57	10.63	3.36	6.56	7.14	7.24	1.12	1.44	0.62	0.85	-2.00
CO2	+20	1.31	1.75	1.28	1.43	1.63	1.63	1.26	1.35	0.78	0.85	0.85	0.77	0.04	0.04	0.07	0.06
	+40	2.62	3.42	2.74	3.45	2.94	3.23	2.68	2.76	1.55	1.73	1.70	1.49	0.09	0.09	0.15	0.11
	+60	3.98	5.20	4.14	4.76	4.23	4.90	4.06	4.09	2.29	1.87	2.55	1.44	0.13	0.14	0.22	0.16
	+80	5.49	6.71	5.67	5.68	5.71	6.61	5.78	5.07	3.16	3.51	3.48	3.13	0.18	0.19	0.30	0.23
	+100	6.71	8.49	7.18	6.99	7.19	8.24	7.26	7.46	4.03	4.46	2.25	4.05	0.23	0.14	0.38	0.28

NOTE: Unit - SRAD (Solar radiation) in MJ/day, Maximum (Tmax), minimum (Tmin) & average temperature (Tavg) in °C and CO₂ concentration in ppm.

Table 4. Matrix of changing weather parameters on LAI (percentage basis) of four rice cultivars for the phenophases.

	Germination – Transplanting				Transplanting – Panicle initiation				Panicle initiation – Flowering				
		Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR- 359	Swarna Sub 1	Sarjoo- 52	Pant Dhan-4	NDR- 359	Swarna Sub 1	Sarjoo -52	Pant Dhan-4	NDR- 359
CTRL		5.06	4.06	5.83	6.79	5.06	4.06	5.84	6.79	5.06	4.06	5.84	6.79
Tmax	+1	-1.58	-1.36	-1.60	-2.40	-1.58	-1.68	-2.38	-1.86	-1.29	-0.44	-1.52	-1.08
	+2	0.40	-2.59	-3.31	-5.35	-0.85	-1.36	-3.23	-5.91	-2.04	-0.35	-3.13	-1.96
	+3	-2.47	-6.04	-4.60	-13.27	-1.84	-5.38	-5.60	-11.24	-3.15	-0.44	-4.52	-2.70
	+4	-5.84	-11.22	-8.06	-20.89	-4.55	-8.14	-6.02	-18.61	-5.04	-1.23	-5.37	-3.19
	+5	-13.75	-18.13	-14.07	-29.88	-12.23	-14.62	-6.88	-25.06	-6.63	-3.16	-6.57	-3.75
Tmin	+1	-0.69	-1.36	-0.51	-1.00	-1.09	-1.11	-0.66	-1.00	-0.69	-0.62	-0.46	-0.31
	+2	-1.48	-2.10	-1.03	-1.74	-1.48	-2.84	-1.35	-1.44	-3.07	-1.11	-1.04	-0.52
	+3	-3.07	-5.06	-2.23	-2.62	-3.07	-6.29	-2.38	-2.33	-7.02	-2.34	-2.03	-0.71
Tavg	+1	-2.57	-3.53	-2.66	-2.40	-3.32	-4.14	-2.80	-2.96	-2.45	-1.16	-2.03	-1.56

	+2	-5.30	-7.84	-6.30	-10.14	-4.55	-5.67	-6.23	-8.85	-3.34	-2.02	-4.26	-2.33
	+3	-9.75	-16.47	-11.23	-22.29	-9.50	-12.77	-11.16	-19.71	-5.88	-3.30	-6.57	-3.31
	+4	-22.12	-29.10	-23.89	-36.28	-20.38	-22.32	-20.36	-30.58	-9.08	-6.14	-8.40	-4.02
	+5	-41.15	-46.07	-42.11	-52.49	-36.46	-38.35	-36.85	-44.61	-12.07	-10.46	-10.90	-4.66
SRAD	+1.0	0.89	3.87	0.17	0.18	0.40	3.58	-0.49	-0.38	0.91	1.53	1.02	0.80
	+1.5	2.37	6.34	0.34	-0.56	1.13	5.43	-0.83	-1.49	1.29	2.02	1.44	1.44
	+2.0	2.67	7.89	0.51	-1.15	1.13	7.89	-1.69	-1.86	1.54	2.86	1.77	0.69
	+2.5	3.11	9.12	0.51	-2.47	1.38	8.20	-2.20	-2.03	2.27	3.45	1.91	2.61
	+3.0	3.86	10.04	0.77	-2.33	2.37	9.12	-2.55	-2.77	2.81	4.07	2.20	2.98
CO2	+20	0.10	1.11	0.34	-1.00	0.30	0.79	0.02	-0.38	0.45	0.57	0.23	0.18
	+40	0.30	2.34	0.51	-1.30	0.69	2.34	0.54	-0.38	0.71	1.13	0.62	0.35
	+60	0.49	3.08	0.56	-1.59	0.69	2.96	0.36	-1.12	0.91	1.41	1.05	0.47
	+80	0.69	3.82	0.77	-1.74	1.29	3.58	0.71	-1.30	1.11	2.00	1.39	1.18
	+100	1.09	4.81	1.03	-2.03	1.88	3.87	1.22	-2.40	1.40	2.64	1.56	1.55

NOTE: Unit - SRAD (Solar radiation) in MJ/day, Maximum (Tmax), minimum (Tmin) & average temperature (Tavg) in °C and CO₂ concentration in ppm.

Table 5. Model output of maximum & average temperature on Anthesis days (percentage basis) of four rice cultivars for the phenophases.

	Germination – Transplanting				Transplanting – Panicle initiation				Panicle initiation – Flowering				
		Swarna Sub 1	Sarjoo- 52	Pant Dhan-4	NDR- 359	Swarna Sub 1	Sarjoo- 52	Pant Dhan-4	NDR- 359	Swarna Sub 1	Sarjoo- 52	Pant Dhan-4	NDR- 359
CTRL		98.75	89.75	105.87	124.87	98.75	89.75	105.87	124.87	98.75	89.75	105.87	124.87
Tmax	+1	1.39	2.09	1.65	1.70	1.27	1.39	1.18	1.30	0.38	0.70	0.47	0.20
	+2	4.94	5.43	4.60	6.41	4.18	3.90	4.01	5.71	1.14	2.09	1.42	1.10
	+3	8.35	8.50	8.03	9.41	7.09	7.10	7.20	8.01	2.66	3.06	2.60	15.32
	+4	12.41	12.26	11.81	11.91	10.63	10.45	5.55	9.71	3.42	4.46	4.60	37.34
	+5	19.75	15.88	16.53	14.01	14.30	13.37	6.02	11.71	4.81	5.57	7.44	45.45
Tavg	+1	1.90	1.81	1.65	1.50	1.27	1.39	1.18	1.20	0.25	0.70	0.47	0.20
	+2	5.70	5.57	4.60	6.21	4.81	4.74	4.49	4.60	1.39	2.23	1.42	1.10
	+3	10.51	10.58	8.03	10.21	8.73	9.05	8.50	9.41	3.04	3.62	2.60	15.32
	+4	17.34	15.88	11.81	11.90	14.18	13.79	14.99	13.91	4.68	5.71	4.60	37.34

	+5	21.77	22.28	16.53	13.41	29.49	18.66	26.09	17.92	7.34	8.77	6.97	45.45
--	----	-------	-------	-------	-------	-------	-------	-------	-------	------	------	------	-------

Table 6. Model output of maximum & average temperature on Maturity days (percentage basis) of four rice cultivars for the phenophases.

	Germination – Transplanting				Transplanting – Panicle initiation				Panicle initiation – Flowering				Flowering – Maturity				
		Swarna Sub 1	Sarjoo- 52	Pant Dhan- 4	NDR- 359	Swarna Sub 1	Sarjoo- 52	Pant Dhan- 4	NDR- 359	Swarna Sub 1	Sarjoo- 52	Pant Dhan- 4	NDR- 359	Swarna Sub 1	Sarjoo- 52	Pant Dhan- 4	NDR- 359
CTRL		128	124.5	131.25	152.37	128	124.5	131.25	152.37	128	124.5	131.25	152.37	128	124.5	131.25	152.37
Tmax	+1	1.66	1.81	1.90	1.15	1.37	1.41	1.52	0.90	0.68	1.00	0.95	0.33	0.29	0.50	0.38	0.06
	+2	4.30	4.72	4.67	5.41	3.52	3.84	4.38	3.88	1.86	2.71	2.19	0.90	0.67	1.00	0.95	0.23
	+3	7.32	7.53	7.90	7.30	6.15	6.53	6.67	9.02	3.13	4.52	3.94	1.97	0.98	1.71	1.24	0.49
	+4	10.45	10.54	10.69	9.43	8.98	9.34	6.95	10.66	4.39	5.84	4.10	33.01	1.46	2.21	1.71	0.66
	+5	13.87	13.55	14.67	10.99	13.28	12.35	7.14	11.80	5.86	7.97	6.19	34.54	1.76	2.91	1.90	0.82
Tavg	+1	1.27	1.51	1.81	0.90	0.78	1.20	1.43	0.49	0.39	0.80	0.76	0.33	0.29	0.10	0.57	0.13
	+2	4.40	4.72	4.67	4.68	3.71	4.22	4.63	3.97	1.76	2.71	2.19	1.48	0.59	1.00	1.05	0.22
	+3	8.59	8.84	7.82	7.43	7.32	7.83	7.71	6.56	3.81	4.92	3.90	12.31	1.17	1.91	1.33	0.32
	+4	14.45	13.45	10.97	8.29	13.38	12.15	12.67	8.45	5.76	6.12	5.90	32.32	1.86	3.11	1.90	0.34
	+5	20.12	19.08	14.86	11.38	19.12	20.08	18.29	11.14	8.40	11.13	8.00	39.87	2.71	4.42	2.38	0.36

NOTE : Unit - SRAD (Solar radiation) in MJ/day, Maximum (Tmax), minimum (Tmin) & average temperature (Tavg) in °c and CO₂ concentration in ppm.

Discussion:

This section describes about the overall analysis of four rice cultivars (Swarna sub-1, Sarjoo-52, Pant dhan 4 & NDR 359) with the phenophases under Prayagraj condition. Calibration and validation of DSSAT CERES – rice model for grain yield was done for all four cultivars. By comparing observed grain yield with simulated yield, the result concluded that all the values were in acceptable range of less than 15% for percent error, in excellent range of less than 10% for normalized root mean square error and Pearson correlation coefficient value is also nearer to 1. Thus, the DSSAT rice crop model can be successfully used for simulating growth and yield of all the four rice varieties for Prayagraj condition. Among all the four varieties NDR 359 yields more followed by pant dhan 4, Swarna sub 1 and sarjoo-52 for Prayagraj condition.

Weather condition affected crop production both directly and indirectly. Temperature is considered to be one of the dominant factors that affect the growth and yield of rice. The impact of climate change on high yield rice varieties was studied by (Karim *et al.*, 1994) using the CERES rice model and several scenarios and sensitivity analysis. They found that high temperatures reduced rice yields in all seasons in most arid locations. Increased CO₂ levels increased rice yields and reduction of rice yields due to high temperature in all season was observed by (Karim *et al.*, 1994). This study showed that increasing temperature results with decreasing yield. To know the effect, Whole crop growth period divided into four stages. Each stage showed that decreasing yield with increasing temperature. For increasing maximum temperature results with decreasing yield. But when we talk about the crop growth period, panicle initiation – flowering (anthesis) stage yields more among all phenophases followed by flowering – maturity, transplanting – panicle initiation and germination – transplanting for four varieties. Hatfield & Pruegar (2015) found that warm temperatures increased the rate of senescence during grain fill and reduced final grain yield. As average temperature increases the yield decreases. For the varieties of Sarjoo-52 and Pant dhan 4 the yield is more during panicle initiation – flowering (Anthesis) period. The yield is more during Maturity period for the varieties Swarna sub 1 and NDR 359. Increasing minimum temperature results with decreasing yield. When we talk about the crop growth period, with the effect of minimum temperature the yield is more in Maturity period followed by panicle initiation – flowering, transplanting- panicle initiation and germination to transplanting Stage for all the varieties. The results of solar radiation and CO₂ concentration is opposite as compared with

temperature. As the solar radiation increased simultaneously the yield of all four varieties also increased. For the variety of Pant dhan 4 and NDR 359 Transplanting – panicle initiation stage yields more and for Swarna sub 1 and sarjoo-52 germination – transplanting stage yields more. Hendrey & Kimball (1994) reported that higher CO₂ concentration increases growth and yield, mainly through their effect on the photosynthetic processes of the crop. Increased CO₂ concentrations and decreased temperature increased growth duration and yield, while increased temperature shortened growth duration and reduced yield. Increasing the CO₂ concentration by 100 ppm and 400 ppm from the standard CO₂ concentration of 380 ppm led to 16.8% and 54.2% increase in grain yield, respectively. (Nyang'au *et al.*, 2014). According to Hunsaker *et al.* (2000) high carbon dioxide concentrations increase water use efficiency which has a major effect on yield. As per our study given in the table, yield increases with increasing CO₂ concentration. Germination – transplanting stage yields more followed by transplanting – panicle initiation, anthesis duration than maturity period for all the four varieties. When we analyse the yield data with changing weather for stage wise growth of crop with SPSS software for significance test, the result presented in the Table 7. showed all the data are significant.

Leaf area index is the projected area of leaves over a unit of land. Changing weather directly or indirectly effect the leaf area index which ultimately effect yield. As we divided the crop period into four phases the flowering – maturity period, has no changes in leaf area index with changing temperature, solar radiation and CO₂ concentration etc. Leaf area index decreased with increase of maximum temperature. When we talk about the growth stages maximum leaf area index found in panicle initiation – flowering duration followed by transplanting – panicle initiation then germination – transplanting for all four varieties. Among all the varieties NDR-359 has more leaf area index followed by pant dhan 4, Swarna sub 1 and Sarjoo-52. For increase in average temperature the leaf area index goes on decreasing. Among the crop periods Panicle initiation – flowering stage has more leaf area index compare to all other stages. As Minimum temperature increases leaf area index decreases. For the variety of Swarna sub 1 leaf area index is more during Germination – transplanting duration and for other three varieties during the period of Panicle initiation – flowering the leaf area index is more. Increasing solar radiation results increase in leaf area index. When we talk about the effect of solar radiation on crop growth period during the period of germination – transplanting Swarna sub 1 and Sarjoo-52 variety have more leaf area index. Pant dhan 4 and NDR 359 variety showed high leaf area index during panicle initiation –

flowering stage. Van Keulen (1984) using a simulation model predicted that an increase of 20% in total global radiation resulted in 10–20% increase in grain yield of rice. In a simulation study on the effect of solar radiation on growth of wheat and rice, it was revealed that the maximum Leaf Area Index was reduced by 7.6% in wheat and 5.9% in rice when the solar radiation was decreased by 10.0% from normal. On the other hand, with increase in radiation by 10%, LAI increased in wheat by 7.1%. Hundal and Kaur (1995). As per the analysis it showed that leaf area index value increases as CO₂ concentration increases. Pant dhan 4 and NDR 359 variety has more leaf area index during panicle initiation – flowering stage and Sarjoo-52 has more leaf area index during germination – transplanting period. By Analyzing the data of Leaf area index with changing weather for stage wise growth of crop using SPSS software for significance test, the results are presented in the Table 8. It showed all the data were significant and r value is nearly equal to 1. The stage of flowering – maturity has no changes in the value of leaf area index with changing weather.

Anthesis days are very crucial period for crop growth. Changing temperature has effect on anthesis days but there is no effect of solar radiation and CO₂ concentration on this. During the period of flowering – maturity there was not seen any changes regarding anthesis days. As the maximum temperature increases the anthesis day time period also increases. On the view of crop growth period there were high anthesis days seen in germination – transplanting followed by transplanting – panicle initiation and panicle initiation – flowering. When we talk about the average temperature anthesis days increases simultaneously with temperature. For the variety of Swarna sub 1 and Pant dhan 4 high anthesis days are found during Transplanting – Panicle initiation, for NDR 359 panicle initiation – Flowering and during Germination – Transplanting phase Sarjoo-52 variety found high anthesis days value. The data of Anthesis days with changing weather were analysed for stage wise growth of crop with SPSS software for significance test, the result are presented in the Table 9. It showed all the data were significant and r value is nearly equal to 1. The stage of flowering – maturity has no changes in the value of Anthesis days with changing weather. Except temperature all other weather parameters has no effect on anthesis days of a crop.

Maturity days comes under last stage in which crop gets mature. There is no effect of solar radiation and CO₂ concentration on maturity days of a crop. Only factor is as the temperature increases maturity days also get increases. When we talk about the crop stages, the varieties like

Swarna sub 1, Sarjoo-52 and Pant dhan 4 showed high maturity days during germination – transplanting. For the period of Panicle initiation – flowering NDR 359 variety has high maturity days. Likewise, as average temperature increases it results in increasing maturity days. The variety Swarna sub 1 and Sarjoo-52 showed high maturity days value during germination – transplanting. For the duration of Panicle initiation – flowering and Transplanting – Panicle initiation NDR 359 & Pant dhan 4 has high maturity value respectively. The data of Maturity days with changing weather were analysed for stage wise growth of crop with SPSS software for significance test, the results are presented in the Table 10. It showed all the data were significant and r value is nearly equal to 1. Except temperature (maximum and average) all other weather parameters has no effect on maturity days of a crop.

CONCLUSION:

The impact of climate change on rice yield, LAI, anthesis & maturity days for the four rice cultivars and for four phenophases has been analyzed by using the DSSAT crop simulation model. The study undertaken could be concluded with the facts that weather changes affect the rice varieties yield and its attributes. Elevated temperature beyond the threshold adversely affects the growth and development of rice crops, which ultimately determines yield. As the temperature increased the yield and LAI decreased and as solar radiation & CO₂ concentration increased the yield and LAI increased. When we discuss about the anthesis and maturity days, as temperature increased the anthesis & maturity days also get increased. There is no effect of solar radiation, CO₂ concentration and minimum temperature on anthesis and maturity days for four rice cultivars as taken. Model calibration and validation showed that all the four varieties (Swarna sub-1, Sarjoo-52, Pant dhan 4, NDR-359) were suitable for Prayagraj condition & NDR-359 was high yielding variety among all the four. The findings will help rice breeders to develop thermo-tolerant rice cultivars to cope with future climate change and obtain high yields for fulfill the demand of the rice-based community.

Table 7. Relationship between weather variables and grain yields for all the phenological stages at Prayagraj.

Weather variables	Correlation coefficient (r)															
	Germination – Transplanting				Transplanting – Panicle initiation				Panicle initiation – Flowering				Flowering – Maturity			
	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359
Tmax	-0.928**	-0.838*	-0.847*	-0.916*	0.944**	-0.854*	-0.888*	0.971**	-0.853*	-0.879*	-0.850*	0.899**	0.959**	0.981**	0.935**	0.995**
Tavg	-0.999*	-0.997*	-1.00*	-0.999*	-1.00*	-0.998*	-0.999*	-0.997*	-0.986*	0.993**	-0.988*	-0.952*	-0.968*	-0.982*	0.994**	-0.960*
Tmin	-0.827*	0.944**	0.963**	0.962**	-0.875*	-0.884*	0.925**	0.956**	0.929**	0.998**	-0.814*	0.983**	0.963**	0.972**	0.973**	0.990**
SRAD	0.998**	0.994**	0.998**	0.898*	0.997**	0.996**	0.993**	0.923**	0.996**	0.999**	0.999**	0.828*	0.999**	0.962**	0.998**	0.992**
CO ₂	1.00**	1.00**	1.00**	0.993**	0.999**	1.00**	0.999**	0.993**	1.00**	0.982**	0.858*	0.968**	1.00**	0.918**	1.00**	0.999**

(* = 5% level of significance & ** = 1% level of significance)

Table 8. Relationship between weather variables and leaf area index for the phenological stages at Prayagraj.

Weather variables	Correlation coefficient (r)											
	Germination – Transplanting				Transplanting – Panicle initiation				Panicle initiation – Flowering			
	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359
Tmax	-0.844*	-0.949**	-0.952**	-0.975**	-0.8348	-0.936**	-0.979**	-0.982**	-0.991**	-0.837*	-0.995**	-0.989**
Tavg	-0.973*	-0.957*	-0.974*	-0.999**	-0.973*	-0.964*	-0.988*	-0.992**	-0.951*	-0.969*	-0.984*	-0.983*
Tmin	-0.919**	-0.960**	-0.928**	-0.974**	-0.925**	-0.946**	0.940**	-0.977**	-0.989**	-0.947**	-0.999**	-0.990**
SRAD	0.986**	0.989**	0.985**	-0.888*	0.952**	0.981**	-0.969**	-0.964**	0.992**	0.998**	0.974**	0.894*
CO ₂	0.990**	0.994**	0.962**	-0.952**	0.975**	0.970**	0.946**	-0.936**	0.984**	0.993**	0.993**	0.969**

(* = 5% level of significance & ** = 1% level of significance)

Table 9. Relationship between weather variables and anthesis days for the phonological stages at Prayagraj.

Weather variables	Correlation coefficient (r)											
	Germination – Transplanting ANTHESIS				Transplanting – Panicle initiation				Panicle initiation – Flowering			
	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359
Tmax	0.976**	0.997**	0.989**	0.992**	0.991**	0.993**	0.870*	0.988**	0.978**	0.994**	0.980**	0.843*
Tavg	0.987**	0.986**	0.989**	0.982**	0.926**	0.987**	0.949**	0.988**	0.969**	0.977**	0.968**	0.932**
Tmin	-	-	-	-	-	-	-	-	-	-	-	-
SRAD	-	-	-	-	-	-	-	-	-	-	-	-
CO₂	-	-	-	-	-	-	-	-	-	-	-	-

(* = 5% level of significance & ** = 1% level of significance)

Table 10. Relationship between weather variables and maturity days for the phonological stages at Prayagraj.

Weather variables	Correlation coefficient (r)															
	Germination – Transplanting				Transplanting – Panicle initiation				Panicle initiation – Flowering				Flowering – Maturity			
	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359	Swarna Sub 1	Sarjoo-52	Pant Dhan-4	NDR-359
Tmax	0.989**	0.994**	0.988**	0.991**	0.973**	0.989**	0.962**	0.981**	0.989**	0.991**	0.980**	0.843*	0.996**	0.994**	0.997**	0.988**
Tavg	0.978**	0.984**	0.995**	0.987**	0.970**	0.966**	0.980**	0.990**	0.978**	0.965**	0.989**	0.930**	0.978**	0.977**	0.997**	0.957**
Tmin	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SRAD	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
CO₂	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

(* = 5% level of significance & ** = 1% level of significance)

References:

1. Adams RM, Hurd BH, Lenhart S, Leary N. Effects of Global Climate Change on Agriculture: An Interpretative View, *Climate Research* 1998; 11:19-30.
2. Baker JT, Allen LH, Boote KJ. Temperature effects on rice at elevated CO₂ concentration. *J Exp Bpt.* 1992; 43:959-964.
3. Basak JK, Ali MA, Biswas JK, Islam MN. Assessment of the effect of climate change on Boro rice yield and yield gaps using DSSAT model. *Bangladesh Rice Journal.* 2012; 16, 67-75.
4. Dharmarathna WRSS, Weerakoon SB, Rathnayaka UR, Herath S. Variation of Irrigated rice yield under the climate change scenarios, SAITM research symposium on Engineering Advancements. 2012; Pp. 31-34.
5. Ducheyne S. Derivation of the parameters of the WAVE model using a deterministic and a stochastic approach. Ph.D Thesis No. 434, Faculty of Agriculture and Applied Biological Sciences, K.U. Leuven, Belgium. 2000; 123.
6. Dwivedi JL. Status Paper on rice in Uttar Pradesh. 2018; 1-2, 8-9.
7. Food and Agriculture Organization of the United Nations (FAO). Water for Sustainable Food and Agriculture—A Report Produced for the G20 Presidency of Germany. 2017.
8. Hardacre AK and Turnbull HI. “The growth and development of maize (*Zea mays* L.) at five temperatures,” *Annals of Botany.* 1986; 58:779–787.
9. Hatfield JL, and Prueger JH. Temperature extremes: effect on plant growth and development. *Weather and Climate Extremes.* 2015; 10: 4-10.
10. Hatfield JL, Boote KJ, Kimball BA, Ziska LH, Izawralde RC, Ort D et al. Climate Impacts on Agriculture: Implications for crops Production. USDA-ARS/UNL Faculty.1350, Nebrasaka,2011.
11. Hendrey GR, & Kimball BA. The FACE program. *Agric. Forest Meteorol.* 1994; 70:3–14.
12. Hoogenboom G, Jones JW, Porter CH, Wilkens PW, Boote KJ, Batchelor WD et al. Decision Support System for Agrotechnology Transfer version 4.0. Volume 1: Overview. University of Hawaii, Honolulu, 2003; HI. 2.
13. Hoogenboom G, Jones JW, Wilkens PW, Porter CH, Boote KJ, Hunt LA. Decision Support System for Agrotechnology Transfer (DSSAT). 2012.
14. Hundal SS and Kaur P. Environment and the effect of environmental stresses on potential production of major cereal crops in Punjab, in Proceedings of the International Conference on Sustainable Agriculture, Haryana Agricultural University, Hisar, India. January 1995; pp. 11–12,
15. Hunsaker DJ, Kimball BA, Pinter PJ et al. CO₂ enrichment and soil nitrogen effects on wheat evapotranspiration and water use efficiency, *Agricultural and Forest Meteorology.* 2000; 104(2):85–105.

16. International Benchmark Sites Network for Agrotechnology Transfer. The IBSNAT Decade. Department of Agronomy and Soil Science, College of Tropical Agriculture and Human Resources, University of Hawaii, Honolulu, Hawaii. 1993.
17. Jones JW, Hoogenboom G, Porter CH, Boote KJ, Batchelor WD, Hunt LA, Wilkens PW, Singh U, Gijsman AJ, Ritchie JT. The DSSAT cropping system model. *European Journal of Agronomy*. 2003; 235- 265.
18. Jones JW, Tsuji GY, Hoogenboom G, Hunt LA, Thornton PK, Wilkens PW et al. Decision support system for agrotechnology transfer; DSSAT v3. In: Tsuji GY, Hoogenboom G, Thornton PK (Eds.), *Understanding Options for Agricultural Production*. Kluwer Academic Publishers, Dordrecht, Netherlands, 1998; pp. 157/177.
19. Karim Z, Ahmed M, Hussain S. and Rashid KB. Impact of climate change on the production of modern rice in Bangladesh. *Implications of Climate Change for International Agriculture Crop Modeling Study*. 1994; 1-11.
20. Moomaw JC, and Vergara BS. *The environment of tropical rice production*, 1965.
21. Nyang'au W.O. , Mati B.M. , Kalamwa K. , Wanjogu R.K. , and Kiplagat L.K. Estimating Rice Yield under Changing Weather Conditions in Kenya Using CERES Rice Model. *International Journal of Agronomy*. (2014); Volume 2014, Article ID 849496, 12 pages.
22. Reyes BG, De Los, Myers SJ, McGrath JM. Differential induction of glyoxylate cycle enzymes by stress as a marker for seedling vigor in sugar beet (*Beta vulgaris*), *Molecular Genetics and Genomics*. 2003; 269(5):692-698.
23. Ritchie JT, Alocilja EC, Singh U, and Uehara G. "IBSNAT and the CERES-rice model," in *Weather and Rice: Proceedings of the International Workshop on the Impact of Weather Parameters on Growth and Yield of Rice*, International Rice Research Institute, Manila, Philippines. 1987; 7–10 April 1986, pp. 271–281
24. Schiller J, Hatsadong and Doungsila K. A history of rice in Laos. In J M Schiller, M B Champhengxay, B Lindquist, and S Appa Rao. eds., *Rice in Laos*. 2006; pp. 9–28. Los Baños, Philippines: IRRI.
25. Tsuji G Y, Hoogenboom G, Thornton PK. *Understanding Options for Agricultural Production*. Berlin: Springer. 1998.
26. Uehara G. and Tsuji, GY. Overview of IBSNAT. In: Tsuji, G.Y., Hoogenboom, G. and Thornton, P.K., Eds., *Understanding Options for Agricultural Production*, Springer, Dordrecht. 1998; 1-7.
27. Van Keulen H. Potential wheat yields in Zambia—a simulation approach, *Agricultural Systems*. 1984; 14(3):171–192.