

Influence of Calcium and Gibberellic Acid on Growth, Yield and Economics of Summer Groundnut (*Arachis hypogea* L.)

ABSTRACT

A field experiment was conducted during *Zaid* season (2022) to determine the Influence of calcium and gibberellic acid on growth, yield and economics of summer groundnut (*Arachis hypogea* L.). Experiment was laid out in Randomized Block Design (RBD) with 3 replicates of 10 treatments. The results showed that treatment 9 of [Ca (40kg/ha) + GA₃ (120ppm)] recorded significantly higher plant height (60.4 cm), more dry weight (45.95g), maximum number of nodules/plant (16.4), maximum number of pods/plant (24.60), maximum number of kernels/pod (1.93), higher seed index (37.8 g), higher seed yield (2.02t/ha), higher haulm yield (4.11 t/ha) and higher harvest index (32.83 %). Maximum gross returns (1,25,778.00 INR/ha), maximum net returns (87,853.50 INR/ha) and highest benefit cost ratio (2.32) was also recorded in treatment 9 of Ca (40kg/ha) + GA₃ (120ppm) as compared to other treatments.

Keywords: Calcium, Economic, Groundnut, Gibberellic acid, Growth and Yield.

1. Introduction

The oilseed sector plays an important role in India's agriculture and economy (DOR, 2013). Nine oil annual seed crops serve as the main source of edible oil (DRMR, 2015). Among soyabeans, peanuts, canola and mustard are the main contributors. Oilseeds are grown mainly on poor soils with a lot of rain. Peanuts are prized for their high oil content and edible seeds. It is the fourth most important source of edible oil and the third most important source. The world's most important vegetable protein source which contain 42-52% oil and 22-30% protein on a dry seed basis, phosphorus, calcium, magnesium and potassium (Savage and keenan, 1994). Peanut are not only a source of income for farmers, but also an excellent source inexpensive source of high-quality dietary protein and oil found in many Ghanaian diets (Asibuo *et al.*, 2008).

Peanut are not only an important oilseed crop of India, but also a major agricultural export. Globally, peanuts cover 31.5 million hectares with a production of 53.6 million tons

and a productivity of 1701 kg/ha (FAOSTAT, 2020). With annual all-season coverage of 55.71 lakh hectares, globally, India ranks first in Groundnut area under cultivation and is the second largest producer in the world with 102 lakh tonnes with productivity of 1831 kg/ha in 2020-21 (agricoop.nic.in). In *Kharif* 2021-22, groundnut production was 82.54 lakh tonnes in an area of 49.14 lakh hectares (agricoop.nic.in). Groundnut is cultivated in one or more (*kharif*, *rabi* and summer) seasons, but nearly 90% of acreage and production comes from *kharif* crop (June-October). During 2019-20 total area coverage under groundnut in Uttar Pradesh 93822.00 hectares with a production of 88371 tonnes and the productivity 940kg/ha (DAC, 2020).

Despite the importance of this crop, yields remain below 1.0 t/ha, a long way off. Less than potential yield of 2-3t/ha. This impacted peanut production, income welfare of peanut farmers. It is not entirely clear whether this low-yield problem is due to declining soil fertility or changing climatic conditions. A lack of calcium increases the rate of seed breakage (empty pods or “pops”) and improperly packed pods (Ntare *et al.*, 2008). It also results in broken or shrivelled fruits, including the darkened plumules and production of seedless pods (Singh and Oswalt, 1995). A sufficient amount of Ca should be present in the soil from early flowering in crop production (Kamara, *et al.*, 2011).

Effect of Plant growth regulators in manipulating physiological processes in plant production, germination, vitality, absorption of nutrients from soil, photosynthesis, respiration, degradation of assimilates, growth inhibition, defoliation, post-harvest ripening (Rahman and Nath (1993); Hossain *et al.* (2015).

Gibberellins are large group of plant hormones that stand alongside auxins are one of the main groups of plant regulators (Bethke, 1998). They are all physiologically different activity and structure, and the first gibberellin identified was gibberellic acid (GA₃). Gibberellins widely involved in all stages of plant growth and development, from seed germination to senescence. They promote seed germination; stimulate stem elongation, leaf enlargement, flowering, pollen and seed development, retardation and inhibition of maturation, aging (Mshelmbula, *et al.*, 2021). Given the above facts, the present investigation entitled, “Influence of calcium and gibberellic acid on the growth, yield and economics of summer groundnut (*Arachis hypogaea* L.)”

2. MATERIALS AND METHODS:

This experiment was conducted at the Crop Research Farm during 2022 *Zaid* season. Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P). The soil of the field constituting a part of central gangetic

alluvium is neutral and deep. The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH 7.8), low level of organic carbon (0.62%), available N (225 Kg/ha), P (38.2 kg/ha), K (240.7 kg/ha) and zinc (2.32 mg/kg). Treatment consists of three calcium stages (20, 30,40kg/ha) and gibberellic acid (40, 80, 120ppm/ha). The treatment combinations are T1- Ca (20kg/ha) + GA₃ (40ppm), T2 - Ca (20kg/ha) + GA₃ (80ppm), T3 - Ca (20kg/ha) + GA₃ (120ppm), T4 - Ca (30kg/ha) + GA₃ (40ppm), T5 - Ca (30kg/ha) + GA₃ (80ppm), T6 - Ca (30kg/ha) + GA₃ (120ppm), T7 - Ca (40kg/ha) + GA₃ (40ppm), T8 - Ca (40kg/ha) + GA₃ (80ppm), T9- Ca (40kg/ha) + GA₃ (120ppm), T10- Control N:P:K (20:60:40 Kg/ha). The growth, yield and economics were recorded at 80 DAS from randomly selected plants in each plot. The data were calculated and analysed by following statistical methods Gomez and Gomez (1984).

3. RESULT AND DISSCUSSION

Growth parameters

Plant height (cm)

The data revealed that significantly higher plant height (60.43 cm) was recorded in treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)]. However, treatment 6 [Ca (30 kg/ha) + GA₃ (120 ppm)] which was found statistically at par with treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)] (Table 1). The significant and higher plant height was observed with the application of calcium (40kg/ha) calcium increases plant nutrient supply, played a important role in photosynthesis, carbohydrate metabolism, protein synthesis and synthesis growth stimulators, consequences of cell division, and cell elongation with would have resulted in increased height. These similar results are agreement with those of Mansingh *et al.* (2018). Additionally, higher plant heights were observed when GA₃ was used (120 ppm) could be due to the application of gibberellic acid *via* leaves, rise the length of the hypocotyl and the length of the two nodes immediately above as a result, it affects the height of the plant. These results are in accordance with Emongor (2007).

Number of nodules/plant

The data found that significantly higher number of nodules/plant (16.40 nodules/plant) was recorded in treatment 9 [Ca (40kg/ha) + GA₃ (120 ppm)]. However, treatment 6 [Ca (30kg/ha) + GA₃ (120 ppm)] which was found statistically at par with treatment 9 [Ca (40kg/ha) + GA₃ (120 ppm)] (Table 1). Significantly higher number of nodules/plant was observed with the application of GA₃ (120ppm), which might be due to foliar application of GA₃ increases plant vitality and strengthens stems. Current results are

close proximity to **Senthil *et al.* (2004)**.

Plant dry weight (g)

Results revealed that significantly higher plant dry weight (45.71 g) was recorded in treatment 9 [Ca (40kg/ha) + GA₃ (120 ppm)]. However, treatment 6 [Ca (30kg/ha) + GA₃ (120 ppm)] which was found statistically at par with treatment 9 [Ca (40kg/ha) + GA₃ (120 ppm)] (Table 1). Significant and greatest plant dry weight was observed with calcium application (40kg/ha), possibly due to the application of calcium sources, which increased the total dry weight and yield of peanuts due to a reduction in percentage of unfilled seeds. There were similar reports as agreed by **Kamara *et al.* (2011)**. Further, higher plant dry weight was observed with the application of GA₃ (120ppm), this may be because GA₃ is one of these growth regulators it as a positive effect on plant as it promotes vegetative growth and ultimately increased plant dry weight. Similar findings were confirmed by **Islam *et al.* (2021)**.

Crop Growth Rate (g/m²/day)

The data recorded that during 60-80 DAS, there was no significant difference between the treatments. However, highest crop growth rate (17.85 g/m²/day) was recorded in treatment 4 Ca (30kg/ha) + GA₃ (40ppm) (Table 1).

Relative Growth Rate (g/g/day)

The data revealed that during 60 – 80 DAS, there was no significant difference between the treatments. However, highest relative growth rate (0.0183 g/g/day) was recorded in treatment 10 [control (RDF)] (Table 1).

Yield Attributes:

Number of pods/plant

The data found that significantly higher number of pods/plant (24.60) was recorded in the treatment 9 [Ca (40kg/ha) + GA₃ (120 ppm)]. However, the treatment 6 [Ca (30 kg/ha) + GA₃ (120 ppm)] which was found statistically at par with treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)] (Table 2). more pods/plant was observed with calcium application (40kg/ha), this might be due to soils that have been amended with gypsum as a calcium source. Contributed to better development of peanut pods and kernels, leading to higher number of pods/plant. Similar findings were earlier reported by **Kabier *et al.* (2013)**

Number of kernels/pod

The data showed that significantly higher kernels/Pod (1.93) were recorded in the treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)]. However, the treatment 6 [Ca (30 kg/ha) + GA₃ (120 ppm)] which was found statistically at par with treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)] (Table 2). Significant and higher kernels/pod was observed with the application of calcium (40kg/ha), this might be due to calcium maintaining cell integrity, membrane permeability activates and participates in a number of enzymes involved in cell division, protein synthesis and carbohydrate transfer, resulted in development of kernels/pod in plant. Similar results were also noticed by **Rajanarasimha *et al.* (2021)**. Further higher kernels/pod was observed with application of GA₃ (120ppm), which might be due to the plant growth regulators such as gibberellic acid may be involved in formation of seeds in pods and their optimum nutrition resulted in fewer number of aborted seeds and thus maximizes fertile seeds/pods survival for oilseed rape and mustard. This result corroborates the one reported by **Akter *et al.* (2007)**.

Seed index (g):

The data inferred that significantly higher seed index (37.80 g) was recorded in the treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)]. However, treatment 6 [Ca (30 kg/ha) + GA₃ (120 ppm)] which was found statistically at par with the treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)] (Table 2). Significant and higher seed index was observed using calcium (40kg/ha), this may be because peanuts are a calcium-loving crop and Ca above 90% in peanut pods is absorbed during the pod formation stage, resulting in calcium absorption from the soil important for peanut embryo and pod development. Similar results were also supported by **Yang *et al.* (2017)**. Further the higher seed index was observed with application of GA₃ (120ppm), this might be due to GA₃ prolonging the grain filling time as a result, it ensures long-term transport of photo-assimilates into grains, increases the 100 seed weight. This is in accordance with previous findings by **Wang *et al.* (2006)**.

Seed yield (t/ha):

The data stated that significant and higher seed yield (2.02 kg/ha) was recorded in the treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)]. However, the treatment 6 [Ca (30 kg/ha) + GA₃ (120 ppm)] which was found statistically at par with treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)] (Table 2). Significant and higher seed yield was observed with application of calcium (40kg/ha) might be due to the fact that calcium plays an important role in reproduction development of peanut crops with increased seed yield. Similar findings were in **accordance** with **Sagar *et al.* (2020)**. Further the higher seed yield was

observed with application of GA₃ (120ppm), which may be due to the positive effect of GA₃ on improving yield by transferring more photo-assimilates to seeds. This is in agreement with the results of **Varshitha et al. (2022)**.

Stover yield (t/ha)

The data reported that significant and higher haulm yield (4.11 t/ha) was recorded with the treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)]. However, the treatment 6 [Ca (30 kg/ha) + GA₃ (120 ppm)] which was found to be statistically at par to the treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)] (Table 2). Significant and higher haulm yield was observed with the application of calcium (40kg/ha), which may be attributed to increased pod yield and haulm and higher Ca concentrations with increased gypsum content as a source of calcium, as a result the uptake of this nutrient by pod and haulm is increased, ultimately resulting in recording. Similar results were supported by **Patro et al. (2016)**. Further the higher haulm yield was observed with the use of GA₃ (120ppm). Gibberellic acid leads to increased plant height and branch numbers which also translated into higher number of leaves which invariably made up the haulm. These results were in conformity with those of **Harb (1992)**.

Harvest index (%)

Significant and higher harvest index (32.83 %) was recorded in the treatment 9 (Ca (40 kg/ha) + GA₃ (120ppm)]. However, the treatments 3 [Ca (20kg/ha) + GA₃ (120ppm)] and treatment 5 [Ca (30 kg/ha) + GA₃ (80ppm)] and treatment 6 [Ca (30kg/ha) + GA₃ (120 ppm)] and treatment 7 [Ca (40kg/ha) + GA₃ (40 ppm)] and treatment 8 [Ca (40kg/ha) + GA₃ (80 ppm)] was found statistically at par with treatment 9 [Ca (40 kg/ha) + GA₃ (120 ppm)]. (Table 1). Significant and higher harvest index was observed with the application of GA₃ (120ppm), which might be due to the higher harvest index indicating GA₃ application acceleration of anabolic feed to the sink. Similar findings have also been reported earlier by **Akter et al. (2007)**.

Economics:

The results stated that maximum gross return (1,25,778.00 INR/ha), higher net return (87,853.50 INR/ha) and higher benefit cost ratio (2.32) was recorded in treatment 9 [Ca (40kg/ha) + GA₃ (120ppm)] as compared to other treatments (Table 3). Significant and maximum B:C ratio was observed with the application of Calcium (40kg/ha), which might be due to the application of gypsum as a source of Ca increase net returns and B:C ratio of harvesting peanuts with optimal nutrient utilization by gypsum during harvest leads to better growth and development of pods. Similar results were found by **Sagar et al. (2020)**.

Table 1: Influence of calcium and gibberellic acid on growth parameters of groundnut

S. No	Treatments	Plant height (cm)	Number of nodules/plant	Plant dry weight (g)	CGR (g/m ² /day)	RGR (g/g/day)
1	Ca 20 kg/ha + GA ₃ 40 ppm	56.60	13.73	41.90	17.23	0.0143
2	Ca 20 kg/ha + GA ₃ 80 ppm	58.23	14.70	43.15	17.39	0.0140
3	Ca 20 kg/ha + GA ₃ 120 ppm	58.90	15.03	43.65	17.47	0.0140
4	Ca 30 kg/ ha + GA ₃ 40 ppm	57.13	14.06	42.44	17.85	0.0147
5	Ca 30 kg/ha + GA ₃ 80 ppm	59.26	15.53	43.95	16.84	0.0130
6	Ca 30 kg/ha + GA ₃ 120 ppm	60.10	16.16	45.95	17.57	0.0130
7	Ca 40 kg/ ha + GA ₃ 40 ppm	58.03	14.23	42.73	17.77	0.0140
8	Ca 40 kg/ha + GA ₃ 80 ppm	59.53	15.86	44.84	17.79	0.0137
9	Ca 40 kg/ha + GA ₃ 120 ppm	60.43	16.40	45.71	17.46	0.0130
10	Control (RDF)	55.23	13.16	41.06	16.49	0.0183
	F-test	S	S	S	NS	NS
	SEm (±)	0.13	0.07	0.15	0.24	0.0002
	CD at 5%	0.41	0.23	0.45	0.72	0.0007

Table 2: Influence of calcium and gibberellic acid on yield and yield attributes of groundnut

S. No	Treatments	Number of pods/plant	Number of kernels/pod	Seed index (g)	seed yield (t/ha)	Haulm yield (t/ha)	Harvest index (%)
1	Ca 20 kg/ha + GA ₃ 40 ppm	20.15	1.27	31.26	1.62	3.54	31.47
2	Ca 20 kg/ha + GA ₃ 80 ppm	22.45	1.48	33.63	1.78	3.75	32.17
3	Ca 20 kg/ha + GA ₃ 120 ppm	23.08	1.59	35.20	1.82	3.82	32.31
4	Ca 30 kg/ ha + GA ₃ 40 ppm	21.23	1.33	31.83	1.66	3.59	31.63
5	Ca 30 kg/ha + GA ₃ 80 ppm	23.55	1.66	35.90	1.85	3.88	32.28
6	Ca 30 kg/ha + GA ₃ 120 ppm	24.05	1.87	37.66	1.96	4.10	32.33
7	Ca 40 kg/ ha + GA ₃ 40 ppm	21.57	1.41	33.03	1.73	3.63	32.31
8	Ca 40 kg/ha + GA ₃ 80 ppm	23.87	1.77	36.43	1.89	3.93	32.56
9	Ca 40 kg/ha + GA ₃ 120 ppm	24.60	1.93	37.80	2.01	4.11	32.83
10	Control (RDF)	18.63	1.20	31.13	1.58	3.46	31.35
	F-test	S	S	S	S	S	S
	SEm (±)	0.19	0.02	0.16	18.19	24.75	0.19
	CD at 5%	0.57	0.07	0.50	0.05	0.07	0.58

Table 3: Influence of calcium and gibberellic acid on the economics of groundnut

S. No.	Treatment combinations	Cost of cultivation (INR/ha)	Gross returns (INR/ha)	Net returns (INR/ha)	B:C ratio
1	Ca 20 kg/ha + GA ₃ 40 ppm	32,515.80	93,504.00	60,988.20	1.88
2	Ca 20 kg/ha + GA ₃ 80 ppm	34,915.80	1,02,748.00	67,832.20	1.94
3	Ca 20 kg/ha + GA ₃ 120 ppm	37,315.80	1,05,284.00	67,968.20	1.82
4	Ca 30 kg/ ha + GA ₃ 40 ppm	32,820.16	95,920.00	63,099.80	1.92
5	Ca 30 kg/ha + GA ₃ 80 ppm	36,220.16	1,06,754.00	70,533.80	1.95
6	Ca 30 kg/ha + GA ₃ 120 ppm	37,620.16	1,13,044.00	75,423.80	2.00
7	Ca 40 kg/ ha + GA ₃ 40 ppm	33,124.52	99,866.00	66,741.50	2.01
8	Ca 40 kg/ha + GA ₃ 80 ppm	35,524.52	1,09,062.00	73,537.50	2.07
9	Ca 40 kg/ha + GA ₃ 120 ppm	37,924.52	1,25,778.00	87,853.50	2.32
10	Control (RDF)	29,507.20	74,552.00	45,044.80	1.53

CONCLUSION

Based on above findings it can be concluded that application of calcium at 40 kg/ha along with foliar application of Gibberellic acid 120 ppm (Treatment 9), has performed better in growth, yield and benefit cost ratio.

REFERENCES

1. **Asibuo, J.Y., Akromah, R., Adu-Dapaah, H.K. and Safo-Katanka, O, (2008).** Evaluation of nutritional quality of groundnut (*Arachis hypogaea* L.) from Ghana. *African Journal of Food and Agriculture Nutrition and development*. **8**(2):133-150.
2. **Akter A, Islam E, Karim R, Razzaque AHM.** (2007) Effect of GA₃ on growth and yield of mustard. *International Journal of Sustainable Crop Production*.**2**(2): 16-20.
3. **Bethke, P.C., Jones, R.L. (1998).** Gibberellin signaling. *Current Opinion in Plant Biology*. **1**: 440–446.
4. **DOR. (2013).** Vision 2050 ICAR- Indian Institute of Oilseeds Research (IIOR) Andhra Pradesh, India p33.
5. **DRMR. (2015).** Vision 2050 ICAR- Directorate of Rapeseed- Mustard Research, Bharatpur India p26.
6. **Department of Agriculture and Cooperation (2020).**
<https://agricoop.nic.in>
7. **Emongor, V. (2007).** Gibberellic acid (GA₃) influence on vegetative growth, nodulation and yield of cowpea (*Vigna unguiculata* L. Walp). *Journal of Agronomy*. **6**(4): 509-517.
8. **Food and Agriculture Organization Corporate Statistical Database (2020).**
<http://www.fao.org.in>
9. **Gomez K.A. and Gomez, A. A. (1984).** Statistical procedures for agricultural research: *John Wiley and Sons. New Delhi*, 680.
10. **Hossain, M.I., Mannam, M.A. and Kareem, M.A. (2015).** Salicylic acid and gibberellic acid ameliorate the adverse effect of salinity on chickpea, *Bangladesh Agronomy Journal*.**18**(1): 81-88.
11. **Harb EZ (1992).** Effect of soaking seeds in some growth regulators and micronutrients on growth, some chemical constituents and yield of faba beans and

- cotton plants. supplement, *Bulletin, Faculty of Agriculture-Cairo University* **43**: 429-452.
- 12. Islam MR, Hasan M, Akter N, Akhtar S. (2021).** Cytokinin and gibberellic acid alleviate the effect of waterlogging in mungbean (*Vigna radiata* L). *Journal Clean WAS.* **5**:21-26.
 - 13. Kamara, A., Olympio, and Asibuo, J. (2011).** Effect of calcium and phosphorus fertilizer on the growth and yield of groundnut (*Arachis hypogaea* L.). *International Research Journal of Agricultural Science and Soil Science.***1**(8) pp. 326-331.
 - 14. Kabir R, Yeasmin S, Islam AKMM, Sarkar MAR (2013).** Effect of phosphorus, calcium and boron on the growth and yield of groundnut (*Arachis hypogaea* L.), *International Journal of Bio-Science and Bio-Technology*, **5**(3): 51-60.
 - 15. Kamara, E. G., Olympio, N. S. and Asibuo, J. Y. (2011).** Effect of calcium and phosphorus fertilizer on the growth and yield of groundnut (*Arachis hypogaea* L.). *International Research Journal of Agricultural Science and Soil Sciences*, **1**:326-331.
 - 16. Mazid, M. and N. Naqvi, (2014).** Key approach of plant growth regulator application in searching of best time for enhancing nitrogen fixation capacity of chickpea cultivar DCP 92-3. *Unique Journal of Ayurvedic and Herbal Medicines*,**2**(05): 8-18.
 - 17. Mshelmbula, B.P., Ogale, E., Bello, S., Kana, H.A., Sulayman, M.Y., Allahnana, M.H., Sirajo, S.A. (2021).** Impact of gibberellic acid (GA₃) on growth, yield and nodulation on two accessions of cowpea (*Vigna unguiculata* (L.) Walp). *Journal of Applied Sciences and Environmental Manage.* **25**(8):1435-1439.
 - 18. Mary Varshitha K, Vikram Singh, Shruthi G G, and Singh A.C. (2022).** Effect of plant growth regulators and spacing on growth and yield of chickpea (*Cicer arietinum* L.). *International Journal of Environment and Climate Change.***12**(10): 614-619.
 - 19. Mansingh, M. D. I. and Suresh S. (2019).** Effect of difference sources and levels of lime application on yield, nutrient availability and uptake of rice in acidic soils. *Journal of Agriculture and Ecology*, **7**: 73-87.
 - 20. Ntare, B.R., Diallo, A.T., Ndjeunga, A.T.W. and aliyar, F. (2008).** Groundnut seed production manual. Patancheru 502324, Andhra Pradesh, India. *International Crops Research institute for the Semi-Arid Tropics* (ICRISAT). 20pp

21. **Patro H, Ray M. (2016).** Effect of rate and time of gypsum application on yield, economics and nutrient uptake in groundnut. *New Agriculturist*. **27**(2): 1-6.
22. **Rahman, M.A. and Nath, K.K. (1993).** Effect of seed treatment of IAA and GA3 on sex expression, fruit character and yield of bottle gourd, *Bangladesh Journal of Sciences*. **5**(2):57-63.
23. **Rajanasimha M, Rajesh Singh and Ekta Singh (2021).** Effect of sulphur and calcium on growth and yield of groundnut (*Arachis hypogea* L.), *The Pharma Innovation Journal*. **10**(9): 1685-1688.
24. **Singh, F. and Oswalt, D.L. (1995).** Groundnut production practices. Skill development series number 3. ICRSAT Training and Fellowship Program *International Crops Institute for the Semi-Arid Tropics*, Patancheru, Andhra Pradesh 502 324, India
25. **Sagar D.R.M.S., Dawson J, and Ramasahayam, Reddy U K. (2020).** Effect of phosphorus and gypsum on growth, yield and economics of groundnut (*Arachis hypogea* L.). *International Journal of Current Microbiology and Applied Sciences*. **9**(10): 1635-1638.
26. **Savage, G. P. and Keenan, J.I. (1994).** The composition and nutritive value of groundnut kernels. ISBN 0 412 408201.
27. **Senthil NK, Jaya KR. (2004).** Effect of growth regulators on yield and quality of greengram (*Vigna radiate*) under graded levels of nitrogen. *Madras Agricultural Journal*, **91** (1-3):92-95.
28. **Wang, F., Cheng, F., and Zhang, G. (2006).** The relationship between grain filling and hormone content as affected by genotype and source-sink relation. *Plant Growth Regul.* **49**, 1-8. Doi: 10.1007/s10725-006-0017-3.
29. **Yang, S., Li, L., Zhang, J., Geng, Y., Guo, F., Wang, J., Meng, J., Sui, N., Wan, S. and Li, X. (2017).** Transcriptome and Differential Expression Profiling Analysis of The Mechanism of Ca²⁺ Regulation in Peanut (*Arachis hypogaea*) pod development. *Frontiers in Plant Science*, **8**: 1609.