

Original Research Article
A study to assess the prevalence of Non-suicidal Self Injury and reasons for that behaviour among adults and their coping

ABSTRACT

ABSTRACT:

Aims: To study prevalence and characteristics of Non-suicidal self-injury (NSSI) and functional purpose of that behaviour among Adults and their level of coping.

Study design: Cross An analytical cross-sectional study.

Place and Duration of Study: The study is conducted in central medical university, Gujarat, India over a period of 30 days.

Methodology: Study was conducted among students of Central Medical University.195 participants were enrolled by consecutive random sampling. They were given a Google form to fill out Demographic details. The form also has an Inventory for NSSI(ISAS) to assess their NSSI behaviour and Brief Resilient Coping Scale (BRCS) to assess coping.

Results: Out of 195 participants, prevalence of NSSI behaviour is 49(25.12%). Among these 20(40.82%) are male & 29(59.18%) are female. Among sociodemographic data type of family is statistically significant(P value- 0.039). Mean age at which participants start self-harming is 16.6 ± 2.89 years. During NSSI behaviour 51.02% of participants always feel pain and 16.32% of participants do not feel pain. Prevalence of NSSI behaviour is higher in 3rd or higher birth order (33.33%) compare to 1st(28.31%) and 2nd (19.67%).The most common method of self-harm was cutting (65.3%), followed by banging (61.22%) or hitting self. The most common functional reason for self-harm was affect regulation followed by self-punishment. of participants who are having NSSI behaviour. In this study we found that there is a statistically significant difference between various resilience coping and NSSI behavior.(P- 0.0008).

Conclusion: NSSI behaviour among youth has been growing. The most common method of self-harm is cutting and functional purpose for this behaviour is affect regulation. Individual with nuclear family have higher chance of NSSI behavior than Joint family. Individual with low resilient coping have higher chances of NSSI behavior and similarly high resilient coping is protective for NSSI behavior.

Keywords: [Non-suicidal self injury, Youth, Coping.]

1. INTRODUCTION

Non-suicidal self-injury (NSSI) is defined as intentional, self-inflicted damage to the surface of the body without suicidal intent, which is not socially sanctioned (1). This definition excludes accidental and indirect self-injurious behaviours(disordered eating or drug abuse), suicidal behaviours, also socially accepted behaviours like tattooing,

piercing, or religious rituals. The most common methods endorsed are cutting, scratching, hitting or banging, carving, and scraping (2).

NSSI is being increasingly identified as an important mental health concern even in youth, NSSI is particularly present in mid-adolescence.(3) A study found that although 41.6% of adolescents reported a h/o NSSI, one study reported that 8.0% of youth aged 7 to 16 reported NSSI. who reported NSSI at least five times in the past year.(4,5)

Suicide attempts are not always lethal and NSSI may be lethal(6). NSSI is one of the strongest predictors of suicide and future suicide attempts(7). NSSIs are thought to serve as a function, being used as a way to escape from pain and regulate emotion(8).

This prevalence, or at least the convention of NSSI, may increase with websites like YouTube. Studies investigating the pervasiveness of NSSI videos online revealed 2,140 videos in 2009 and over 5,000 in 2012. A Google search in 2009 revealed more than 15 million results, and almost 87 million in 2014(9,10).

Self-harm is associated with stress and emotional dysregulation. Most of the studies focused on adolescents. So, this study intended to assess the behaviour and characteristics of non-suicidal self-harm in adults as stress was found to be high among them and which is the strongest predictor of suicide and future suicide attempts. So we can identify and early intervention can be done to prevent suicide.

In Adolescence, most common cause of death is Suicidal, and NSSI is a very strong predictor of Suicide.(11) Pearson's correlation analysis revealed that mindfulness was significantly and positively associated with students' perceived level of coping. (14) In Adolescence higher amount of successful suicidal attempts is being performed. This research will help to find the prevalence and characteristics of non-suicidal self-injury(NSSI) and the functional purpose of that behaviour among Adults and their level of coping.

- **NSSI in India:**

There is very less existing literature on self-harming behaviour in India. Though a small number of studies have investigated behaviours similar to NSSI, inferring the prevalence of NSSI using these studies may be difficult, as they use definitions that may not allow for distinctions between different kinds of self-injurious behaviour. 1-year prevalence of NSSI in India has been reported to vary between 31% and 34%. Bholra and colleagues (2017) point out , both studies used a very broad definition of NSSI. Therefore, the resulting one-year prevalence may be inflated. Evidence regarding gender differences is also conflicting, as one of the studies reported a lack of gender differences (12), whereas the other study reported a higher one-year prevalence of NSSI in females than in males. Consistent with the international literature, the age of onset of NSSI in India has been reported to be around 15.9 years. Given the small number of studies and their inconsistent findings, further research is necessary to understand NSSI in India(13).

It has been evident in the available literature that self-bruising behaviour was more commonly reported in India and their prevalence has been found out but the reason for their behaviour has been overlooked. As we are moving forward in the 20th century NSSI has been increasing among youth because of multiple reasons. As NSSI is a predictor of Suicide (11) it can be helpful to find out the reason as it can help to prevent future suicide attempts. Yet there have been few studies stating NSSI and Coping relations. The present study is focused on finding a correlation between NSSI and coping.

This study is conducted to determine the Prevalence & to identify the characteristics of NSSI and also to find the functional purpose of NSSI. We will also be able to assess the level of coping among adults and its association between coping and NSSI.

1.1 OBJECTIVES

1. This study Prevalence of NSSI among youth of central university students.
2. To identify the characteristics of NSSI.
3. To find the functional purpose of NSSI.
4. To assess the level of coping among youth and its association between coping and NSSI.

2. METHODOLOGY

The study is a cross-sectional study. The study is conducted by the Department of Psychiatry of Central medical university, Gujarat, India over a period of 30 days.

Inclusion criteria adopted for the present study were participants in an age group between 18 and 25 years, those who were well versed in English Language and who belong to university Campus. We Included students of Medical(Bachelor of Medicine and Bachelor of Surgery), Dental(Bachelor of Dental Surgery), Ayurveda(Bachelor of Ayurvedic and Medicine and Surgery), Physiotherapy(Bachelor of Physiotherapy), Nursing(Bachelor of Nursing Science) and Pharmacy(Bachelor of Pharmacy and Diploma of Pharmacy). Exclusion criteria adopted for the present study were those who did not wish to participate.

The sample size was calculated by using the sample size calculation formula for cross-sectional study with help of a sample size calculator. The formula is $n' = NZ^2 P(1-P)/d^2 (N-1) + Z^2 P(1-P)$. In this formula "n" is Sample size (for finite population), "N" is the population size, "Z" is the statistic for level of confidence, "P" is expected proportion and "d" is Precision. As per calculation required Sample size is 153.

Data were collected from study participants by using Google Forms. Before starting the survey, all participants were provided with details of the time taken to complete the survey, the nature of the survey and the information that fills in the

survey implies the provision of informed consent by participants. A survey questionnaire was circulated using Google links. Identity of every participant was strictly kept anonymous.

A survey was conducted in the English language. In the survey, two scales are used for data collection:

1) NSSI(ISAS): The ISAS assesses 13 functions of NSSI, as well as the frequency of 12 NSSI behaviours. ISAS factors exhibited excellent internal consistency and expected correlations with both clinical constructs (e.g., borderline personality disorder, suicidality, depression, anxiety) and contextual variables (e.g., a tendency to self-injure alone). The ISAS may be useful in research and treatment contexts as a comprehensive measure of NSSI functions.

2) The Brief Resilient Coping Scale(BRCS): captures tendencies to cope with stress adaptively. The scale focuses on the tendency to effectively use coping strategies in flexible, committed ways to actively solve problems despite stressful circumstances. It contains 4 items and each item contains a 5-point Likert scale response, ranging from 1=does not describe me at all to 5 describes me very well.

Data was then analysed and the mean along with the standard deviation was calculated to find out the prevalence. Chi-square test was used to determine the level of significance. The level of Significance was set at 0.05.

3. RESULTS

In the present study 199 participants approach for the study, 4 participants denied giving consent hence 195 participants were studied. The mean age of participants is 21.65(\pm 1.85) yrs. 46.15% are Male while 53.85% are female. Among Participants, 79.48% belong to the Medical field while 20.52% belong to paramedical and other fields (such as Pharm D, MBA). 67.69% belong to a nuclear family whereas 32.3% belong to joint families. 89.74% belong to upper socioeconomic status, 8.71% belong to middle socioeconomic status and 1.53% of participants belong to lower socioeconomic status according to Modified Prasad Classification. 57.94% of participants have 1st birth order, 31.28% of participants have 2nd birth order whereas 10.78% of participants are of the third birth order or above.(Table 1)

Table 1: Association between Sociodemographic variables & NSSI

Sociodemographic variables		NSSI		Total	Chi-square	P value
		Present	Absent			
Gender	Male	20	70	90	0.75	0.386
	Female	29	76	105		
Education	Medical	41	114	155	0.703	0.401
	Paramedical& Others	8	32	40		
Family	Nuclear	39	93	132	4.24	0.039*

	Joint	10	53	63		
Socio-economic class	Upper	40	135	175	-	-
	Middle	7	10	17		
	Lower	2	1	3		
Birth order	1 st	30	81	113	1.9	0.38
	2 nd	12	49	61		
	≥ 3 rd	7	14	21		

(p-value<0.05 is statistically significant)

Out of 195 participants, 49 participants were having NSSI behaviour, so the prevalence of NSSI behaviour is 25.12%. Among these 20(40.82%) are male & 29(59.18%) are female. Out of the participants doing NSSI, 83.67% belong to the medical field, and 16.33% belong to the paramedical field and other professions. Among sociodemographic data type of family is statistically significant(P value- 0.039). All other demographic variable are not statistically significant.

Non-Suicidal Self- Harming Behaviour:

Characteristic of NSSI behavior:

The mean age of starting self-harm is 16.6(± 2.89) years. Among participants with NSSI behaviour, 25(51.02%) of participants always feel pain, 16(32.63%) of participants sometimes feel pain and 8(16.32%) participants do not feel pain at all while performing self-harming behaviour. 34(69.38%) of participants perform NSSI always when they are alone, 11(22.44%) do it sometimes when they are alone and 4(8.16%) perform it when they are with someone. 24(48.97%) of participants performed self-harm within 1 hour of their urge to do NSSI, 16(32.65%) performed between 1-12 hours of the urge to do NSSI and 9(18.36%) of participants performed NSSI after 12 hours of their urge.

Table -2: Prevalence of different methods of Self Harm.

Self Harm	Numbers	Percentage
Cutting	32	65.30%
Banging or hitting self	30	61.22%
Pinching	28	57.14%
Interfere with wound healing	26	53.06%
pulling hair	24	48.97%
Biting	22	44.89%
severe scratching	19	38.77%
Rubbing skin with rough surface	15	30.61%
Sticking self with needles	14	28.57%
Burning	12	24.48%
Carving	11	22.44%
Swallowing dangerous substance	9	18.36%

The most common method of self-harm is cutting(65.30%) followed by banging or hitting self(61.22%) while the least common method is swallowing a dangerous substance(18.36%).(Table 2) Among individuals with NSSI behavior there is more than one self harming behavior found.

Functional Purpose of NSSI Behaviour:

Table-3: Functional purpose of NSSI Behaviour:

Functional purpose of NSSI	Mean Score
Affect Regulation	4.14
Interpersonal Boundaries	2.16

Self-Punishment	3.44
Self-Care	1.97
Anti-Dissociation/Feeling-Generation	2.77
Anti-Suicide	2.81
Sensation-Seeking	1.18
Peer-Bonding	1.2
Interpersonal Influence	1.75
Toughness	2.3
Marking Distress	3
Revenge	1.16
Autonomy	2

In this study we found that, most common reason for self-harm was affect regulation(mean-4.14) followed by self-punishment(mean- 3.44). where the least common reason was revenge(mean-1.16). (Table-3)

Association between Coping and NSSI behavior:

Table-4 Association between Coping and NSSI behaviour:

	NSSI	NO NSSI	Total	Chi square	P- value
Low resilient coping	21	33	54	14.058	0.0008*
Medium resilient coping	18	40	58		
High resilient coping	10	73	83		
Total	49	146	195		

(p-value<0.05 is statistically significant)

In this study we found that there is a statistically significant difference between various resilience coping and NSSI behavior.(P- 0.0008). (Table- 4)

4. DISCUSSION

The overall prevalence of NSSI in our study, considering the mean age of the students to be 21.65(±1.85)years. Consistent with the international literature, the age of onset of NSSI in India has been reported to be around 15.9 years. We found that the onset of NSSI was at the mean age of 16.6(±2.89) years. Evidence regarding gender differences is also conflicting, as one of the studies reported a lack of gender differences (12), whereas the other study reported a higher one-year prevalence of NSSI in females than in males. In our study, we do not found statistically significant difference between male and female. In the Demographic variable, only the type of family is statistically significant for NSSI behavior. All other demographic variable are not found statistically significant.

In this study, the highest prevalence of NSSI has been noted amongst Medical field students (83.67%) in comparison to other streams. Although this cannot be considered as standard out of 199 participants who took part in the study, 155 belonged to the medical field.

In this study, the Prevalence of NSSI behaviour is higher in the 3rd or higher birth order(33.33%) compare to subsequently in the 1st birth order(28.31%) and 2nd birth order(19.67%). which is not statistically significant. No other similar study found for the same.

In our study, the most common method of self-harm was found to be cutting (65.3%), followed by banging (61.22%) or hitting self while the least common method was by swallowing a dangerous substance (18.36%).

We also found that the most common reason for self-harm was affect regulation followed by self-punishment where as the least common reason was revenge.

In this study we found that there is a statistically significant difference between various resilience coping and NSSI behavior.(P- 0.0008). Stating that individual with low resilient coping have higher chances of NSSI behavior and similarly individual with high resilient coping is protective for NSSI behavior.

5. CONCLUSION

It has been concluded that, NSSI behaviour among youth has been growing. The most common method of self-harm was found to be cutting. The most common functional purpose for this behaviour is affect regulation. Individual with nuclear family have higher chance of NSSI behavior than Joint family. Individual with low resilient coping have higher chances of NSSI behavior and similarly individual with high resilient coping is protective for NSSI behavior.

6. LIMITATIONS

- Cross sectional study design precludes any causal inferences.
- Subjective bias.
- All the participants are selected from one demographic region that restricting the generalizability of the data.

CONSENT

Prior informed consent was obtained through software 'google form' from the participants before collecting data for using data in future for analysis and publication.

ETHICAL APPROVAL

Prior permission of Sumandeep Vidyapeeth institutional ethics committee (SVIEC) was taken to start the study SVIEC/ON/MEDI/BNPG21/Dec22/216

REFERENCES

1. Ruch DA, Sheftall AH, Schlagbaum P, et al.: Trends in Suicide Among Youth Aged 10 to 19 Years in the United States, 1975 to 2016. *JAMA Netw Open*. 2019; 2(5): e193886.
2. Curtin SC, Warner M, Hedegaard H: Increase in Suicide in the United States, 1999-2014. *NCHS Data Brief*. 2016; (241): 1–8.
3. Center for Disease Control and Prevention: 10 leading causes of death by age group, United States-2017.
4. Zetterqvist M, Lundh L-G, Dahlström Ö, Svedin CG. Prevalence and function of non-suicidal self-injury (NSSI) in a community sample of adolescents, using suggested DSM-5 criteria for a potential NSSI disorder. *Journal of Abnormal Child Psychology*. 2013; 41:759–773. [PubMed: 23344701]
5. Barrocas AL, Hankin BL, Young JF, Abela JRZ. Rates of nonsuicidal self-injury in youth: Age, sex, and behavioral methods in a community sample. *Pediatrics*. 2012; 130:39–45. [PubMed: 22689875]
6. Miller, A., Rathus, J., & Linehan, M. (2007). *Dialectic behavior therapy with suicidal adolescents*. New York: Guilford.
7. Shain, B. (2016). *Suicide and suicide attempts in adolescents*. American Academy of Pediatrics clinical report.
8. Linehan, M. (1993). *Cognitive behavioral therapy of borderline personality disorder*. New York: Guilford.

9. Purington, A., & Whitlock, J. (2010). Non-suicidal self-injury in the media. *The Prevention Researcher*, 17(1), 11-13.
10. Lewis, S. P., Heath, N. L., Michal, N. J., & Duggan, J. M. (2012). Non-suicidal self-injury, youth, and the internet: What mental health professionals need to know. *Child and Adolescent Psychiatry and Mental Health*, 6(13), 1- 13.
11. Poudel A, Lamichhane A, Magar KR, Khanal GP. Non suicidal self injury and suicidal behavior among adolescents: co-occurrence and associated risk factors. *BMC Psychiatry*. 2022 Feb 9;22(1):96. doi: 10.1186/s12888-022-03763-z. PMID: 35139825; PMCID: PMC8827284
12. Bhola P, Manjula M, Rajappa V, Phillip M. Predictors of non-suicidal and suicidal self-injurious behaviours, among adolescents and young adults in urban India. *Asian J Psychiatr*. 2017 Oct;29:123-128. doi: 10.1016/j.ajp.2017.04.024. Epub 2017 May 6. PMID: 29061408.
13. Gandhi A, Luyckx K, Adhikari A, et al. Non-suicidal self-injury and its association with identity formation in India and Belgium: A cross-cultural case-control study. *Transcultural Psychiatry*. 2021;58(1):52-62. doi:10.1177/1363461520933759
14. Heath, N.L., Joly, M. & Carsley, D. Coping Self-Efficacy and Mindfulness in Non-Suicidal Self-Injury. *Mindfulness* 7, 1132–1141 (2016).

UNDER PEER REVIEW