

Antimicrobial Activity of Herbal Plants used for Treatment of Gastroenteritis Infection in Tharaka Nithi County, Kenya

Abstract

Herbals have been used for decades to treat gastrointestinal infections across the world including Tharaka-Nithi County in Kenya. Crude extracts from herbals have long been prescribed by traditional healers for treatment of typhoid, cholera and shigellosis. However, the effectiveness of extracts obtained using different extraction solvents such as methanol, acetone and hexane has not fully been evaluated. This study aimed at evaluating the effect of solvents (Ethanol, hexane and methanol) on the yield of crude extract from plants (*Erythrina abyssinica*, *Aspilia pluriseta*, *Vangueria infausta*, *Ficus sycomorus*, *Carissa edulis*), assess the effect of the solvents on extracted active metabolites and determine the effect of these solvents on extract's bioactivity against *Escherichia coli*, *Vibrio cholerae*, *Shigella flexneri*, and *Salmonella typhi*. Crude extracts were obtained by soaking dried ground plant parts in individual solvents then concentrated by rotary evaporator. The phytochemical screening to detect plant metabolites was done qualitatively. Bioassays to analyze the efficacy of the plant crude extracts against the microbes was carried out in 4x5x3x3 factorial experiment laid out in completely randomized design. Determination of bioactivity of herbal extracts was carried out using minimal inhibition concentration and minimum bactericidal concentration methods. Data obtained on the bioactivity assay (Count on the bacteria colony forming units) was analyzed using Kruskal Wallis test at $\alpha = 0.05$ and medians were compared by Wilcoxon rank sum test in Scientific analysis System version 9.4. Methanol solvent produced higher quantities of crude extracts for all the herbal samples used with *Vangueria infausta* producing the highest extract (5.06g). Most phytochemicals were present in Methanolic extracts compared to hexane and acetone extracts. There was a significant ($p < 0.05$) difference in the bioactivity of different herbal plants against bacterial pathogens at different concentrations of crude extract. The efficacy of plant extract increased progressively from 100ppm to 1000ppm concentration. Methanol is recommended for use in extraction of medicinal plant extracts as it leads to an improved potency as compared to hexane and acetone.

Keywords: Herbal-plants, antimicrobial-activity, cholera, typhoid, shigellosis, Tharaka-Nithi County

I. INTRODUCTION

Medicine is fundamental to healthcare in management and prevention of illnesses given that healthcare is a basic human need to reduce mortality and disease burdens [1] [2]. Access to safe, effective and affordable medicine is among the components of the United Nations' Sustainable Development Goals [3]. Yet, the occurrence of low-quality medicines continues to be a pervasive challenge due to increased drug resistance [4]. Comparative research on alternative antimicrobial agents with greater efficacy is needed to counteract resistant pathogens that are continually selected by current therapeutic regimen [5]. Traditional medication from plant extracts have been in use for years to cure many ailments, thus, herbal are alternative and relatively safer sources of treatment [6]. Traditional medicinal herbs contain bioactive compounds which have few side effects, have low propensity to develop resistance, are less toxic and are associated with improved efficacy [7]. Recent advances in research involving traditional herbal medicine have resulted in identification of several bioactive compounds with antimalarial, anti-cancer and antibiotic properties [8]. Important antimicrobial agents of plant include alkaloids, phenylpropanoids i.e., flavonoids and terpenoids that include saponins [9] [10].

Kenya has a wide range of flora with over 7,000 plant species [11]. Up to 70% of the rural populace use home remedies from plant parts as the first source of medicine to treat infections [12] [13]. Different plant parts that include flowers, leaves and fruits are used as sources of home remedies [14]. The current study evaluated the antimicrobial activity of *Carissa edulis* roots, *Aspilia pluriseta* leaves, *Vangueria infausta* leaves and *Erythrina abyssinnica* stem that are traditionally used in treatment of gastrointestinal infections in Tharaka-Nithi County. The health benefits of plants mentioned above are attributed to their richness in phytochemical compounds with physiological effects against gastroenteritis bacteria [15]. However, the phytochemicals present in these crude extracts including those used by herbalists in Tharaka-Nithi County is not fully determined.

Solvents that include water, acetone, hexane, methanol and ethanol, are used to extract bioactive compounds from plant materials [16] [8]. Studies has indicated that solvent used in extraction significantly affects extract's yield, quality, extraction velocity, and bioactivity which is attributed to differences in polarity [16] [17] [18]. Use of ethanol has become popular because beside being safe for infusing food items, it does not corrode container used in addition to being easily recoverable with consistent outcome in bioassays involving crude extract [19]. Some of the active metabolites that have been extracted using ethanol as solvent are; alkaloids, glycosides, flavonoids, terpenoids, steroids, tannins, saponins, and reducing sugars [20] [19]. Efficiency of alcohol has been attributed to its polarity that resulted in extraction of more active ingredients [21] [20]. As reported by Abarca-Vargas *et al.* [22] and Hikmawanti *et al.* [20], ethanol is considered a protic organic solvent that has a polarity index value of 5.2 and with the dielectric constant of 24.55. Methanol as an extraction solvent has resulted in amino acid, carbohydrate, flavonoids, phenolic compounds, phytosterol, proteins, saponins and tannin [23] [24]. Hexanoic plant crude extracts have been reported not to contain Phenol, Saponins and Carbohydrates [25]. The levels of phenolics and flavonoids is determined by the solvent and has been reported (butanol > methanol > ethyl acetate > hexane > chloroform [24]).

Understanding of the medicinal herbs' aspects such as phytochemical, pharmacological and standardization is necessary the for constitution of dosage in an informed manner [11]. It is therefore necessary to evaluate different solvent concentrations that will produce the most efficacious extract against the test pathogens. In this respect, this study evaluated effect of different organic solvents (Methanol, hexane, and acetone) on the efficacy of crude extract from five plant species (*Carissa edulis* roots, *Aspilia pluriseta* leaves, *Ficus sycomorus* stems, *Erythrina abyssinnica* barks and *Vangueria infausta* leaves) against human pathogenic bacteria (*Vibrio cholerae*, *Salmonella typhi* and *Shigella flexneri*) associated with gastrointestinal infections.

II. MATERIALS AND METHODS

2.1 Study Area

The plants used in this study were collected in Tharaka-Nithi County in Kenya which is located at a latitude coordinate of 0°9'25.03''S and longitude coordinates of 37°58'41.48''E. Tharaka-Nithi County has four sub counties; Maara, Tharaka North, Tharaka South and Meru South, covering a total area of 2,662.1 km². The human population in Tharaka-Nithi County is estimated at 365,330 [26]

2.2 Experimental Design

Bioassays to analyze the efficacy of the plant crude extracts against the microbes was carried out in 3×3×3×5 factorial experiment laid out in completely randomized design (CRD) having three factors, with factor A being the microbial pathogens (At three levels-*Vibrio cholerae*, *Salmonella typhi* and *Shigella* species). Factor B was the concentrations of crude plant extracts at three levels (1000ppm, 500ppm, 100ppm). Factor C was solvent at three levels (Hexane, acetone and methanol). Lastly factor D was plant extract at 5 levels (*Erythrinna abyssinnica*, *Aspilia pluriseta*, *Vangueria infausta*, *Ficus sycomonus*, *Carissa edulis*).

2.3 Collection of plant samples, Extraction of crude extracts and Phytochemical Analysis

The commonly used herbs were collected with assistance from traditional herbal dispensers in Tharaka North based on the information gathered through questionnaires on the herbs [27]. Information gathered included the part of the plant used, method of extraction and dose. The plant samples were wrapped in aluminum foils and put in Ziplock bags and transported to Chuka university chemistry laboratories for extraction of the metabolites. In the laboratory, the samples were cleaned using running water and rinsed by use of distilled water. They were then chopped in to small sizes using a sterilized panga after which they were left on the bench so as to dry for 21 days. The dried samples were then blended in to soft powder. Extraction of crude extract was performed using methanol, acetone and hexane solvents. 100 grams of each ground sample was weighed and dissolved in 350 ml of methanol, acetone and hexane. They were then allowed to stand for 48 hours so that extraction can take place. Thereafter, filtration was done by use of filter papers number 1. The extracts of the three solvents were then concentrated by use of Rotary evaporator at 40 degrees Celsius and were left to dry in the fume chamber. The concentrated extracts were then weighed using a beam balance and wrapped in aluminum foils, placed in stoppered containers and then transported to KEMRI CDC Kisumu for antimicrobial analysis.

Each plant sample was analyzed for presence of tannins, flavonoids, alkaloids, cardiac glycosides, phenolics, saponins, steroids and terpenoids. Qualitative Test for tannins was done according to Ejikeme *et al.* [28]. Five grams of individual plant powder sample was dissolved in five ml of methanol and then filtered. To the filtrate, 3ml of ferric chloride was added. The occurrence of brownish-green or a blue-black color indicated positive test. Test for flavonoids was done following procedure used by Hossain *et al.* [29] where 2% of the sodium hydroxide was dispensed to each plant sample and followed by addition of dilute hydrochloric acid. The presence of a golden yellow precipitate showed a positive test. Test for saponins was done according to Ejikeme *et al.* [28] where distilled water and each sample filtrate was vigorously shaken till a stable persistent froth was formed. To the mixture, drops of olive oil was added. Formation of an emulsion upon addition of olive oil showed a positive test. Salkowski test as explained by Yadav and Agarwala [30] was used to test for terpenoids presence where three milliliters of each plant extract were dissolved in three milliliters of chloroform. Three milliliters of concentrated sulphuric acid were added where a grayish colour indicated a positive test.

Wagner test procedure used by Ghosh *et al.* [31] was adopted to test for the presence of alkaloids where two milliliters of plant extract was dissolved in two milliliters of ethanol. Two drops of 1M Hydrochloric acid were added to each test tube and boiled for a minute. Three drops of Mayors reagent were added. Formation of a reddish -brown precipitate indicated a positive test. Ferric chloride test as described by Kumar *et al.* [32] was used to test for the presence of phenolics where one milliliter of ferric chloride solution was added to three milliliters of each plant extract sample. Formation of a green colour indicated a positive test for phenolics. Steroids were tested by adding two milliliters of acetic anhydride were added to two milliliters of each plant extract sample followed by

addition of two milliliters of sulphuric acid. Formation of a blue or green colour indicated a positive test for steroids. Kellar-Kiliani test as described by Parekh and Chanda 2007 was used to test for glycosides where two milliliters of glacial acetic acid were added to five milliliters of each extract. One milliliter of ferric chloride was then added followed by addition of one milliliter of concentrated sulphuric acid. A green blue coloration of the solution indicated a positive test for glycosides.

2.4 Antimicrobial activity of the Plant Crude Extracts against test bacteria organisms

a. Preparation of the Crude extract, Culture Media and Microbial Culture Inoculant

Different concentrations of each plant extracts were made dissolving 0.1grams of each crude plant extract to 1 ml 0.2% DMSO making a concentration of 1 g/ml [33]. A 1000 ppm solution was prepared by 1ml of the stock solution into 1000ml of distilled water from which 500ppm and 100ppm concentrations was prepared. The TSB was prepared using a procedure described in the literature [34] where 15 grams of TSB powder was suspended in 500ml of distilled water and then mixed thoroughly. It was slightly warmed to completely dissolve and then autoclaved at 121°C for 15 minutes prior to cooling at room temperature.

b. Minimum Inhibitory Concentration and Minimum bactericidal Concentration of the Extracts

Pure colonies of type cultures of the microorganisms were obtained from KEMRI CDC Kisumu and was sub cultured on Tryptic Soy Broth as per the Clinical Laboratory Standards of US [35]. *Shigella* species which is a member of Enterobacteriaceae family was incubated at a temperature of 35 °C in air for 18 hours while *Vibrio cholerae* and *Salmonella typhi* which are anaerobes was incubated at 10% carbon (IV) oxide, 10% hydrogen and 80% nitrogen. Standardization of bacterial colony cells was done using 0.5 McFarland standard. The McFarland standard was prepared from 0.5ml of 0.048M BaCl₂ (1.17% w/v) added to 99.5ml of 0.18M sulphuric acid (1% v/v) and stirred [36]. The optical density of the turbidity was measured using a spectrophotometer [37] to obtain the final inoculum concentration of bacterial colony forming units (cfu)/ml. The optical density of the colonies was evaluated using a spectrophotometer and adjusted to equal the 0.5 McFarland standard (10⁸ CFU/ml) by adding sterile distilled water.

Inoculations was done by adding 20 µL of each bacterial suspensions to individual 100 µL of TSB in a microtiter plate where each of the treatment was replicated three times. The TSB + DMSO was used while TSB alone as negative control. At the end of the incubation period, A 100 µl of the bacteria under different treatments were cultured by spread plate method on the nutrient agar and was incubated for up to 18 hours. At the end of the re-incubation session, each of the plates was divided into four equal parts using a maker pen and counting of the bacteria CFUs was done for every treatment. The extract concentration at which the CFUs counted was < 10 colonies was assumed to be MBC [38].

2.5 Data analysis

Data obtained on the bacteria colony forming units (Counts) was analyzed using Kruskal Wallis procedure method $\alpha = 0.05$ in Statistical Analysis System (SAS) version 9.4. The medians from the

results were compared by ranking (Wilcoxon rank sum test) in SAS. Analysis only compared plant extracts and excluded results for positive and negative control set ups.

III. RESULTS

3.1 Phytochemical Screening of Extracts from Herbal Plants to Treat Gastrointestinal Infections in Tharaka Nithi County

Methanol as solvent produced higher quantities of crude extract for all the herbal samples used. *Vangueria infausta* produced the highest crude extract with methanol (5.06g) while *Aspilia pluriseta* produced lower yield (0.73g). *Carissa edulis* produced the highest crude extract with acetone (3.7g) while *Aspilia pluriseta* produced lower yield (0.53g). *Carissa edulis* produced the highest crude extract with hexane (1.81g) while *Erythrina infausta* produced lower yield (0.06g). The masses of crude extracts obtained are presented in (Table 1).

Table 1: Mass of crude extracts obtained using methanol, acetone and hexane solvents

Plant sample	Hexane	Acetone	Methanol
<i>Carissa edulis</i>	1.81g	3.7g	4.02g
<i>Aspilia pluriseta</i>	0.45g	0.6g	0.73g
<i>Physalis peruviana</i>	1.25g	2.4g	3.74g
<i>Vangueria infausta</i>	1.32g	1.13g	5.06g
<i>Erythrina abyssinnicca</i>	0.06g	1.61g	1.76g
<i>Ficus sycomorus</i>	0.5g	1.22g	1.6g

Several phytochemical compounds were identified from the herbs (Table 2). In methanolic extracts all the seven herbs contained tannins, saponins and phenolic compounds. Flavonoids were only absent in *Ficus sycomorus*, terpenoids were absent only in *Aspilia pluriseta* and *Vangueria infausta* leaves. Alkaloids were present in *Carissa edulis*, *Vangueria infausta* leaves and barks. Steroids were present in *Ficus sycomorus*, *Carissa edulis*, *Erythrina abyssinnicca* barks only while *Vangueria infausta* leaves were the only containing glycosides.

Table 2: Phytochemicals present in crude extracts of some medicinal plants used for treatment of gastrointestinal infections

Sample	Tnns ^a	Flv ^b	Spnn ^c	Tpn ^d	Alk ^e	Phen ^f	Ster ^g	Glyc ^h
methanolic extracts								
<i>Fs</i> ¹	+	-	+	+	-	+	+	-
<i>Ce</i> ²	+	+	+	+	+	+	+	-
<i>Ap</i> ³	+	-	+	-	-	+	-	-
<i>Ea</i> ⁴	+	+	+	+	-	+	+	-
<i>Vil</i> ⁵	+	+	+	-	+	+	-	+
Hexanoic extracts								
<i>Fs</i> ¹	+	-	+	-	-	+	-	-
<i>Ce</i> ²	+	+	+	+	-	-	+	-
<i>Ap</i> ³	+	-	-	-	-	+	-	-
<i>Ea</i> ⁴	+	+	+	-	-	+	+	-
<i>Vil</i> ⁵	+	+	+	-	-	-	-	+
Acetonic extracts								
<i>Fs</i> ¹	+	-	+	-	-	-	-	-
<i>Ce</i> ²	+	+	+	-	-	-	+	-
<i>Ap</i> ³	+	-	-	-	-	-	-	-

Ea ⁴	+	+	+	-	-	+	-	-
Vil ⁶	+	+	-	-	+	-	-	+

¹F. sycomorus, ²C. edulis, ³A. pluriseta, ⁴E. abyssinnica (barks). ⁵V. infausta (leaves). ^aTannins, ^bFlavonoids, ^cSaponins, ^dTerpenoids, ^eAlkaloids, ^fPhenolics, ^gSteroids, ^hGlycosides

3.2 Effect of Solvents Choice and Plant on Efficacy of Plant's Bacterial Colony Forming Units

The solvent choice significantly ($p < 0.05$) affected the bacteria colony forming units formed by *E. coli*, *Vibrio cholerae* and *Shigella flexneri* subjected to *Erythrina abyssinnica* extracts. The highest bacteria cfu (Median = 217 cfu) and reduced bacteria cfu (Median = 0 cfu) were observed in *E. coli* hexanoic and Methanoic extract respectively when bacteria were subjected to *Erythrina abyssinnica* extracts (Table 3). The solvent choice significantly ($p < 0.05$) affected the bacteria colony forming units formed by *E. coli*, *Salmonella typhi*, *Vibrio cholerae* and *Shigella flexneri* subjected to *Aspilia pluriseta* extracts. The highest bacteria cfu (Median = 276 cfu) and reduced bacteria cfu (Median = 1 cfu) were observed in *E. coli* and *Vibrio cholerae* in hexanoic and acetonic extracts respectively when bacteria were subjected to *Aspilia pluriseta* extracts (Table 3). The solvent choice significantly ($p < 0.05$) affected the bacteria colony forming units formed by *E. coli*, *Salmonella typhi*, *Vibrio cholerae* and *Shigella flexneri* subjected to *Vangueria infausta* extracts. The highest bacteria cfu (Median = 278 cfu) and a reduced bacteria cfu (Median = 17 cfu) were observed in *Shigella flexneri* and *Salmonella typhi* in hexanoic and acetonic extract respectively when bacteria were subjected to *Vangueria infausta* extracts (Table 3). The solvent choice significantly ($p < 0.05$) affected the bacteria colony forming units formed by *E. coli*, *Salmonella typhi*, *Vibrio cholerae* and *Shigella flexneri* subjected to *Ficus sycomorus* extracts. The highest (Median = 140 cfu) was observed in *Salmonella typhi* in hexanoic extract while a reduced cfu (Median = 10 cfu) were observed in *E. coli* in both hexanoic and acetonic extract when bacteria were subjected to *Ficus sycomorus* extracts (Table 3). The solvent choice significantly ($p < 0.05$) affected the bacteria colony forming units formed by *E. coli*, *Salmonella typhi*, *Vibrio cholerae* and *Shigella flexneri* subjected to *carissa edulis* extracts. The highest bacteria cfu was observed in *Salmonella typhi* and *Shigella flexneri* (Median = 198 cfu) in hexanoic extract. Likewise, a reduced cfu (Median = 65 cfu) was observed in *E. coli* in hexanoic extract when bacteria were subjected to *carissa edulis* extracts (Table 3).

Table 3: Effect of different solvents on efficacy plant's extracts against bacteria growth

Plant Extract	Test Bacteria	Hex ^b	Ace ^f	Met ^g	Kruskal-Wallis statistics			
					N	H	df	P value
Eryth ¹	E ^k	217 ^a	200 ^a	0 ^b	9	19.376	2	<.0001
	S ^t	187	177	167	9	2.084	2	0.352
	V ^l	211 ^a	6 ^c	154 ^b	9	19.323	2	<.0001
	S ^r	199 ^a	56 ^b	146 ^a	9	19.142	2	<.0001
Aspi ²	E ^k	276 ^a	232 ^a	3 ^b	9	17.801	2	0.0001
	S ^t	178 ^a	165 ^a	146 ^b	9	15.737	2	0.0004
	V ^l	200 ^a	1 ^c	7 ^b	9	20.544	2	<.0001
	S ^r	222 ^a	166 ^b	28 ^b	9	13.302	2	0.0013
Vang ³	E ^k	250 ^c	243 ^b	187 ^a	9	19.441	2	<.0001
	S ^t	256 ^a	234 ^b	17 ^c	9	20.538	2	<.0001
	V ^l	265 ^a	213 ^b	187 ^b	9	19.413	2	<.0001
	S ^r	278 ^a	220 ^b	187 ^b	9	20.711	2	<.0001
Ficus ⁴	E ^k	55 ^a	10 ^b	10 ^b	9	12.622	2	0.0018
	S ^t	140	123	111	9	6.091	2	0.05
	V ^l	87	109	30	9	2.309	2	0.3153

	S ^t	79 ^b	111 ^a	27 ^b	9	12.169	2	0.0023
Cari ^b	E ^k	65 ^c	155 ^a	143 ^b	9	16.414	2	0.0002
	S ^t	198 ^a	168 ^b	148 ^c	9	18.618	2	<.0001
	V ^l	30 ^b	176 ^a	148 ^a	9	6.909	2	0.0316
	S ^t	198 ^a	156 ^b	140 ^b	9	12.384	2	0.002

^aMedian values followed by the same letters in rows are not significantly different at $\alpha = 0.05$. ¹ *Erythrina abyssinica*, ² *Aspilia pluriseta*, ³ *Vangueria infausta*, ⁴ *Ficus sycomorus*, ⁵ *Carissa edulis*, ^k *E. coli*, ^t *S. typhi*, ^l *V. cholerae*, ^l *S. flexneri*, ^b Hexane extract, ⁷ Acetonic extract, ⁸ Methanolic extract

There was a significant ($p < 0.05$) effect of exposing test bacteria to various plant extracts at different concentration on the bacteria colony forming units formed (Table 4). Exposing *E. coli* to 1000ppm plant extracts resulted in significantly reduced bacteria cfu counts in methanolic extracts of *Erythrina abyssinica* (Median = 0 cfu), *Aspilia pluriseta* (Median = 3 cfu) and *Ficus sycomorus* (Median = 0 cfu) while remaining significantly higher in hexanoic extract of *Aspilia pluriseta* (Median = 276 cfu) and *Vangueria infausta* (Median = 250 cfu). Exposing *S. typhi* to 1000ppm of plant extracts resulted in significantly lower bacteria cfu counts in methanolic extracts of *Ficus sycomorus* (Median = 13 cfu) and *Vangueria infausta* (Median = 17 cfu) while remaining significantly higher in hexanoic (Median = 256 cfu) and acetonic (Median = 256 cfu) extracts of *Vangueria infausta* respectively. Exposing *V. cholerae* to 1000ppm of plant extracts resulted in significantly lower bacteria cfu counts in acetonic extracts of *Erythrina abyssinica* (Median = 0 cfu) and *Aspilia pluriseta* (Median = 1 cfu) while significantly higher was observed in hexanoic (Median = 265 cfu) extract of *Vangueria infausta*. Exposing *Shigella flexneri* to 1000ppm of plant extracts resulted in significantly lower bacteria cfu counts in acetonic extracts of *Erythrina abyssinica* (Median = 2 cfu) and methanolic extract of *Ficus sycomorus* (Median = 1 cfu) and significantly higher cfu was observed in hexanoic (Median = 278 cfu) extract of *Vangueria infausta* (Table 4).

Exposing *E. coli* to 500ppm plant extracts resulted in significantly reduced bacteria cfu counts in methanolic extracts of *Erythrina abyssinica* (Median = 0 cfu) and *Aspilia pluriseta* (Median = 3 cfu) but remained significantly higher in hexanoic extract of *Vangueria infausta* (Median = 258 cfu). Exposure of *S. typhi* to 1000ppm of plant extracts resulted in significantly reduced bacteria cfu counts in methanolic extracts of *Vangueria infausta* (Median = 62 cfu) but remaining significantly higher in hexanoic (Median = 266 cfu) extracts of *Vangueria infausta*. Exposing *V. cholerae* to 500ppm of plant extracts resulted in significantly reduced cfu counts in acetonic extracts of *Ficus sycomorus* (Median = 4 cfu) and *Aspilia pluriseta* (Median = 5 cfu) while bacteria cfu counts were significantly higher in hexanoic (Median = 287 cfu) extract of *Vangueria infausta*. Exposing *Shigella flexneri* to 500ppm of plant extracts resulted in reduced bacteria cfu counts in methanolic extracts of *Carissa edulis* (Median = 27 cfu) and higher cfu was observed in acetonic (Median = 265 cfu) extract of *Aspilia pluriseta* (Table 4).

When *E. coli* was exposed to 100ppm plant extracts, significantly reduced bacteria cfu counts occurred in methanolic extracts of *Aspilia pluriseta* (Median = 0 cfu) but remained significantly higher in hexanoic extract of *Vangueria infausta* (Median = 296 cfu). Exposure of *S. typhi* to 100ppm of plant extracts resulted in significantly reduced cfu counts in acetonic extracts of *Erythrina abyssinica* (Median = 22 cfu) but remaining significantly higher (Median = 284 cfu) in hexanoic extracts of

Vangueria infausta. Exposing *V. cholerae* to 100ppm of plant extracts resulted in significantly reduced cfu counts in acetonic extracts of *Aspilia pluriseta* (Median = 9 cfu) while significantly higher in hexanoic (Median = 265 cfu) extract of *Vangueria infausta*. Exposing *Shigella flexneri* to 100ppm of plant extracts resulted in reduced cfu counts in acetonic extracts of *Vangueria infausta* (Median = 22 cfu) and higher bacteria cfu was observed in hexanoic (Median = 288 cfu) extract of *Vangueria infausta* (Table 4).

Table 4: Effect of different solvents and extract concentrations on efficacy of plant's extracts against bacteria growth

Plant Extract	Sol	N	Plant extracts at 1000ppm					Kruskal-Wallis statistics		
			Eryth ¹	Aspi ²	Vang ³	Fic ⁴	Cari ⁵	H	Df	p-value
<i>E^k</i>	Hex ⁶	3	217 ^b	276 ^a	250 ^a	57 ^c	21 ^c	12.433	4	0.014
	Ace ⁷	3	154 ^b	232 ^a	243 ^a	11 ^c	165 ^b	11.750	4	0.019
	Met ⁸	3	0 ^c	3 ^c	187 ^a	3 ^c	80 ^b	11.835	4	0.018
<i>S^t</i>	Hex ⁶	3	163 ^b	178 ^b	256 ^a	67 ^c	189 ^b	11.145	4	0.025
	Ace ⁷	3	22 ^d	165 ^b	234 ^a	122 ^c	168 ^b	12.833	4	0.012
	Met ⁸	3	167 ^a	146 ^a	17 ^c	13 ^b	145 ^a	11.212	4	0.024
<i>V^l</i>	Hex ⁶	3	144 ^c	200 ^b	265 ^a	57 ^d	8	13.524	4	0.009
	Ace ⁷	3	0 ^d	1 ^d	213 ^a	122 ^c	176 ^b	13.099	4	0.011
	Met ⁸	3	164 ^b	7 ^c	220 ^a	7 ^c	38 ^c	12.077	4	0.016
<i>S^t</i>	Hex ⁶	3	144 ^c	222 ^b	278 ^a	28 ^d	200 ^b	13.257	4	0.010
	Ace ⁷	3	2 ^d	166 ^b	202 ^a	110 ^c	20 ^d	12.9	4	0.011
	Met ⁸	3	109 ^b	28 ^d	187 ^a	4 ^d	89 ^c	12.923	4	0.012
Plant extracts at 500ppm										
<i>E^k</i>	Hex ⁶	3	215 ^b	234 ^a	258 ^a	47 ^c	65 ^c	11.479	4	0.022
	Ace ⁷	3	200 ^a	188 ^a	222 ^a	11 ^d	167 ^c	10.444	4	0.034
	Met ⁸	3	0 ^c	0 ^c	198 ^a	10 ^c	143 ^b	13.68	4	0.008
<i>S^t</i>	Hex ⁶	3	187 ^b	233 ^a	266 ^a	140 ^c	203 ^b	11.979	4	0.017
	Ace ⁷	3	177 ^a	200 ^a	195 ^a	133 ^b	177 ^a	10.627	4	0.031
	Met ⁸	3	133 ^a	117 ^a	62 ^c	111 ^b	136 ^a	10.961	4	0.027
<i>V^l</i>	Hex ⁶	3	234 ^b	225 ^b	287 ^a	89 ^c	30 ^d	11.979	4	0.017
	Ace ⁷	3	6 ^b	5 ^b	189 ^a	4 ^b	176 ^a	10.627	4	0.031
	Met ⁸	3	123 ^b	132 ^a	152 ^a	89 ^c	148 ^a	10.961	4	0.027
<i>S^t</i>	Hex ⁶	3	238 ^b	223 ^b	278 ^a	80 ^d	185 ^c	10.252	4	0.036
	Ace ⁷	3	41 ^c	265 ^a	201 ^b	132 ^c	172 ^b	12.533	4	0.013
	Met ⁸	3	146	154	142	27	130	8	4	0.092
Plant extracts at 100ppm										
<i>E^k</i>	Hex ⁶	3	258 ^a	276 ^a	296 ^a	64 ^b	74 ^b	11.487	4	0.022
	Ace ⁷	3	217 ^b	283 ^a	208 ^b	6 ^d	99 ^c	12.833	4	0.012
	Met ⁸	3	3 ^d	0 ^d	87 ^b	38 ^c	159 ^a	13.597	4	0.008
<i>S^t</i>	Hex ⁶	3	220 ^b	266 ^a	284 ^a	160 ^c	189 ^b	12.022	4	0.017
	Ace ⁷	3	22 ^d	165 ^b	234 ^a	122 ^c	168 ^b	12.833	4	0.012
	Met ⁸	3	204 ^b	267 ^a	202 ^b	115 ^d	166 ^c	12.433	4	0.014
<i>V^l</i>	Hex ⁶	3	211 ^b	265 ^a	89 ^c	188 ^b	8 ^d	13.233	4	0.010
	Ace ⁷	3	76 ^d	9 ^e	243 ^a	113 ^c	178 ^b	13.548	4	0.008
	Met ⁸	3	193	154	162	27	171	8.956	4	0.062
<i>S^t</i>	Hex ⁶	3	267 ^a	279 ^a	288 ^a	103 ^c	198 ^b	11.215	4	0.024
	Ace ⁷	3	84 ^c	76 ^c	22 ^d	110 ^b	156 ^a	12.9	4	0.012
	Met ⁸	3	189 ^a	167 ^a	181 ^a	120 ^b	165 ^a	12.714	4	0.013

^aMedian values followed by the same letters in rows are not significantly different at $\alpha = 0.05$. ¹ *Erythrina abyssinica*, ² *Aspilia pluriseta*, ³ *Vangueria infausta*, ⁴ *Ficus sycomonus*, ⁵ *Carissa edulis*, ^k *E. coli*, ^t *S. typhi*, ^l *V. cholerae*, ^s *S. flexneri*, ⁶ Hexane extract, ⁷ Acetonioic extract, ⁸ Methanoic extract

IV. DISCUSSION

4.1 Phytochemical Extraction Using Different Solvent and Analysis of Crude Extracts

A higher yield is observed in methanol then acetone and then in hexane extracts. These observations are comparable to other results obtained in related studies such as those of Do *et al* (2014) on the extraction yield of *Limophila aromatica* and that of Kuppusamy *et al.* [39] on *Commelina nudiflora*. Further, these results are supported by studies which have demonstrated that the efficiency of extraction process is dependent on type of the solvent used [40] [41]. Variation on the amount of extract obtained by different solvent is attributed to difference in their polarity [42]. This could be that the medicinal herbs contain a higher number of polar compounds. High polar compounds are soluble in solvents that have high polarity like in the case of methanol [42]. Liu *et al.* [43] describes methanol, a polar solvent as a solvent with improved efficiency of solvation as a result of interaction of hydrogen bonds of the polar sites in the metabolites and the solvent than the non-polar.

The phytochemical analysis results showed the occurrence of secondary metabolites such as phenols, saponins, tannins, flavonoids, terpenoids and glycosides which are known for medicinal attributes as well as physiological activities of the herb [44]. These phytochemicals may be produced in response to inversion of the plant by the microbes. The phytochemicals have been observed to have antimicrobial properties against numerous micro-organisms [45] [46]. Extracts from *Carissa edulis* had compounds such as terpenoids, tannins, steroids flavonoids, saponins, alkaloids, phenolics and absence of glycosides. The current results are in line other related studies such as those of Nawaz [47] who reported occurrence of such compounds in stem bark of *Ficus sycomorus*. The results are further in agreement with a study carried out by Madivoli *et al.* [48].

Extracts from *Ficus sycomorus* had tannins, saponins, terpenoids, phenolics, steroids while flavonoids, alkaloids and glycosides were absent. The occurrence of these metabolites was similar to a study by Nawaz *et al.* [47] who observed tannins, flavonoids, saponins and phenolics. This was also in agreement with a study carried out in Sudan by Osama and Awdekarim [49]. However, in their study flavonoids and alkaloids were present, this difference could be linked to associated to the difference in climatic conditions or even the soil type in which this herb grow from where the samples were collected.

Aspilia pluriseta extracts had tannins, saponins and phenolics while flavonoids, terpenoids, alkaloids, steroids and glycosides were absent. This was similar to a study carried out by Njeru and Muema [50] in Mbeere community, Embu County which is used in treatment of tuberculosis. The results of this study differed with those carried out around Maseno University, Kenya by Musyimi *et al.* [51] who found the presence of alkaloids, steroids and glycosides which were absent in this study. In both studies methanol as a solvent was used for extraction, therefore the difference could be attributed to different rainfall amount or even the soil type in these different regions. *Erythrina abyssinnica* extracts had tannins, flavonoids, saponins, terpenoids, phenolics, steroids while alkaloids and glycosides were absent. These results are supported by a similar study conducted in Tharaka-Nithi county on utilization of herbal medicine among children under five by Nzuki [52] who identified presence of tannins, terpenoids, alkaloids, phenolics and flavonoids.

Vangueria infausta leaf extract had tannins, flavonoids, saponins, alkaloids, phenolics and glycosides while terpenoids and steroids were absent. These results agree with report of Obakiro *et al.* [53] who observed the presence of flavonoids, alkaloids, phenolics and saponins. These results also are in line with results from a study carried in Zimbabwe by Manyarara *et al.* [54] where the extract is used in treatment of candidiasis. Saponins are present in almost all medicinal herbs which grounds for their curative potential. Biological role of phytochemicals in plants is solely to protect them against invasion by injurious pathogens and organisms. That notwithstanding, certain saponins have been reported to be toxic to endotherms but such toxicity is lower towards mammals. Since they are toxic to quite a number of organisms, saponins find their use in pharmacological, antibiotic and fungicidal applications [55]. Flavonoids which are based on a 15-carbon skeleton is comprised of two benzene ring connected through a heterocyclic pyrane ring. The basic flavonoid structure is the aglycone. These flavonoids have many biochemical properties including antioxidant property which is governed by function group arrangement. The substitution, configuration and the total number of hydroxyl groups determines the antioxidant property mechanisms such as the ability to chelate metal ions and in scavenging of radical [56] [57]. The mechanism of antioxidant action involves upregulation or protection of antioxidant defences, suppression of injurious reactive oxygen species which is formed by either inhibition of the enzymes or through chelation of the trace elements that are involved in free radical generation and also by scavenging reactive oxygen species [46].

Studies by Meng *et al.* [58] as well as that by Kumar and Pandey [59] show that flavonoids are present in herbal medicines and are reported to be effective as cardio protective agents, antibacterial, antioxidants and an important part of pharmaceutical and medical applications. In plants, the living and non-living factors bring about production of ROS leading to oxidative stress [59]. The production of flavonoids is improved as a result of this oxidative stress and they exhibit the ability to hinder ROS generation, ability to quench ROS immediately they are formed [59]. Further, they have the potential of absorbing the energetic solar wavelengths. The antioxidant capability as well as the ability of absorbing ultra violet wavelengths depend on how different rings of flavonoids are substituted [59]. According to Cazorolli *et al.* [60], flavonoids inhibit enzymes-based activities including peroxidase and those of xanthine oxidase which take part in generation of free radical thus reducing the oxidative damage brought about by the macromolecules.

The phenolic compounds are among the highest and common plant phytochemicals [61]. Phenolic compounds have anti-inflammation properties, anti-aging properties, and have the advantage of improvement of endothelial function and causing the inhibition of the cell proliferation and angiogenesis activities [62]. Phenolic compounds accounts for the antioxidant properties exhibited by most plants used as herbal medicine [63]. Phenolic compounds in plants comprises of chemicals like flavonoids, tocopherols and phenolic acids [63] and are available in almost all plant parts. Phenolics have roles such as scavenging of free radicals, have anticarcinogenic effect as well as antioxidant properties. Phenolics are useful in prevention of human chronic diseases like those associated with bacterial as well as parasitic infections [64].

Steroids have been reported to be endowed with antibacterial phytochemicals [65]. Further, the steroids are considered to be essential phytochemicals as they are associated with sex hormones [30]. Despite having cytotoxic properties, steroids have medical significance [30]. Reports indicate that alkaloids obtained from plants are used as therapy for anaesthetics, antioxidants, antidiabetic, antimicrobial and muscle relaxants. In plants, alkaloids offer protection against injurious singlet reactive oxygen which may be produced in plant tissues and excited by light source [66]. According to report by Ayoola *et al.* [67], alkaloids possess anti-bacterial properties that are exploitable for treatment of bacterial related diseases. Additionally, alkaloids have anti-plasmodic as well as the analgesic significance [67]. Tannins comprises of complex derivatives of the acid (polyhydroxy benzoic) an organic or non-nitrogenous compound endowed with the ability of precipitating proteins. They are capable of dissolving in water to form colloidal solutions. Further, tannins are soluble in acetone alcohol but are however slightly soluble in organic solvents including chloroform and ethyl acetate. Presence of groups such as carboxylic as well as the free phenolic gives tannins solutions acidic characteristics [68]. Tannins are medicinally used as antidiarrheal, as haemostatic and as antihemorrhoidal metabolites. Due to their anti-inflammatory activities, are useful in the treatment of enteritis, bowel disorders and gastritis. Further, tannins may be used to inhibit viral infections [69], bacterial infection as well as to treat parasitic infections [70]. According to Awuchi [71] glycosides are molecules that have their sugar attached to a different functional group by way of glycosidic bond. Reports by Njoku and Obi [72] indicate that glycosides lower blood pressure.

4.2 Effect of Solvents Choice and Plant on Efficacy of Plant's Bacterial Colony Forming Units

Ficus sycomorus extracts gave the reduced colonies growth of tested bacteria that included *Escherichia coli* (5 cfu). *Vibrio cholerae* 7.08 cfu and *Shigella flexneri* (13.28 cfu). Reduced colony growth of *Salmonella typhi* (28.34 cfu) was observed with crude extract of *Erythrina abyssinnica*. Slightly higher colonies of *Shigella flexneri* and *Vibrio cholera* were observed in the crude extract from *Vangueria infausta* while higher colony units of *Escherichia coli* were observed with the extracts from *Carissa edulis*. According to Korir *et al.* [73]. *Erythrina abyssinnica* extracts are active against gram positive bacteria as well as on the gram-negative bacteria but have lower efficacy against *E. coli*.

Differences in response of different bacteria species towards extracts of some plant species have been reported in other work [74]. Specifically, evaluation of antimicrobial activity of *Ficus* sp extracts inhibit different bacteria species at different levels [75]. Variability of plant extracts in reducing the growth of bacterial colony units in this study could be attributed to plants species with different strength of phytochemicals. This argument is supported by related work of Odunbaku *et al.* [76] and Chilufya *et al.* [77] on the minimal inhibition of sycomorous. According to Mostafa *et al.* [78] inhibition of different plant extracts against microorganisms may vary depending on plant's chemical constituents and volatility of such chemicals.

Environmental and climatical variations have been cited to influence the concentrations of plant's phytochemicals thus affecting plant's antibacterial activity [77]. Antimicrobial constituents in plant extracts include compounds such as alkaloids, flavonoids, terpenoid and phenolics [79]. These antimicrobial contents have the potential to disrupt microbial cells limiting cell functions by inducing

cell death or inhibit biosynthesis of important enzymes and amino acids [80] [79]. Additionally, bioactivity of extracts from the plants could further be attributed to the hydrophobicity attributes enabling extracts to react with cell membrane and mitochondria and altering cells permeability [81] [82]. Variation in colony forming units observed is an indication that different bacteria species differ in their susceptibility to different plant extracts.

4.3 Effect of Varied Concentrations of Crude Plant Extracts on Bacteria Growth

Reduction of bacteria colony forming units in all test bacteria evaluated reduced progressively with increasingly higher concentration of extracts. Minimum inhibition concentration in this study was at 0.5 ppm for all the micro-organisms studied and for all the plants extracts used. The minimum inhibition concentration of *Vangueria infausta* stem bark and crude extracts of *Physalis peruviana* leaves may be higher than 1000ppm and may be the reason why there was no inhibition in all dilutions of the tested gastroenteritis pathogens. In the case of *Aspilia pluriseta*, according to reports, methanolic extracts exhibit broad-spectrum bio-activity against microbes (*Escherichia coli*, *Staphylococcus aureus*) at the concentration of 6.26 – 25 µg/mL [50]. Musa *et al.* [83] indicated a minimal inhibitory concentration of the extracts from *F. sycomorus* bark ethanolic extracts on *S. typhi* and *E. coli* range of 6.25-1.56 mg/ml which were higher than the observed minimal inhibition concentration in this study. The concentration in this study remains lower than 50mg/ml and 300mg/ml documented earlier by Odunbaku *et al.* [76]. Variation in phytochemical activity may have been attributed to variability of environment and climate which have effect on plant phytochemical constituents. According to Li *et al.* [84] phytochemical constituents of same plant species growing under different environmental conditions can be different.

The methanolic extracts from *Erythrina abyssinnica*, *Aspilia pluriseta*, *Ficus sycomorus* performed better against *Escherichia coli*. Better performance of methanolic extract against bacteria species observed in this study are supported by other studies [85] [75] [86]. Manimozhi *et al.* [85] reported that MeOH extract of *Ficus species* exhibited better activity against all bacteria tested that included *E. coli*, *B. subtilis*, and *S. aureus*. Variability of the extracts obtained by different solvent may be attributed to their differences in polarity status. Accordingly, it has been observed that Polarity of solvents has influence on the extraction efficiency of plants phytochemicals [16]. Hence, active phytochemical constituents that may not extract efficiently into acetone and hexane extract, methanol extraction is likely to provide more consistent with wider antimicrobial activity.

References

- [1] A. Degu, P. Njogu, I. Weru and P. Karimi, "Assessment of drug therapy problems among patients with cervical cancer at Kenyatta National Hospital, Kenya," *Gynecologic Oncology Research and Practice*, vol. 4, no. 1, p. 15, 2017.
- [2] A. Alhedethe, K. Alhudaithy and A. Zloh, "An evaluation of prevalence of low quality of medicines in Saudi Arabia and factors associated an analytical comparative study," *Archives in Chemical Research*, vol. 2, no. 1, pp. 1-18, 2017.
- [3] UN, "Transforming our world: The 2030 agenda for sustainable development," New York:

United Nations, Department of Economic and Social Affairs, 2015.

- [4] F. Khuluza, S. Kigera and L. Heide, "Low prevalence of substandard and falsified antimalarial and antibiotic medicines in public and faith-based health facilities of southern Malawi," *The American Journal of Tropical Medicine and Hygiene*, vol. 96, no. 5, pp. 1125 - 1135, 2017.
- [5] WHO, "Prioritization of pathogens to guide discovery, research and development of new antibiotics for drug-resistant bacterial infections, including tuberculosis (No. WHO/EMP/IAU/2017.12)," World Health Organization, 2017.
- [6] C. Nunes, M. Arantes, S. Pereira, L. Cruz, M. Passos and L. Moraes, "Plants as sources of anti-inflammatory agents," *Molecules*, vol. 25, no. 16, p. 3726, 2020.
- [7] M. A. Mir, N. Bashir, A. Alfaify and M. D. Oteef, "GC-MS analysis of Myrtus communis extract and its antibacterial activity against Gram-positive bacteria," *BioMed Central. complementary medicine and therapies*, vol. 20, no. 1, pp. 1-9, 2020.
- [8] P. A. Uwineza and A. Waśkiewicz, "Recent advances in supercritical fluid extraction of natural bioactive compounds from natural plant materials," *Molecules*, vol. 25, no. 17, p. 3847, 2020.
- [9] J. Sampedro and E. R. Valdivia, "New antimicrobial agents of plant origin," in *Antimicrobial Compounds: Current Strategies and New Alternatives*, T. G. Villa and P. Veiga-Crespo, Eds., Heidelberg, Germany, Springer-Verlag, 2014, pp. 83-114.
- [10] A. Altemimi, N. Lakhssassi, A. Baharlouei, D. G. Watson and D. A. Lightfoot, "Phytochemicals: Extraction, isolation, and identification of bioactive compounds from plant extracts," *Plants*, vol. 6, no. 4, p. 42, 2017.
- [11] Y. M. Mbuni, S. Wang, B. N. Mwangi, N. J. Mbari, P. M. Musili, N. O. Walter and Q. Wang, "Medicinal plants and their traditional uses in local communities around Cherangani Hills, Western Kenya," *Plants*, vol. 9, no. 3, p. 331, 2020.
- [12] K. R. Cheruiyot, D. Olila and J. Kateregga, "In-vitro antibacterial activity of selected medicinal plants from Longisa region of Bomet district, Kenya," *African Health Sciences*, vol. 9, no. 2, pp. 42-46, 2009.
- [13] J. K. Muthee, D. W. Gakuya, J. M. Mbaria, P. G. Kareru, C. M. Mulei and F. K. Njonge, "Ethnobotanical study of anthelmintic and other medicinal plants traditionally used in Loitokitok district of Kenya," *Journal of Ethnopharmacology*, vol. 135, no. 1, p. 15, 2011.
- [14] D. W. Gakuya, S. M. Itonga, J. M. Mbaria, J. K. Muthee and J. K. Musau, "Ethnobotanical survey of biopesticides and other medicinal plants traditionally used in Meru central district of Kenya," *Journal of Ethnopharmacology*, vol. 145, no. 2, pp. 547-553, 2013.
- [15] D. W. Gakuya, M. O. Okumu, S. G. Kiama, J. M. Mbaria, P. K. Gathumbi, P. M. Mathiu and J. M. Nguta, "Traditional medicine in Kenya: past and current status, challenges, and the way forward," *Scientific African*, vol. 8, pp. 360-367, 2020.
- [16] A. Wakeel, S. A. Jan, I. Ullah, Z. K. Shinwari and X. Ming, "Solvent polarity mediates phytochemical yield and antioxidant capacity of *Isatis tinctoria*," *PeerJournal*, vol. 7, pp. 7857-7879, 2019.
- [17] K. Rafińska, P. Pomastowski, J. Rudnicka, A. Krakowska, A. Maruśka, M. Narkute and B. Buszewski, "Effect of solvent and extraction technique on composition and biological activity of *Lepidium sativum* extracts," *Food Chemistry*, vol. 289, p. 16–25, 2019.
- [18] H. Zhang, J. Birch, J. Pei, I. A. Mohamed Ahmed, H. Yang, G. Dias and A. E. D. Bekhit, "Identification of Six Phytochemical Compounds from *Asparagus officinalis* L. Root Cultivars from New Zealand and China Using UAE-SPE-UPLC-MS/MS: Effects of Extracts on H₂O₂-Induced Oxidative Stress," *Nutrients*, vol. 11, no. 1, p. 107, 2019.

- [19] L. Sikder, M. Khan, S. Z. Smrity, M. T. Islam and S. A. Khan, "Phytochemical and pharmacological investigation of the ethanol extract of *Byttneria pilosa* Roxb," *Clinical Phytoscience*, vol. 8, no. 1, pp. 1-8, 2022.
- [20] N. P. E. Hikmawanti, S. Fatmawati and A. W. Asri, "The effect of ethanol concentrations as the extraction solvent on antioxidant activity of Katuk (*Sauropus androgynus* (L.) Merr.) leaves extracts," in *In IOP Conference Series: Earth and Environmental Science*, 2021.
- [21] R. Sankeshwari, A. Ankola, K. Bhat and K. Hullatti, "Soxhlet versus cold maceration: Which method gives better antimicrobial activity to licorice extract against *Streptococcus mutans*?," *Journal of the Scientific Society*, vol. 45, no. 2, p. 67, 2018.
- [22] R. Abarca-Vargas, C. F. P. Malacara and V. L. Petricevich, "Characterization of chemical compounds with antioxidant and cytotoxic activities in *bougainvillea x buttiana* holttum and standl, (Var. rose) extracts," *Antioxidants*, vol. 5, no. 45, p. 1–11, 2016.
- [23] B. Pandey, R. Thapa and A. Upreti, "Chemical composition, antioxidant and antibacterial activities of essential oil and methanol extract of *Artemisia vulgaris* and *Gaultheria fragrantissima* collected from Nepal," *Asian Pacific Journal of Tropical Medicine*, vol. 10, no. 10, p. 952–9, 2017.
- [24] M. D. Shah, J. S. S. Seelan and M. Iqbal, "Phytochemical investigation and antioxidant activities of methanol extract, methanol fractions and essential oil of *Dillenia suffruticosa* leaves," *Arabian Journal of Chemistry*, vol. 13, no. 9, pp. 7170-7182, 2020.
- [25] Y. Isah and M. P. Ibrahim, "Analysis of Phytocomponents in the n-Hexane Extract from the Stem Bark of *Parinari polyandra* (Benth) using FTIR and GC-MS," *Bayero Journal of Pure and Applied Sciences*, vol. 14, no. 2, pp. 15-19, 2021.
- [26] K. Kenya National Bureau of Statistics, "Kenya Population and Housing Census : Population by County and Sub-County," Government of Kenya, Nairobi, 2019.
- [27] M. P. Kiteme, O. B. Onyango, C. E. Njagi and O. F. Ogolla, "Medicinal Plants Used for Treatment of Gastrointestinal Infections in Tharaka-Nithi County," *International Journal of Pathogen Research*, vol. 10, no. 4, pp. 1-11, 2022.
- [28] C. Ejikeme, C. S. Ezeonu and A. N. Eboatu, "Determination of Physical and Phytochemical Constituents of some Tropical Timbers Indigenous to niger delta area of nigeria," *European Scientific Journal*, vol. 10, no. 18, pp. 247-270, 2014.
- [29] M. Hossain, K. AL-Raqmi, Z. Al-Mijizy, A. Weli and Q. Al-Riyami, "Study of total phenol, flavonoids contents and phytochemical screening of various leaves crude extracts of locally grown *Thymus vulgaris*," *Asian Pacific journal of tropical biomedicine*, vol. 3, no. 9, pp. 705-710, 2013.
- [30] R. N. S. Yadav and M. Agarwala, "Phytochemical analysis of some medicinal plants," *Journal of phytology*, vol. 3, no. 12, pp. 10-14, 2011.
- [31] A. Ghosh, S. De and Y. N. Dey, "Phytochemical investigation and chromatographic evaluation of the different extracts of tuber of *Amorphaphallus paeoniifolius* (Araceae)," *International Journal on Pharmaceutical and Biomedical Research*, vol. 1, no. 5, pp. 150-1, 2010.
- [32] P. P. Kumar, S. Kumaravel and C. Lalitha, "Screening of antioxidant activity, total phenolics and GC-MS study of *Vitex negundo*," *African Journal of Biochemistry Research*, vol. 4, no. 7, pp. 191-195, 2010.
- [33] D. Konkolewicz, A. J. Magenau, S. E. Averick, A. Simakova, H. He and K. Matyjaszewski, "ICAR ATRP with ppm Cu Catalyst in Water," *Macromolecules*, vol. 45, no. 11, pp. 4461-4468, 2012.
- [34] M. Lalitha, *Manual on Antimicrobial Susceptibility Testing*, Washington DC, USA: American Society for Microbiology, 2009.

- [35] S. Puttaswamy, S. K. Gupta, H. Regunath, L. P. Smith and S. Sengupta, "A comprehensive review of the present and future antibiotic susceptibility Testing (AST) systems," *Archives of Clinical Microbiology*, vol. 9, no. 3, pp. 83-92, 2018.
- [36] J. M. Andrews, "Determination of minimum inhibitory concentrations," *Journal of antimicrobial Chemotherapy*, vol. 48, no. 1, pp. 5-16, 2001.
- [37] I. Wiegand, K. Hilpert and R. E. Hancock, "Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances," *Nature Protocols*, vol. 3, no. 2, pp. 163-175, 2008.
- [38] R. Mogana, A. Adhikari, M. N. Tzar, R. Ramliza and C. Wiart, "Antibacterial activities of the extracts, fractions and isolated compounds from *Canarium patentinervium* Miq. against bacterial clinical isolates. BMC Complementary Medicine and Therap," *BMC Complementary Medicine and Therapies*, vol. 20, no. 1, pp. 1-11, 2020.
- [39] P. Kuppusamy, M. M. Yusoff, N. R. Parine and N. Govindan, "Evaluation of in-vitro antioxidant and antibacterial properties of *Commelina nudiflora* L. extracts prepared by different polar solvents," *Saudi journal of biological sciences*, vol. 22, no. 3, pp. 293-301, 2015.
- [40] N. Turkmen, F. Sari and Y. S. Velioglu, "Effects of extraction solvents on concentration and antioxidant activity of black and black mate tea polyphenols determined by ferrous tartrate and Folin-Ciocalteu methods," *Food chemistry*, vol. 99, no. 4, pp. 835-841, 2006.
- [41] T. V. Ngo, C. J. Scarlett, M. C. Bowyer, P. D. Ngo and Q. V. Vuong, "Impact of different extraction solvents on bioactive compounds and antioxidant capacity from the root of *Salacia chinensis* L.," *Journal of Food Quality*, pp. 1-8, 2017.
- [42] D.-H. Truong, D. H. Nguyen, N. T. A. Ta, A. V. Bui, T. H. Do and C. N. Hoang, "Evaluation of the Use of Different Solvents for Phytochemical Constituents, Antioxidants, and In Vitro Anti-Inflammatory Activities of *Severinia buxifolia*," *Journal of Food Quality*, vol. 2019, 2019.
- [43] X. Liu, M. Dong, X. Chen, M. Jiang, X. Lv and G. Yan, "Antioxidant activity and phenolics of an endophytic *Xylaria* sp. from *Ginkgo biloba*," *Food chemistry*, vol. 105, no. 2, pp. 548-554, 2007.
- [44] M. F. Kadir, M. S. B. Sayeed and M. M. K. Mia, "Ethnopharmacological survey of medicinal plants used by indigenous and tribal people in Rangamati, Bangladesh," *Journal of ethnopharmacology*, vol. 144, no. 3, pp. 627-637, 2012.
- [45] A. Mishra, S. Kumar, A. Bhargava, B. Sharma and A. K. Pandey, "Studies on in vitro antioxidant and antistaphylococcal activities of some important medicinal plants.," *Cellular and Molecular Biology*, vol. 57, no. 1, pp. 16-25, 2011.
- [46] A. Mishra, S. Kumar and A. K. Pandey, "Scientific validation of the medicinal efficacy of *Tinospora cordifolia*," *The Scientific World Journal*, pp. 1-8, 2013.
- [47] H. Nawaz, R. Waheed and M. Nawaz, "Phytochemical composition, antioxidant potential, and medicinal significance of ficus," in *Modern Fruit Industry*, I. Kahramanoglu, A. Küden and S. Çömlekçioglu, Eds., 2019, pp. 553-573.
- [48] E. S. Madivoli, M. K. Murigi, P. G. Patrick G, J. Karanja, K. Cheruiyot, S. O. Rechab and E. G. Maina, "Phytochemical and antimicrobial evaluation of selected medicinal plants in Meru community of Kenya," *Journal of Medicinal Plants for Economic Development*, vol. 2, no. 1, pp. 1-4, 2018.
- [49] A. Osama and S. Awdelkarim, "Phytochemical screening of *Ficus sycomorus* L. bark and *Cleome gynandra* L. aerial parts," *Journal of Pharmacognosy and Phytochemistry*, vol. 4, no. 4, pp. 24-27, 2015.
- [50] S. N. Njerua and J. M. Muema, "Antimicrobial activity, phytochemical characterization and gas chromatography-mass spectrometry analysis of *Aspilia pluriseta* Schweinf. extracts," *Heliyon*,

- vol. 6, no. 10, pp. 5195-5165, 2020.
- [51] D. M. Musyimi, J. A. Ogur and P. M. Muema, "Effects of leaf and root extracts of *Aspilia* plant (*Aspilia mossambicensis*) (Oliv) Wild. on some selected microorganisms," *International Journal of Biological Chemistry*, vol. 1, no. 4, pp. 213-220, 2007.
- [52] D. M. Nzuki, "Utilization of herbal medicine among children under 5 years of age in Tharaka Nithi County, Kenya," *Doctoral dissertation, Kenyatta university, Kenya*, 2016.
- [53] S. B. Obakiro, A. Kiprop, E. Kigundu, I. K'Owino, M. P. Odero, S. Manyim and L. Bunalema, "Traditional medicinal Uses, phytoconstituents, bioactivities, and toxicities of *Erythrina abyssinica* Lam. ex DC. (fabaceae): a systematic review," *Evidence-Based Complementary and Alternative Medicine*, pp. 1-43, 2021.
- [54] T. E. Manyarara, J. Chifamba and F. T. Tarugarira, "Antifungal activity of *Ziziphus mucronata* and *Erythrina abyssinica* bark crude extracts on *Cryptococcus neoformans* and *Candida albicans* species," *Journal of Pharmaceutical Research International*, vol. 10, no. 3, pp. 1-11, 2016.
- [55] S. D. Desai, D. G. Desai and H. Kaur, "Saponins and their biological activities," *Pharma Times*, vol. 41, no. 3, pp. 13-16, 2009.
- [56] K. E. Heim, A. R. Tagliaferro and D. J. Bobilya, "Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships," *The Journal of nutritional biochemistry*, vol. 13, no. 10, pp. 572-584, 2002.
- [57] A. K. Pandey, A. K. Mishra and A. Mishra, "Antifungal and antioxidative potential of oil and extracts derived from leaves of Indian spice plant *Cinnamomum tamala*," *Cellular and Molecular Biology*, vol. 58, no. 1, pp. 142-147, 2012.
- [58] X. H. Meng, C. Liu, R. Fan, L. F. Zhu, S. X. Yang, H. T. Zhu and Y. J. Zhang, "Antioxidative flavan-3-ol dimers from the leaves of *Camellia fangchengensis*," *Journal of agricultural and food chemistry*, vol. 66, no. 1, pp. 247-254, 2018.
- [59] S. Kumar and A. K. Pandey, "Chemistry and biological activities of flavonoids: an overview," *The scientific world journal*, pp. 1-16, 2013.
- [60] L. H. Cazarolli, L. Zanatta, E. H. Alberton, M. S. R. Bonorino Figueiredo, P. Folador, R. G. Damazio and F. R. M. Barreto Silva, "Flavonoids: prospective drug candidates. Mini reviews in medicinal chemistry," vol. 8, no. 13, pp. 1429-1440, 2008.
- [61] R. Singh, S. Singh, S. Kumar and S. Arora, "Evaluation of antioxidant potential of ethyl acetate extract/fractions of *Acacia auriculiformis*," *Food and chemical toxicology*, vol. 45, no. 7, pp. 1216-1223, 2007.
- [62] X. Han, T. Shen and H. Lou, "Dietary polyphenols and their biological significance," *International journal of molecular sciences*, vol. 8, no. 9, pp. 950-988, 2007.
- [63] S. S. Ali, N. Kasoju, A. Luthra, A. Singh, H. Sharanabasava, A. Sahu and U. Bora, "Indian medicinal herbs as sources of antioxidants," *Food research international*, vol. 41, no. 1, pp. 1-15, 2008.
- [64] A. Canini, D. Alesiani, G. D'Arcangelo and P. Tagliatesta, "Gas chromatography-mass spectrometry analysis of phenolic compounds from *Carica papaya* L. leaf," *Journal of food composition and analysis*, vol. 20, no. 7, pp. 584-590, 2007.
- [65] R. F. S. P. B. Epand and R. M. Epand, "Bacterial lipid composition and the antimicrobial efficacy of cationic steroid compounds (Ceragenins)," *Biochimica et Biophysica Acta (BBA)-Biomembranes*, vol. 1768, no. 10, pp. 2500-2509, 2007.
- [66] W. A. Kukula-Koch and J. Widelski, "Alkaloids," in *Pharmacognosy*, S. Badal and R. Delgoda, Eds., London, UK, Academic Press, 2017, pp. 163-198.
- [67] G. A. Ayoola, H. A. Coker, S. A. Adesegun, A. A. Adepoju-Bello, K. Obaweya, E. C. Ezennia and T.

- O. Atangbayila, "Phytochemical screening and antioxidant activities of some selected medicinal plants used for malaria therapy in Southwestern Nigeria," *Tropical Journal of Pharmaceutical Research*, vol. 7, no. 3, pp. 1019-1024, 2008.
- [68] I. Ahmad, Z. Mehmood and F. Mohammad, "Screening of some indian medicinal plants for their antimicrobial properties," *Journal of Ethnopharmacology*, vol. 62, pp. 183- 9, 1998.
- [69] L. Lu, S. W. Liu, S. B. Jiang and S. G. Wu, "Tannin inhibits HIV-1 entry by targeting gp41," *Acta Pharmacologica Sinica*, vol. 52, no. 2, pp. 213-218, 2004.
- [70] K. Funatogawa, S. Hayashi, H. Shimomura, T. Yoshida, T. Hatano and H. H. Y. Ito, "Antibacterial activity of hydrolyzable tannins derived from medicinal plants against *Helicobacter pylori*," *Microbiology and immunology*, vol. 48, no. 4, pp. 251-261, 2004.
- [71] C. G. Awuchi, "The Biochemistry, Toxicology, and Uses of the acologically Active Phytochemicals: Alkaloids, Terpenes, Polyphenols, and Glycosides," *Merit Research Journals*, vol. 5, no. 1, pp. 6-21, 2020.
- [72] V. O. Njoku and C. Obi, "Phytochemical constituents of some selected medicinal plants," *African journal of pure and applied chemistry*, vol. 3, no. 11, pp. 228-233, 2009.
- [73] K. Korir, C. Bii, C. Kiiyukia and C. Mutai, "Antimicrobial activities of *Clusia abyssinica* and *Erythrina abyssinica* plants extracts used among the Kipsigis community of Bomet district in Kenya," *Natural Products*, vol. 7, no. 5, pp. 247 - 252, 2011.
- [74] S. N. Njeru and M. A. Obonyo, "Potency of extracts of selected plant species from Mbeere, Embu County-Kenya against *Mycobacterium tuberculosis*," *Journal of Medicinal Plants Research*, vol. 10, no. 12, pp. 149-157, 2016.
- [75] M. Z. Salem, A. Z. Salem, L. M. Camacho and H. M. Ali, "Antimicrobial activities and phytochemical composition," *African Journal of Microbiology Research*, vol. 7, no. 33, pp. 4207-4219, 2013.
- [76] O. Odunbaku, O. Ilusanya and K. Akasoro, "Antibacterial activity of ethanolic leaf extract of *Ficus exasperata* on *Escherichia coli* and *Staphylococcus albus*," *Scientific Research and Essay*, vol. 3, no. 11, pp. 562-564, 2008.
- [77] G. Chilufya, G. Masaiti, C. Malambo and S. Mudenda, "Antibacterial Properties of *Ficus sycamoros* Bark Extract Against *Staphylococcus aureus* and *Escherichia coli*," *International Journal of Biomedical Investigation*, vol. 2, no. 121, pp. 1-8, 2019.
- [78] A. A. Mostafa, A. A. Al-Askara, K. S. Almaary, T. M. Dawoud, E. N. Sholkamy and M. M. Bakric, "Antimicrobial activity of some plant extracts against bacterial strains causing food poisoning diseases," *Saudi Journal of Biological Sciences*, vol. 25, no. 2, pp. 361-366, 2018.
- [79] S. Alibi, C. Dámaso and J. Navas, "Plant-Derivatives Small Molecules with Antibacterial Activity," *Antibiotics*, vol. 10, no. 3, pp. 23-30, 2021.
- [80] B. Khameneh, M. Iranshahy, V. Soheili and B. S. Bazzaz, "Review on plant antimicrobials: a mechanistic viewpoint," *Antimicrobial Resistance and Infection Control*, vol. 8, no. 1, pp. 1-28, 2019.
- [81] M. Friedman, P. Henika, C. Levin and R. Mandrell, "Antibacterial activities of plant essential oils and their components against *Escherichia coli* O157:H7 and *Salmonella enterica* in apple juice," *Journal of Agricultural and Food Chemistry*, vol. 52, pp. 6042-6048, 2004.
- [82] B. Tiwari, V. Valdramidi, C. O'Donnell, K. Muthukumarappan, P. Bourke and P. Cullen, "Application of natural antimicrobials for food preservation," *Journal of Agricultural and Food Chemistry*, vol. 57, pp. 5987-6000, 2009.
- [83] I. Musa, S. S. Manga, A. Muhammad, Z. Y. Tondi and M. Saadu, "The In vitro Antibacterial

- Effects of *Ficus sycomorus* Stem Bark Extracts on *Salmonella typhi* and *Escherichia coli*," *BiblioMed*, vol. 7, no. 1, pp. 108-113, 2020.
- [84] Y. Li, D. Kong, Y. Fu, M. R. Sussman and H. Wu, "The effect of developmental and environmental factors on secondary metabolites in medicinal plants," *Plant Physiology and Biochemistry*, vol. 148, pp. 80-89, 2020.
- [85] D. Manimozhi, S. Sankaranarayanan and G. Sampathkumar, "Evaluating The Antibacterial Activity Of Flavonoids Extracted From *Ficus Benghalensis*," *International Journal of Pharmaceutical and Biological Research*, vol. 3, no. 1, 2012.
- [86] C. Mudzengi, A. Murwira, M. Tivapasic, C. Murungweni, J. Burumue and T. Halimani, "Antibacterial Activity of Aqueous and Methanol Extracts of Selected Species Used in Livestock Health Management," *Pharmaceutical Biology*, vol. 55, pp. 1054-1060, 2017.

UNDER PEER REVIEW