

Role of Intestinal Parasitic Infection in Stunting

ABSTRACT

Aim: to review recent findings in the context of the relation between intestinal parasitic infection (IPI) induce the host's chronic immune activation and systemic inflammation that contributes to stunting

Discussion: to some extent, based on the parasite species (with the account of the condition of co-infections), combined with its timing (when was the initial infection took place), the duration of the disease, its intensity, and also must be considered the pathophysiology effects, can cause serious effect to their host. Prolonged IPI contributes to the formation of stunting. Active IPI can cause direct dysregulation of growth factors which are important for prenatal and postnatal growth. Furthermore, IPI opens the door for the entry of secondary infections, bacterial and viral, and allowed them to established definite co-infection which can cause prolonged inflammation, locally and systematically. Several potential interrelated mechanistic routes also could be connected through the host's imbalanced nutritional status, possible environmental enteric dysfunction (EED) due to persistent IPI and prolonged inflammation, hormonal derangement and perhaps also due to metabolic disturbances, immune dynamics, the consequences of definite anaemia, changed microbiota composition, and also changes in epigenetic signature.

Conclusion: Persistent and prolonged IPI induced both chronic immune activation and systemic inflammation that contributes to the formation of stunting, directly and indirectly.

Keywords: helminths, protozoan, co-infection, children, nutrition, environmental enteric dysfunction, microbiota composition

INTRODUCTION

Intestinal parasitic infection (IPI) can contribute to the formation of stunting, with *Ascaris lumbricoides* and hookworms contributed the most notable to stunting.[1] Individuals who experienced childhood stunting are more likely to encounter higher risk of morbidity [2], mortality [3], and its unnoticed consequences, such as less optimal cognitive function [4] and serious delay in motoric advancement. [5] According to The UNICEF-WHO-World Bank Joint Child Malnutrition Estimates (JME) Working Group, in the year of 2020, 149.2 million children under 5 years of age were too short for their age (stunting).

Unfortunately, the progression to eliminate stunting is not as expected, even reduction in prevalence are still uneven and still shows wide gap of differences in many countries. [7] Fortunately, because reduction of stunting is part of the MDG's targets, global research and networking has been established and has produced a lot of thought-provoking products for stunting alleviation. Understanding is developed regarding the causes of stunting, one of them is that enormous fraction of stunting is not caused by just diarrhea or inappropriate diet

[9], it remains that other factors must explain continued growth faltering.[9,10]. Persistent IPI contributes to the origination of stunting.

The pathognomonic esplanade by which IPI lead to stunting must always be considered multi-factorial [1] and milieu-specific, [11] and dependent on various interrelated factors such as parasite species, personal hygiene practice, geographic location, and a multitude of other circumstantial elements. The aim of this mini-review is to re-visit and refresh knowledge about how IPI contributes to the genesis of stunting.

THE VICIOUS CYCLE OF STUNTING

Until nowadays, the most globally accepted pathway to stunting is a theory named 'vicious cycle'. [12] Stunting is a result of continuous cyclical process based on women who were themselves stunted in childhood, in the future tend to have stunted offspring; this generating an intergenerational cycle of poverty and diminished human capital that is almost impossible to break by themselves, unless there is assistance from outside parties in power (the government) in breaking the chain of this cycle.




Fig. 1. The vicious cycle of stunting

Stunting vicious cycle revealed the connection between double burden of deteriorating nutritional status and infection [13] which is in turn progresses to into the following:

- (i) Encompass dysbiosis of the intestinal normal microbiota [14,15]. The vertebrate intestine be overflowing with a huge, miscellaneous, and vigorous inhabitants microbiota community [15] that has ubiquitous outcomes on overall intestine function, physiologically, biochemically, and last but not least immunity [14,15]. In normal situations, these inhabitants microbiota dividend their habitation with a correspondingly spirited neighborhood of eukaryotes organisms (e.g., helminths, protozoans, and also fungi), many of which are well-known parasites. This sudden combo group of the existing prokaryotic microbiota and the 'new comer' parasites can fiercely convert the actual physical and immune topography of the intestine, and fabricating tolerable possibilities for both parties (normal microbiota and parasites) to merge or at least to interact.[16] Such an unique relationship may substantially affect

the infection courses and outcomes and definitely influenced the condition of host's health and disease, comprehensively. For example, sudden parasite invasion can alter normal host association with its normal bacterial flora,[15] either compelling or guarding against the condition of dysbiosis and inflammatory disease.[17,18] Conversely, the microbiota can alter a parasite's colonization success, replication, and virulence, shifting it along the parasitism-mutualism spectrum [14,16,17].

- (ii) Initiating locally intestinal restricted inflammation [19,20] Helminth parasites are the classic perverters of Th2 responses. Immunologically, the Th2-polarized T cell response driven by helminth infection has been linked to the attenuation of some damaging Th1 driven inflammatory feedback, definitely blocking several Th1-mediated autoimmune conditions in the host, including experimentally induced colitis.[21] Scientifically, there is an antithetical interdependence between the amount of revelation to the eukaryotes intestinal helminths and the occurrence of unequivocal immune-mediated conditions, including inflammatory bowel diseases (IBD). intestinal helminth infection can prevent the outgrowth of a common intestinal bacterium that causes IBD in genetically susceptible mice.[20] However, recent studies have provided evidence indicating the exacerbating effects of helminths on bacterial as well as non-infectious colitis in animal models [18,19]. According to study conducted by Brosschot et al, [22] intestinal helminth infection can impair host resistance to co-infection with enteric bacterial pathogens Salmonella and the administration of anthelmintic regiment prior to Salmonella challenge is adequate to reestablish host initial resistance to Salmonella.
- (iii) Initiating systemic inflammation [18,23]. Intestinal helminths can manage deleterious inflammatory reactions and in the same time keeping homeostasis by way of stimulating systemic immune responses. The induction of distinct elements of immunity awakened by the existence of certain helminths, which includes Type 2 and immune regulatory responses, that can both contribute toward the reduction in harmful Type 1 immune responses that drive certain inflammatory diseases. Regardless of inducing type 2 responses, IPI may also down regulate harmful Type 2 immune responses including allergic responses. We consider the possibility that intestinal helminth infection may indirectly affect inflammation by influencing the composition of the intestinal microbiome.[25] Some previous study analyzed the effect of intestinal parasite to their host's systemic inflammation. According to Caudet et al, [24] enteric eukaryotic unicellular intestinal parasites may play an important role in modulating the antioxidant defenses of an obese host, thus could have beneficial effects with respect to the development of systemic metabolic disorders. While other study by Rajamanickam et al [23] revealed that helminth infection is associated with attenuation of systemic inflammation and microbial translocation in Type 2 Diabetes Mellitus and its reversal following anthelmintic therapy.
- (iv) The loss of energy due to hormonal and metabolic consequences of IPI.[26,27] A variety of helminths spend some part of their life cycle in the gastrointestinal tract and even entirely enteral nematode infections exert beneficial effects on glucose and lipid metabolism. Parasitic nematodes can impact metabolism including effects gastrointestinal hormones, altering epithelial function, and changing the number and/or phenotype of immune cells in metabolic tissues. Nematodes can also exert their beneficial effects through Th2 cytokines that activate the transcription factor STAT6, which upregulates genes that regulate glucose and lipid metabolism. Parasitic infections, even those restricted to the intestine, increase circulating levels of IL-4, IL-5, and IL-13 which may act to blunt or reverse the Th1-induced

inflammation in metabolic tissues. In the context of hormonal consequences, many parasites produce ecdysteroids, but restricted evidence is convenient on sex steroid and corticosteroid amalgamation. In most, but not all cases of parasite infection, the host's hormonal milieu regulates the susceptibility, the course, and severity of parasite infections. [28] In most cases, parasitic infection disturbs the normal existing host immune environment,[15] and actually triggers immune reactions that become stirring the host's endocrine armamentarium. In addition, sex steroids and corticosteroids may also straightly reshape the parasite reproductive ability and its molting.[29] Available information stipulates that living parasitizing organisms synthesize some steroid hormones, e.g., ecdysteroids and sex steroids, and the presence and activity of related enzymes have been studied elsewhere. [27]

THE MECHANISM BY WHICH PARASITIC INFECTION CONTRIBUTES TO STUNTING FORMATION

The human GI tract accommodate diverse but dynamic normal microbiota population since very early of life. These organisms contributes positively on its host's well-being, biochemically and physiologically, in terms of local milieu and also systemic function. Whenever there is invading parasitic eukaryotes, these normal microbiota must share their milieu with one main goal, namely to maintain homeostasis, whatever the cost. The bidirectional relationship between intestinal parasites and the existing microbiota [15] if happens very early (childhood early life), and their combined effects, [15,30] could play a key role in stunting. Intestinal parasite infection may contribute to stunting formation by way of:

- a. directly induce prolonged and persistent sub clinical systemic inflammation [1,2] and having an immunomodulatory properties that interfere with all features of the host immune response to ensure their persistence within the host [3] which has a detrimental effect on the individual who experiences it, or
- b. indirectly by triggering persistent local gut inflammation as a result of Environmental enteric dysfunction (EED), [31,32] an uninterrupted inflammatory state, persistently, derived via the adaptive immune reaction, which also has a serious impact on the digestive and absorptive process. The presence of persistent yet active sub-clinically intestinal parasitic infections was unquestionably linked with stunting but not with wasting. [11] Intestinal parasites can also collaborate with normal gut microbiota (and it can alter each other's pathogenicity), and this interaction also leading to malnutrition [5]

The condition of EED, even though initially it only took place locally and causing subclinical condition in term of disease progression, but we must realize that the condition of this intestinal inflammation, can occur since infants-age, most common found across underdeveloped countries, and is proposed as an immediate causal factor connecting poor sanitation and stunting.[31] A result of long term pathogen exposure (often found in types of infection by multicellular parasites that inhabited organs for a very long time), EED presents multiple causal pathways.[33] Although the definite pathogenesis of EED and the mechanism by which stunting occurs are yet to be defined, but evidence regarding contribution of intestinal parasitic infection cannot be ignored.

HOW CALORIES AND NUTRIENTS ARE LOST DUE TO THE COMBINATION OF IPI AND STUNTING?

One consideration governing immune-metabolic interactions includes sharing versus competition for energetic resources. [34] Immune activation cannot be denied responsible for

the lost of calorie intake and vital nutrients, that normally should be used for other routine non-immune activity (e.g., study, exercise etc).

Unfortunately, active immune armamentarium is actually high priced, in terms of both energetically and metabolically; and this may directly (or indirectly via other pathway) divert calories and nutrients away from other normal and routine physiological processes such as linear growth, in an energetic trade-off.[35,36] A study conducted by Urlacher et al in Amazonian youngster found out that substantial increased in total IgE level, antibody against parasite as an adaptive immune reactions, was correlated with definite growth inhibitions. [35]




Figure 2 Possible mechanistic pathway of stunting that begins with intestinal parasitic infection in the very early phase of life.

The sequence of events can overlap and occur simultaneously . These chains of events can exacerbate the course of the disease. note that short-term effects can occur immediately and quickly, but if not handled adequately they can continue to become serious long-term effects and may be irreversible clinically

Energetic tradeoffs implying human immune armamentarium activation are to be expected to arise since the very early of life, [37] based on exposure to outside world, that shape the way humans respond efficiently to energy needs while synchronizing with resilience to stimuli and even promote tolerance (e.g. to extreme heat).[38] The more active the immune system, the higher the energy needed.[39] This is a characterization in human evolution by considerable controlled energy allotment. [40] To some extent, physical development during growth spur phase is predicted to be exceptionally subtle to immune-related tradeoffs at this occasion. [36,37,40] It is not surprising that ordinary people also know that children who are often sick usually do not have good physical growth.

The direct consequences of IPI is that it may increase gut permeability, local (gut) inflammation, and systemic inflammation. [18-21] Parasites not inhabiting in the gastrointestinal tract may still cause systemic inflammation directly via activation of the immune response. [17] Changes in the gut microbiome and epigenetic signature of the host may also be implicated in the pathway to stunting, via the gut and/or systemic inflammatory

response. [41,42] Gut permeability and inflammation (associated with environmental enteric dysfunction) and systemic inflammation may be causative in the pathway to stunting. [31,32]

Furthermore, any multicellular parasites may manipulate amino acids, essential substance needed for the host's growth, by way of:

- 1) competing with their host and utilize them up for their own benefit, [43] or
- 2) via sudden and escalating requirement by the immune system during infection; usually bacterial based that it is usually a bacterial-based type of infection that occurs as a secondary infection or co-infection with the primary IPI. [44]

LOSS OF ESSENTIAL AMINO ACIDS AND ITS CONSEQUENCES

Essential amino acids are critical for nucleic acid and hormone biosynthesis and cellular replication, [45] and in turn for the long term, amino acids, especially that comes from animal [46], are very good for child development and growth, Anatomically and Neuro-cognitively.

According to Parikh et al, [45] the high incidence of growth faltering among children under 5 years of age, even if those kids are lucky enough not to be stunted, but still they underwent the condition of unable to outstretch their maximum growth potential. The time between conception and 2 years of age is critical for development. [47] It is often found, especially among mothers who are having children for the first time, at the age of 6-23 months is actually a critical period for inhibiting child growth due to the initial introduction and administration of complementary foods for their children, but unfortunately it concur mostly with the initial formation of growth faltering and accompanied with postponed neurocognitive developments; two most common condition found which could have preceded the stunting event. [48] Fortunately, this is also the time period of which diet exercises is familiarized, culturally. [49]

According to Semba et al, [50] child stunting is associated with definite poor circulating essential amino acids., systematically Furthermore they stated that unfortunate children with a high possibility of developing stunting may not be obtaining sufficient daily dietary intake that consist of essential amino acids and choline, an important nutrient for the synthesis of sphingolipids and glycerophospholipids.

Unfortunate children who grow up in a poor milieu, with a constant long-term deficient and inappropriate dietary intake, as typically seen in poor countries, inhibits growth and development of children and prevents them from achieving their maximum potential developmental and economic future potential. [51] The availability of adequate nutrient, in the context of quality and quantity, and its main implementation which is to make sure child's growth and development, especially in the post- breastfeeding period. [52] Children from poor country, although it is possible that children consuming sufficient total protein, but unfortunately it might fail to fulfill their essentials amino acids requirements.[53] The underlying condition for that perhaps as a consequences of poor diet heterogeneity and under-quality dietary protein; conditions that are very likely to be found in low level socio-economic society with low purchasing power and almost non-existent bargaining power in the community; it is a very good example of a marginalized poor society.

Amino acids deficiencies that took place in a very early life can cause limitations in linear growth-spur and neuro-cognitive advancement. [36] Assuring the availability and adequacy of amino acids in daily foods for children, especially through incorporation of nutrient-dense

animal source foods for children aged 6 to 23 months, is forcibly reassured in order to reimburse for previously less than optimal growth during complementary feeding.

Parasites may also utilize host amino acids; for example, intestinal protozoan *Giardia lamblia* directly consumes arginine as an energy source, and by doing so they showed their competitiveness and interactivity. [30] Furthermore, amino acids also contribute significantly for intestinal barrier function, with its shortage likely intensifying the EED cycle by way of diminishing the ability of intestinal cells to repair injury. [54] Amino acids ameliorate the (a) functions of intestinal barrier and expressions of anti-inflammatory cytokines and (b) tight junction proteins. Besides that, amino acids actively reduce (a) oxidative stress and (b) the apoptosis of enterocytes as well as (c) the expressions of proinflammatory cytokines in the intestinal inflammation.

Active IPI can cause direct dysregulation of growth factors which are important for prenatal and postnatal growth. [55] The consequences of immune activation upregulate acute-phase proteins, which prevent IGF-1 in the liver, directing to the condition of growth hormone (GH) resistance and inhibition of longitudinal bone growth via direct effects on the growth plate, potentially reducing optimal linear growth. [56] According to Maleta et al, that conducted a study among Malawian and children, infection caused by protozoan named *Cryptosporidium* spp. was correlated indirectly with lower length-for-age; a condition obtained by way of developed systemic inflammation and diminished plasma IGF-1 concentrations, [57] Another study conducted in Gambia also traversing these previously mentioned theories. [58] Mbuya et al [58] found that repeatedly appraised EED and its immune markers (representing normal gut microbiota translocation, intestinal permeability, and inflammatory reaction), together the three of them as a mechanism, forecasted of growth faltering condition up to 55%.

This condition can also occur in other types of parasitic infections, depending on the species, location, number and severity of the disease. Further exploration is needed to increase awareness, prevent fatalities and most importantly break the chain of infection so that stunting can be prevented.

CONCLUSION

Intestinal parasitic infection definitely contributes to the formation of stunting. The definite underlying mechanism is still being explored extensively. But in general speaking, the condition of EED and its successive inflammation triggered by IPI, these three together may be an underappreciated contributor to stunting, at least until recently. Fortunately, findings and achievements in the fields of nutrition, immunology, Internal Medicine, Public Health and clinical Parasitology/Microbiology contribute significantly to the understanding of the course of the disease with their consequences for stunting, and all of these has shed light on the global effort to prevent IPI and eliminate stunting.

CONSENT

Not needed

ETHICAL APPROVAL (WHERE EVER APPLICABLE)

Not needed

REFERENCES

1. Deka S, Kalita D, Hazarika NK. Prevalence and Risk Factors of Intestinal Parasitic Infection in Under-Five Children With Malnutrition: A Hospital Based Cross-Sectional Study. *J Family Med Prim Care*. 2022;11(6):2794-2801. https://doi.org/10.4103/jfmprc.jfmprc_1742_21.
2. Soekatri MYE, Sandjaja S, Syauqy A. Stunting Was Associated with Reported Morbidity, Parental Education and Socioeconomic Status in 0.5-12-Year-Old Indonesian Children. *Int J Environ Res Public Health*. 2020;17(17):6204. <https://doi.org/10.3390/ijerph17176204>.
3. Wright CM, Macpherson J, Bland R, Ashorn P, Zaman S, Ho FK. Wasting and Stunting in Infants and Young Children as Risk Factors for Subsequent Stunting or Mortality: Longitudinal Analysis of Data from Malawi, South Africa, and Pakistan. *J Nutr*. 2021;151(7):2022-2028. <https://doi.org/10.1093/jn/nxab054>.
4. Handryastuti S, Puspongoro HD, Nurdadi S, Chandra A, Pramita FA, Soebadi A, Widjaja IR, Rafli A. Comparison of Cognitive Function in Children with Stunting and Children with Undernutrition with Normal Stature. *J Nutr Metab*. 2022;2022:9775727. <https://doi.org/10.1155/2022/9775727>.
5. Mustakim MRD, Irwanto, Irawan R, Irmawati M, Setyo-boedi B. Impact of Stunting on Development of Children between 1-3 Years of Age. *Ethiop J Health Sci*. 2022;32(3):569-578. <https://doi.org/10.4314/ejhs.v32i3.13>.
6. The UNICEF-WHO-World Bank Joint Child Malnutrition Estimates (JME) Working Group. Technical Notes from the background document for country consultation on the 2021 edition of the UNICEF-WHO-World Bank Joint Malnutrition Estimates SDG Indicators 2.2.1 on stunting, 2.2.2a on wasting and 2.2.2b on overweight. 2021
7. Vaivada T, Akseer N, Akseer S, Somaskandan A, Stefopoulos M, Bhutta ZA. Stunting in childhood: an overview of global burden, trends, determinants, and drivers of decline. *Am J Clin Nutr*. 2020; 112 (Suppl 2):777S-91S. <https://doi.org/10.1093/ajcn/nqaa159>.
8. Bhutta ZA, Akseer N, Keats EC, Vaivada T, Baker S, Horton SE, et al. How countries can reduce child stunting at scale: lessons from exemplar countries. *Am J Clin Nutr*. 2020;112(Suppl 2):894S-904S. <https://doi.org/10.1093/ajcn/nqaa153>
9. Akseer N, Vaivada T, Rothschild O, Ho K, Bhutta ZA. Understanding multifactorial drivers of child stunting reduction in Exemplar countries: a mixed-methods approach.

The American Journal of Clinical Nutrition, 2020;112, 792S - 805S. <https://doi.org/10.1093/ajcn/nqaa152>

10. Isanaka S, Hitchenings MDT, Berthé F, Briend A, Grais RF. Linear growth faltering and the role of weight attainment: Prospective analysis of young children recovering from severe wasting in Niger. *Matern Child Nutr.* 2019;15(4):e12817. doi: <https://doi.org/10.1111/mcn.12817>.
11. Yoseph, A., Beyene, H. The high prevalence of intestinal parasitic infections is associated with stunting among children aged 6–59 months in Boricha Woreda, Southern Ethiopia: a cross-sectional study. *BMC Public Health*, 2020; 20: 1270. <https://doi.org/10.1186/s12889-020-09377-y>
12. de Onis M, Branca F. Childhood stunting: a global perspective. *Matern Child Nutr.* 2016;12 Suppl 1(Suppl 1):12-26. <https://doi.org/10.1111/mcn.12231>.
13. Prendergast AJ, Humphrey JH. The stunting syndrome in developing countries. *Paediatr Int Child Health.* 2014;34(4):250-65. <https://doi.org/10.1179/2046905514Y.0000000158>.
14. Leung JM, Graham AL, Knowles SCL. Parasite-Microbiota Interactions With the Vertebrate Gut: Synthesis Through an Ecological Lens. *Front Microbiol.* 2018;9:843. <https://doi.org/10.3389/fmicb.2018.00843>.
15. Sunarti LS. Microbial Normal Flora: Its Existence And Their Contribution To Homeostasis. *Journal of Advances in Microbiology*, 2022; 22(9): 1-15 <https://doi.org/10.9734/JAMB/2022/v22i930483>
16. Llinás-Caballero K, Caraballo L. Helminths and Bacterial Microbiota: The Interactions of Two of Humans' "Old Friends". *International Journal of Molecular Sciences.* 2022; 23(21):13358. <https://doi.org/10.3390/ijms232113358>
17. Maizels RM, McSorley HJ. Regulation of the host immune system by helminth parasites. *J Allergy Clin Immunol.* 2016;138(3):666-675. <https://doi.org/10.1016/j.jaci.2016.07.007>.
18. Zavala GA, García OP, Camacho M, Ronquillo D, Campos-Ponce M, Doak C, Polman K, Rosado JL. Intestinal parasites: Associations with intestinal and systemic inflammation. *Parasite Immunol.* 2018;40(4):e12518. <https://doi.org/10.1111/pim.12518>.
19. Giacomini P, Croese J, Krause L, Loukas A, Cantacessi C. Suppression of inflammation by helminths: a role for the gut microbiota? *Philos Trans R Soc Lond B Biol Sci.* 2015;370(1675):20140296. <https://doi.org/10.1098/rstb.2014.0296>.
20. Harris, N. Intimate gut interactions: helminths and the microbiota. *Cell Res* 2016;26: 861–2. <https://doi.org/10.1038/cr.2016.72>
21. Wang LJ, Cao Y, Shi HN. Helminth infections and intestinal inflammation. *World J Gastroenterol.* 2008;14(33):5125-32. <https://doi.org/10.3748/wjg.14.5125>.
22. Brosschot TP, Lawrence KM, Moeller BE, Kennedy MHE, FitzPatrick RD. Impaired host resistance to Salmonella during helminth co-infection is restored by anthelmintic

treatment prior to bacterial challenge. *PLOS Neglected Tropical Diseases* 2021; 15(1): e0009052. <https://doi.org/10.1371/journal.pntd.0009052>

23. Rajamanickam A, Munisankar S, Menon PA, Dolla C, Nutman TB, Babu S. Helminth Mediated Attenuation of Systemic Inflammation and Microbial Translocation in Helminth-Diabetes Comorbidity. *Front Cell Infect Microbiol.* 2020;10:431. <https://doi.org/10.3389/fcimb.2020.00431>.
24. Caudet J, Trelis M, Cifre S, Tapia G, Soriano JM, Rodrigo R, Merino-Torres JF. Do Intestinal Unicellular Parasites Have a Role in the Inflammatory and Redox Status among the Severely Obese? *Antioxidants.* 2022; 11(11):2090. <https://doi.org/10.3390/antiox11112090>
25. Mishra P, Palma M, Bleich D. Systemic impact of intestinal helminth infections. *Mucosal Immunol* 2014;7: 753–62. <https://doi.org/10.1038/mi.2014.23>
26. Shea-Donohue T, Qin B, Smith A. Parasites, nutrition, immune responses and biology of metabolic tissues. *Parasite Immunol.* 2017;39(5):10.1111/pim.12422. <https://doi.org/10.1111/pim.12422>.
27. Romano MC, Jiménez P, Miranda-Brito C, Valdez RA. Parasites and steroid hormones: corticosteroid and sex steroid synthesis, their role in the parasite physiology and development. *Front Neurosci.* 2015;9:224. <https://doi.org/10.3389/fnins.2015.00224>.
28. Wesołowska, A. Sex—the most underappreciated variable in research: insights from helminth-infected hosts. *Vet Res,* 2022; 53: 94. <https://doi.org/10.1186/s13567-022-01103-3>
29. Hernández-Bello R, Ramirez-Nieto R, Muñiz-Hernández S, Nava-Castro K, Pavón L, Sánchez-Acosta AG, Morales-Montor J. Sex steroids effects on the molting process of the helminth human parasite *Trichinella spiralis*. *J Biomed Biotechnol.* 2011:625380. <https://doi.org/10.1155/2011/625380>.
30. Siagian FE. Intestinal Microflora Vs Protozoan Parasites: From Interaction to Competition. *South Asian Journal of Research in Microbiology,* 2022; 13(1): 36-46. <https://doi.org/10.9734/SAJRM/2022/v13i130290>
31. Budge S, Parker AH, Hutchings PT, Garbutt C. Environmental enteric dysfunction and child stunting. *Nutr Rev.* 2019;77(4):240-53. <https://doi.org/10.1093/nutrit/nuy068>.
32. Harper KM, Mutasa M, Prendergast AJ, Humphrey J, Manges AR. Environmental enteric dysfunction pathways and child stunting: A systematic review. *PLoS Negl Trop Dis,* 2018; 12(1): e0006205. <https://doi.org/10.1371/journal.pntd.0006205>
33. Chen C, Lee WC. Attributing diseases to multiple pathways: a causal-pie modeling approach. *Clin Epidemiol.* 2018;10:489-97. <https://doi.org/10.2147/CLEP.S160205>.
34. Zmora N, Bashirdes S, Levy M, Elinav E. The Role of the Immune System in Metabolic Health and Disease. *Cell Metab.* 2017;25(3):506-21. <https://doi.org/10.1016/j.cmet.2017.02.006>.

35. Urlacher SS, Ellison PT, Sugiyama LS, Pontzer H, Eick G, Liebert MA, et al. Tradeoffs between immune function and childhood growth among Amazonian forager-horticulturalists. *Proc Natl Acad Sci USA*. 2018;115(17):E3914-21. <https://doi.org/10.1073/pnas.1717522115>.
36. Inzaghi E, Pampanini V, Deodati A, Cianfarani S. The Effects of Nutrition on Linear Growth. *Nutrients*. 2022;14(9):1752. <https://doi.org/10.3390/nu14091752>.
37. Garcia AR, Blackwell AD, Trumble BC, Stieglitz J, Kaplan H, Gurven MD. Evidence for height and immune function trade-offs among preadolescents in a high pathogen population. *Evol Med Public Health*. 2020;2020(1):86-99. <https://doi.org/10.1093/emph/eoaa017>.
38. Ganeshan K, Nikkanen J, Man K, Leong YA, Sogawa Y, Maschek JA, et al. Energetic Trade-Offs and Hypometabolic States Promote Disease Tolerance. *Cell*. 2019;177(2):399-413.e12. <https://doi.org/10.1016/j.cell.2019.01.050>.
39. Bird, L. Getting enough energy for immunity. *Nat Rev Immunol*, 2019; 19: 269. <https://doi.org/10.1038/s41577-019-0159-y>
40. Pontzer H. The crown joules: energetics, ecology, and evolution in humans and other primates. *Evol Anthropol*. 2017 Jan;26(1):12-24. <https://doi.org/10.1002/evan.21513>.
41. Durack J, Lynch SV. The gut microbiome: Relationships with disease and opportunities for therapy. *J Exp Med*. 2019 Jan 7;216(1):20-40. <https://doi.org/10.1084/jem.20180448>.
42. Woo V, Alenghat T. Epigenetic regulation by gut microbiota. *Gut Microbes*. 2022 Jan-Dec;14(1):2022407. <https://doi.org/10.1080/19490976.2021>.
43. Krishnan A, Soldati-Favre D. Amino Acid Metabolism in Apicomplexan Parasites. *Metabolites*. 2021; 11(2):61. <https://doi.org/10.3390/metabo11020061>
44. Zhang YJ, Rubin EJ. Feast or famine: the host-pathogen battle over amino acids. *Cell Microbiol*. 2013;15(7):1079-87. <https://doi.org/10.1111/cmi.12140>.
45. Lopez MJ, Mohiuddin SS. Biochemistry, Essential Amino Acids. [Updated 2023 Mar 13]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK557845/>
46. Parikh P, Semba R, Manary M, Swaminathan S, Udomkesmalee E, Bos R, et al. Animal source foods, rich in essential amino acids, are important for linear growth and development of young children in low- and middle-income countries. *Maternal & Child Nutrition*, 2021;18:e13264. <https://doi.org/10.1111/mcn.13264>
47. Likhar A, Patil MS. Importance of Maternal Nutrition in the First 1,000 Days of Life and Its Effects on Child Development: A Narrative Review. *Cureus*. 2022;14(10):e30083. <https://doi.org/10.7759/cureus.30083>.
48. D'Auria E, Borsani B, Pendezza E, Bosetti A, Paradiso L, Zuccotti GV, Verduci E. Complementary Feeding: Pitfalls for Health Outcomes. *International Journal of Environmental Research and Public Health*. 2020; 17(21):7931. <https://doi.org/10.3390/ijerph17217931>

49. Puspitorini P, Lestari P, Paramashanti B. The risk of inappropriate timing of complementary foods introduction is increased among first-time mothers and poor households. *Jurnal Gizi Klinik Indonesia*, 2021; 17(3), 96-102. <https://doi.org/10.22146/ijcn.53443>
50. Semba RD, Shardell M, Sakr Ashour FA, Moaddel R, Trehan I, Maleta KM, Ordiz MI, Kraemer K, Khadeer MA, Ferrucci L, Manary MJ. Child Stunting is Associated with Low Circulating Essential Amino Acids. *EBioMedicine*. 2016;6:246-252. <https://doi.org/10.1016/j.ebiom.2016.02.030>.
51. Gubbels JS. Environmental Influences on Dietary Intake of Children and Adolescents. *Nutrients*. 2020;12(4):922. <https://doi.org/10.3390/nu12040922>.
52. Infant and Young Child Feeding: Model Chapter for Textbooks for Medical Students and Allied Health Professionals. Geneva: World Health Organization; 2009. SESSION 3, Complementary feeding. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK148957/>
53. Maulidiana AR, Sutjiati E. Low intake of essential amino acids and other risk factors of stunting among under-five children in Malang City, East Java, Indonesia. *J Public Health Res*. 2021;10(2):2161. <https://doi.org/10.4081/jphr.2021.2161>.
54. He F, Wu C, Li P, Li N, Zhang D, Zhu Q, et al. Functions and Signaling Pathways of Amino Acids in Intestinal Inflammation. *Biomed Res Int*. 2018;2018:9171905. <https://doi.org/10.1155/2018/9171905>.
55. Błaszowska J, Góralska K. Parasites and fungi as a threat for prenatal and postnatal human development. *Ann Parasitol*. 2014;60(4):225-34.
56. Dixit M, Poudel SB, Yakar S. Effects of GH/IGF axis on bone and cartilage. *Mol Cell Endocrinol*. 2021;519:111052. <https://doi.org/10.1016/j.mce.2020.111052>.
57. Maleta K, Fan YM, Luoma J, Ashorn U, Bendabenda J, Dewey KG, et al. Infections and systemic inflammation are associated with lower plasma concentration of insulin-like growth factor I among Malawian children. *Am J Clin Nutr*. 2021;113(2):380-390. <https://doi.org/10.1093/ajcn/nqaa327>.
58. Mbuya MN, Humphrey JH. Preventing environmental enteric dysfunction through improved water, sanitation and hygiene: an opportunity for stunting reduction in developing countries. *Matern Child Nutr*. 2016;12 Suppl 1(Suppl 1):106-20. <https://doi.org/10.1111/mcn.12220>