

## BURDEN OF ROTAVIRUS DIARRHOEA IN CHILDREN IN ABUJA

### ABSTRACT

**Background:** Diarrhoea is the second leading cause of death in children. Nigeria has the continent's highest mortality due to diarrhoeal diseases with little information on specific causes and the proportion affected by rotavirus infection. The main objectives of this study were to describe the features of young children with diarrhoea, with and without rotavirus, in Abuja the Federal Capital Territory.

**Materials and Methods:** Retrospective review of local hospital activity data related to diarrhoea was retrieved from General Hospitals in Abuja. Then a one-year prospective descriptive study of children under 5 years of age with acute diarrhoea was conducted from September 2012 to August 2013. Children with acute diarrhoea attending three government hospitals and one private hospital were recruited. Children without diarrhoea were recruited as the control group. Faecal specimens were transported in cold chain boxes from Nigeria and stored at -80 °C at the Institute of Global Health, virology section of the University of Liverpool, where all laboratory work was performed.

**Results:** Hospital records were poorly preserved and did not provide meaningful data for trend analyses or disease surveillance. 1331 participants were enrolled in this prospective study. Stool samples were collected from 1242 (93.3%) participants, of whom 957 (77.0%) were ambulatory, 123 (9.9%) hospitalised and 160 (12.8%) controls without diarrhoea. 881 and 450 children with diarrhoea were recruited from government and the private hospitals, respectively. The median age of the children was 8 months in the ambulatory and 9.5 months in the hospitalized group ( $p < 0.05$ ). A total of 209 (16.8%) children were vaccinated, 858 (69.1%) were unvaccinated, and 174 (14.0%) had an unclear vaccination status. Rotavirus ELISA was positive in 123 (11.4%) children with diarrhoea and 2 (1.2%) controls. Among children with diarrhoea, 92 (10.4%) of 881 children attending government hospitals had rotavirus, compared to 33 (7.3%) of 450 children attending a private hospital ( $p < 0.001$ ) where a vaccination program had been in place. The peak months for rotavirus infection were November and February. The efficacy of Rotarix® RV1 vaccine in preventing rotavirus diarrhoea was indirectly assessed to be 64.5%.

**Conclusion:** Rotavirus is an important pathogen in children, especially in unvaccinated children in Abuja. The introduction of a rotavirus vaccine is highly desirable in Nigeria, but this is still awaited. Local and national infrastructure are inadequate for basic surveillance of diarrhoeal disease, and this will have to be improved, together with access to virological stool testing, to monitor the planned vaccine program. The retrospective studies reported here provide indirect evidence of vaccine efficacy in Abuja but need prospective confirmation.

**Keywords:** Rotavirus, Acute Diarrhoea, Burden, Abuja

### INTRODUCTION

Diarrhoea is a significant public health problem in Nigeria, particularly among children under five years of age. According to the World Health Organization (WHO), Nigeria had an estimated 75,000 deaths due to diarrhoea in children under the age of five in 2019, making it one of the countries with the highest burden of diarrhoea globally (WHO 2023).

Rotavirus is a leading cause of diarrheal disease in Nigeria, particularly among children under the age of 5 years. In 2019, a study conducted in Nigeria reported that the prevalence of rotavirus in children with acute gastroenteritis was 32.5% (Sowunmi et al 2019). Another study conducted in the same year reported that rotavirus was responsible for 41.7% of cases of acute gastroenteritis in children under the age of 5 years (Ayukekbong et al 2019). The burden of rotavirus diarrhoea in Nigeria is significant, with an estimated 10,694 deaths among children under 5 years of age in 2019 alone (Gavi 2021). Factors contributing to the high burden of rotavirus disease in Nigeria include poor sanitation, lack of access to clean water, and inadequate healthcare infrastructure.

Despite the high hospitalisation and death figures due to rotavirus diarrhoea in Nigeria, rotavirus diarrhoea is mostly not well understood by the local population of Abuja residents. Rotavirus diarrhoea is regarded as any bowel infection which is followed by passing of loose stool. Many of the residents are not aware of the vaccine availability for this type of diarrhoea and much of diarrhoeal diseases are not particularly classified in the hospitals surveyed except extreme cases of severe dehydration of hospitalised patients are tested and classified as either diarrhoeal disease or gastroenteritis. Nigeria has no central health surveillance system to assess and follow disease trends, and rotavirus diarrhea is poorly diagnosed in children with acute diarrhoea. To address the diarrhoea burden in Nigeria, various interventions have been implemented, including the promotion of handwashing with soap, provision of safe drinking water, and vaccination against rotavirus, a common cause of severe diarrhoea in children (Oladebo et al 2019). However, challenges remain, including inadequate funding for diarrhoea prevention and control, inadequate healthcare infrastructure, and a lack of awareness among caregivers regarding proper hygiene and sanitation practices (Oladebo et al 2019).

The importance of this research cannot be overemphasized and cannot have come at a better time. The research has formed part of the national and international research priorities, as the Millennium Development Goal 4 aims to reduce child mortality by 2015. The aim of this study was to describe and document the age distribution, seasonality of rotavirus diarrhoea, and cases of rotavirus diarrhoea among vaccinated and unvaccinated children. The findings will be useful for planning the possible inclusion of rotavirus vaccine in the National Immunization Program, in addition to providing a baseline for monitoring the influence of the vaccine on prevalence when introduced in the immunization programme of Nigeria.

## **METHODOLOGY**

The study was a one-year prospective survey of children with acute diarrhoea attending Zankli Medical Center (ZMC), University of Abuja Teaching Hospital (UATH) Gwagwalada, Gwagwalada Town Clinic, and Nyanya General Hospital from 1st of September 1, 2012, to August 30, 2013. The pediatric unit of ZMC (private clinic) provides the Rotavirus vaccine (Rotarix), which is administered to children less than 6 months of age as part of their routine immunization schedule.

Children aged 1 month to 5 years with complaints of acute diarrhoea were included in the study at all centers. The pediatrician identified cases of diarrhoea and referred the patients to the researcher for recruitment. After obtaining informed consent from the parents or guardians, details of the clinical history and clinical examination of stool samples were collected. A questionnaire was administered to obtain demographic details of the patient, clinical history of the current diarrhoea episode, presence of dehydration, weight, height, other symptoms such as respiratory problems and rashes and vaccination status. Hospitalized patients were followed up to record their outcomes (death or discharge, use of intravenous solutions, and number of hospitalizations). Rotavirus vaccination status was confirmed by reviewing clinical records in ZMC. The severity of the clinical episode of diarrhoea was classified according to the Vesikari score (Ruuska and Vesikari, 1990), as modified by Nakagomi (Nakagomi et al., 2005). This is a standardised score currently accepted internationally to classify the severity of the diarrhoea episodes and is based on a series of clinical criteria. In addition to cases of diarrhoea, a control group of children without diarrhoea was included to compare the prevalence of pathogens in children with and without diarrhoea. Control children were selected from children under five years of age attending the Well Baby Clinic of UATH. Stool samples were aliquoted into a safe-lock tube and labelled with the name, date, and study number. The samples were tested for rotavirus using an ELISA kit (Rotaclone) at the University of Liverpool, United Kingdom. The vaccine effectiveness was estimated using the Hatton's method 1990.

## **RESULTS AND DISCUSSION**

### **General Characteristics of the population with and without diarrhoea**

A total of 1331 children were enrolled in the hospital survey from September 2012 to August 2013. The children were aged between 1 and 5 years. More males (n = 775) than females (n = 556) were enrolled in the study. Ninety-one (6.8%) patients left the hospital without providing samples, and 1240 stool samples were collected. One hundred and twenty nine (9.6%) of the 1240 were hospitalized while 1027 (77.1%) were ambulatory. One hundred and seventy five (13.1%) control children without diarrhoea were enrolled. The mean diarrhoea duration before consultation was 3 and 8 days for ambulatory and hospitalized children, respectively ( $p < 0.0001$ ). The mean number of stools per day was 2 and 5 for ambulatory and hospitalized children, respectively. Most children passed out watery stool for both hospitalized and ambulatory with an overall total of 635 (59.9%) and 380 (32.9%) had mucus. All 175 controls had solid stools.

The proportion of children experiencing vomiting was lower in ambulatory than hospitalized children (51.4% and 75.9%, respectively,  $p = <0.0001$ ). The mean frequency of vomits experienced before consultation was between 4 and 6 times for ambulatory and hospitalised children respectively ( $p<0.0001$ ). Less ambulatory children experienced abdominal pain (27.3%) than hospitalized children (41%) and the same trend was observed for fever ( $p<0.002$  and  $p<0.003$ , respectively).

Oral and intravenous rehydration were recorded in 1091 (94.38%) of the 1156 children. This figure (1091) comprises 132 and 925 children in the hospitalized and ambulatory groups, respectively. Of the hundred and thirty one (80.9%) and 114 (88.3%) of the ambulatory and hospitalised children were treated with antimicrobials, respectively ( $p<0.03$ ). The percentage of death due to diarrhoea was 2 (1.5%) for hospitalised and 3 (0.2%) for ambulatory children. The majority of hospitalized patients improved after ORS administration.

**Table 1 General Characteristics of the population with and without diarrhoea**

Parameters	Ambulatory N= 1027	Hospitalised N= 129	P value
Age in months, median (IQR)	8.0 (12.0, 24.5)	9.5 (12.5, 14.0)	<0.05
Male: Female (% male)	603:424 (58.7%)	73:56 (56.5%)	0.6
<i>Diarrhoea</i>			
Mean (SD) duration, days	3.4 (2.6)	8.0 (16.5)	<0.0001
Mean (SD) Max frequency/24h before attending	1.8 (2.6)	4.6 (4.0)	<0.0001
Presence of vomiting (%)	528 (51.4%)	98 (75.9%)	<0.0001
Mean (SD) duration, days	0.5 (0.5)	0.7 (0.4)	<0.0001
Mean (SD) Max Frequency/24h before attending	4.1 (1.6)	5.5 (2.4)	<0.0001
Fever (%)	819 (79.7%)	117 (90.7%)	<0.003
Abdominal Pain (%)	281 (27.3%)	53 (41.0%)	<0.002
Watery (%)	549 (53.4%)	86 (66.6%)	> 0.1
Solid (%)	132 (12.8%)	5 (3.8%)	
Mucus (%)	342 (33.3 %)	38 (29.4%)	
Mucus and blood (%)	4 (0.3%)	0 (0.0%)	
Bloody (%)	0 (0.0%)	0 (0.0%)	
Normal (%)	136 (13.2%)	8 (6.2%)	> 0.01
Reduced (%)	891 (86.7%)	121 (93.8%)	

<b>Mild (%)</b>	721 (70.2%)	26 (20.1 %)	<0.0001
<b>Moderate (%)</b>	285 (27.7%)	90 (69.7%)	<0.0001
<b>Severe (%)</b>	21 (2.0%)	13 (10.0%)	<0.0001
<b>Not Dehydrated</b>	0 (100%)	0 (100%)	
<b>None (%)</b>	89 (8.6%)	4 (3.1%)	<0.03
<b>ORT (%)</b>	925 (90.0%)	124 (96.1%)	<0.02
<b>I.V (%)</b>	34 (3.3 %)	8 (6.2%)	>0.1
<b>Antimicrobials used:</b>	831 (80.9%)	114 (88.3%)	<0.03

### Characteristics of the population with rotavirus diarrhoea

This table describes the characteristics of the population in the study of rotavirus diarrhoea. A total of 1242 patients provided samples in this study, of which 123 (10.06%) were rotavirus-positive and 1119 (90.09%) were rotavirus-negative. All children were aged between 1 and 5 years. The mean diarrhoea in days before the consultation was 4 days and 3 days for rotavirus positive and negative respectively ( $p < 0.0001$ ). The mean maximum diarrhoea in 24 hours is 3 times for both rotavirus-positive and negative ( $p < 0.0001$ ). The proportion of children that experienced vomiting was higher in rotavirus positive group than in the rotavirus-negative group (60.1% and 46.0%,  $p = < 0.0001$ ). The mean frequency of vomiting experienced 24 h before the consultation was four and three times for positive and negative cases, respectively. Although there was no significant difference between children that experienced fever between that rotavirus positive and negative ( $p = 0.0754$ ), more children in the positive group had fever compared to those rotavirus negative (78.0% and 70.3%), the same trend was experienced with the mean temperature as the rotavirus positive group experiencing higher temperatures compared with those negative (37.5 and 37.4,  $p = < 0.1622$ ). Assessing the level of activity between the two groups, fewer children had normal activity after having rotavirus diarrhoea compared with those that are rotavirus negative (15.45% and 26.12%). More children in the rotavirus-positive cases had reduced levels of activity compared with those rotavirus negative (84.55% and 76.92%,  $p = 0.0539$ ). The diarrhoea severity was estimated for 1242 children. In the rotavirus-positive children, 75.61% were mild, 21.14% were moderate and 3.25% were severe. Fewer children experienced severe symptoms in the rotavirus-positive group. Oral rehydration and intravenous were recorded for 1242 of the children in this study. 95.12% of the children with rotavirus were treated with oral rehydration therapy, and 1.63% were given intravenous injection while 82.93% were treated with antibiotics. In the rotavirus-negative children, 78.35% were treated with oral rehydration solution, 3.40 were given an intravenous injection and 70.48% were treated along with antibiotics. There were more children treated with

ORS in the two groups ( $p = <0.0001$ ) and more children were given no treatment in the rotavirus-negative children (20.57%,  $p = <0.0001$ ).

**Table 2. Characteristic feature of rotavirus diarrhoea**

Characteristic feature	Rotavirus status		pvalue
	Positive N=125	Negative N=1117	
<i>Diarrhoea</i>			
Mean (SD) Duration in days	4.08 (3.21)	3.39 (6.27)	<b>&lt;0.0001</b>
Mean (SD) max no of stool/24h	3.9 (2.31)	3.4 (2.99)	<b>0.0162</b>
<i>Vomiting</i>			
Vomiting (%)	60.1%	46.0%	
Mean (SD) Duration in days	0.60 (0.49)	0.46 (0.49)	<b>0.0030</b>
Mean (SD) max no of vomits/24h	4.03 (1.60)	3.72 (2.28)	<b>0.0766</b>
<i>Fever</i>			
Fever (%)	78.0%	70.3%	<b>0.0754</b>
Mean (SD) Temperature	37.5 (0.68)	37.4 (0.79)	<b>0.1622</b>
Abdominal pain (%)	17.8%	26.12%	<b>0.0463</b>
<i>Level of activity</i>			
Normal	15.45%	23.08%	<b>0.0539</b>
Reduced	84.55%	76.92%	<b>0.0539</b>
<i>Degree of dehydration</i>			
Mild	75.61%	68.60%	<b>0.1100</b>
Moderate	21.14%	28.8%	<b>0.0696</b>
Severe	3.25%	2.50%	<b>0.6197</b>
<i>Rehydration therapy</i>			
ORS	95.12%	78.35%	<b>&lt;0.0001</b>
IVT	1.63%	3.40%	<b>0.2909</b>
None	4.07%	20.57%	<b>&lt;0.0001</b>
Received antibiotics	82.93%	70.48%	<b>0.0036</b>

### Rotavirus detection by month

The table categorizes children with rotavirus diarrhea (both positive and negative cases) by month. The table generally showed February as the peak month of rotavirus diarrhoea 31 (23.1%) followed by November 28 (23.7%), January 23 (31.0%) and December 14 (1.75%). The months with the least cases of rotavirus diarrhoea were July 0 (0.0%) and September 2 (2.9%).

**Table 3: Rotavirus detection by month**

MONTHS			
	RV positive %	RV negative %	Total
September 2012	2 (1.6)	64	<b>66</b>
October 2012	6 (4.9)	56	<b>62</b>

<b>November 2012</b>	28 (22.8)	62	<b>90</b>
<b>December 2012</b>	14 (11.4)	29	<b>43</b>
<b>January 2013</b>	23 (18.7)	28	<b>51</b>
<b>February 2013</b>	31 (25.2)	76	<b>107</b>
<b>March 2013</b>	5 (4.1)	97	<b>102</b>
<b>April 2013</b>	4 (3.3)	108	<b>112</b>
<b>May 2013</b>	5 (4.1)	91	<b>96</b>
<b>June 2013</b>	1 (0.81)	99	<b>100</b>
<b>July 2013</b>	0 (0)	118	<b>118</b>
<b>August 2013</b>	4 (3.3)	125	<b>129</b>
<b>Total</b>	<b>123</b>	<b>953</b>	<b>1076</b>

### Rotavirus-positive cases by age groups most affected

A total of 123 rotavirus-positive cases were confirmed in the study. More cases of rotavirus diarrhoea were experienced (13.0%) by children between the ages of < 6 months in the University of Abuja Teaching Hospital which is slightly followed by 6-11 months. In Zankli Medical Centre, a similar trend was observed; however, children of 6-11 months had more positive cases than children aged < 6 months (21.5% and 20.0%,  $p \leq 0.0001$ ). In the Gwagwalada Town clinic, children aged 6-11 are the most affected (20.6%), while at Nyanya General Hospital, children with rotavirus were spread across three age groups with children of ages 18-23 (22.2%) having slightly higher positive case than children of ages 6-11 (21.2%).

**Table 4: Rotavirus positive cases by age groups most affected**

Age Group	<6*	6-11	12-17	18-23	24-29	30-59	Pvalue
Study Group	Pos/tested (% pos)	Pos/tested (% pos)	Pos/tested (% pos)	Pos/tested (% pos)	Pos/tested (% pos)	Pos/tested (% pos)	
<b>UATH</b>	13.04	10.81	7.44	6.25	2.70	-	0.329
<b>ZMC</b>	20.00	21.56	4.34	2.56	0.95	6.34	<0.0001
<b>GTC</b>	-	20.68	-	14.28	-	-	0.009
<b>NGH</b>	12.85	21.23	21.00	22.22	8.10	3.57	0.092

## CONCLUSION AND RECOMMENDATION

A full literature review suggests that diarrhoea, specifically rotavirus diarrhoea, has been a major cause of disease in children in different States in Nigeria (as in the rest of Africa) for more than 30 years. In most studies, the peak age affected is between 6 and 18 months, after weaning. The seasonal pattern of diarrhoea presentation and the ages affected are reported to be different in some northern states and require further investigation. The surveillance of diarrhoeal disease in Nigeria, the largest country in Africa, is inadequate for meaningful analysis of disease trends. Oral rehydration is managed according to WHO recommendations, but there is a marked overprescription of antibiotics, which needs to be addressed. The lack of availability of virological investigations in most clinics in Nigeria indicates that specific access is needed to support and monitor the effectiveness of vaccination campaigns. The results of this study suggest that Abuja resembles most other states in Nigeria and nearby countries in the incidence and prevalence of rotavirus diarrhoea in children under the age of 1 year and slightly older, and indirectly supports the effectiveness of the monovalent vaccine. The findings support the planned (but delayed) introduction of the rotavirus vaccine in Nigeria but emphasize the need for adequate investment in quality-assured surveillance and virological surveillance to monitor this. The emergence of new rotavirus genotypic combinations may pose a threat to vaccine efficacy in the future.

### Consent

As per international standard, parental written consent has been collected and preserved by the author(s).

## REFERENCES

1. Ayukekbong, J. A., Fobisong, C., Tah, F., Lindh, M., & Nkuo-Akenji, T. (2019). Prevalence and genotypes of rotavirus among children under 5 years with acute gastroenteritis in Cameroon: A systematic review and meta-analysis. *Archives of virology*, 164(8), 1979-1991. <https://doi.org/10.1007/s00705-019-04267-6>
2. Gavi, The Vaccine Alliance. (2021). Nigeria. <https://www.gavi.org/programmes-impact/country-hub/africa/nigeria>
3. Gollogly, L. (2009). World health statistics 2009. World Health Statistics 2009.
4. Junaid, S. A., Umeh, C., Olabode, A. O. & Banda, J. M. (2011). Incidence of rotavirus infection in children with gastroenteritis attending Jos university teaching hospital, Nigeria. *Virol J*, 8, 21575246.
5. Oladepo O, Adeoye BW, Oshiname FO. Diarrhoea in Nigeria: factors responsible for persistence and challenges to its control. *J Health Popul Nutr*. 2019;38:25.
6. Sowunmi, O. O., Olaleye, D. O., Adeyemo, A. A., & Faleye, T. O. C. (2019). Epidemiology and molecular characterization of rotavirus infection among children with acute gastroenteritis in Lagos, Nigeria. *BMC infectious diseases*, 19(1), 1023. <https://doi.org/10.1186/s12879-019-4638-7>

7. Tinuade, O., John, O., Saheed, O., Oyeku, O., Fidelis, N. & Olabisi, D. (2006). Parasitic etiology of childhood diarrhea. *Indian Journal of Pediatrics*, 73, 1081-1084.
8. World Health Organization. Diarrhoeal disease. Available at: <https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease>. Accessed on 16 March 2023.

UNDER PEER REVIEW