

Genetic correlation and path coefficient analysis of yield attributing parameters in Indian mustard

ABSTRACT

Indian mustard (*Brassica juncea* L. Czern. & Coss) is a natural amphidiploid which is the greatest pre-dominating crop of oilseed *Brassica* group. A study was undertaken to estimate the genetic variability, correlation and path coefficient analysis of yield and its contributing traits in 75 mustard cultivars grown in Randomized Block Design with two replications. The analysis of variance were highly significant for all the characters investigated. All thirteen characters were showed higher values of phenotypic coefficients of variation than genotypic coefficients of variation. The higher heritability in broad sense was estimated for all the characters. High value of heritability indicates that it may be due to higher contribution of genotypic components. High heritability coupled with high genetic advance as percent of means were recorded for days to 50% flowering, plant height (cm), number(s) of secondary branches per plant, length of main raceme (cm), siliquae length (cm), seed yield per plant (g), yield per plot (g), harvest index and biological yield that indicated predominance of additive gene action in the inheritance of these traits. The higher direct positive genotypic and phenotypic correlations for the biological yield, numbers of primary branches, numbers of siliquae on main raceme and number of secondary branches was documented. Whereas, days to maturity and siliquae length showed direct negative correlations with grain yield. Seventy-five varieties, included in study were grouped into 6 clusters. The maximum inter cluster D^2 value indicated that genotypes of cluster III and IV are not so closely related while the genotypes of cluster I and III are closely related. It is apparent therefore; the genotypes of various clusters differ so significantly with regards to their relative genetic distance as indicated from the high variation of D^2 values. This makes it clear that the genotypes included in these clusters have a wide range of genetic diversity and may be used in a mustard hybridization programme to develop higher yielding cultivars.

Key words: Genetic variability, correlation, path analysis and Indian mustard

INTRODUCTION

Indian mustard (*Brassica juncea* L. Czern. & Coss) is a natural amphidiploid ($2n = 36$, AABB genome) which is the greatest pre-dominating crop of oilseed *Brassica* group. It is self-pollinated with some degree of cross-pollination and with a physical genome size of 920 Mb [1]. It is the most significant oilseed crop of India having substantial economic, nutritional, and industrial applications [2-5]. Mustard is primarily cultivated throughout Asia, Europe, Canada, and the former Soviet Union, but the cultivars

planted there differ from those grown in India. The oil content varies slightly depending on the type of oil used [6-9]. India is the fourth-largest contributor of oilseeds in the world and Rapeseed and mustard contributes about 28.6% of total oil seeds production. During 2021-22 the area under coverage has been pegged at 87.44 lakh hectares while the average yield is seen at 1,270 kg ha⁻¹. Mustard is an important cash crop for farmers in Rajasthan, Haryana, Madhya Pradesh, and Uttar Pradesh, among others. Rajasthan is the largest producing state in the country [10].

Indian mustard yield and its constituents are quantitative in nature, thus learning more about the type, scope, and effects of genetic variability on the environment may be proved obliging. Therefore, it is more reasonable to assess heritability that considers the genotype environment interaction variation for the overall variance when predicting genetic advance as a result of selection [11-15]. It is necessary to identify the stable genotypes suitable for wide range of environmental conditions [16-20].

Different yield-contributing features are associated with grain yield in Indian mustard. Additionally, these characteristics are related to one another [21,22]. Through path coefficient, the intricate web of such a relationship is further simplified for analysis [23,24]. The path coefficient breaks the correlation coefficient of the yield with its contributing traits into direct and indirect effects [25]. Estimates of genetic variability, heritability and genetic advance facilitated selection may help to devise efficient selection criteria [26]. The presence of high-genetic diversity in the germplasm material provides the basis for plant breeding and developing new cultivars and assists in selecting desirable agronomic traits [27]. The extent and amount of genetic diversity present among available germplasm are of greatest importance while formulating any breeding program [28,29]. The loss of genetic diversity is a universal phenomenon for all crops and almost all the cultivated varieties of Indian mustard have a narrow genetic base [30,31]. Further, the identification and selection of genetically diverse parents are the most vital criteria for hybrid breeding programmes [32].

Genetic diversity among individuals or populations can be determined using morphological [33-36], biochemical [2,37,38], and molecular approaches [39-54]. It is now possible to quantify the magnitude of diversity among germplasm for use in breeding programme evaluation by employing biometrical methods like multivariate analysis [55] based on Mahalanobis [56], D² statistics, and Ward's no-hierarchical squared Euclidean distance method.

The present study was conducted to accomplish the presence of the genetic variability and diversity among various Indian mustard genotypes for yield accrediting traits.

MATERIALS AND METHODS

The current investigation was undertaken on a total of 75 Indian mustard genotypes (Table 1) acquired from the Zonal Agricultural Research Station, Morena, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya (RVSKVV), Gwalior, India (AICRP on Rapeseed and Mustard). All the genotypes were grown in randomized block design with two replications in Rabi 2021 at the experimental field of Department of Genetics & Plant Breeding, College of Agriculture, RVSKVV, Gwalior, India. Each genotype was planted in a plot of one row of 2-meter length with an arrangement of 30 cm apart between rows and 15 cm plant to plant. The crop was enchanted with protective irrigations and recommended packages of practices right through the growing season. Five arbitrarily chosen plants from each treatment were marked for taking the observations for the parameters, viz., plant height (cm), numbers of primary and secondary branches per plant, days to 50 percent flowering, days to maturity, length of the main raceme (cm), numbers of silique per the main raceme, numbers of silique per plant, silique length (cm), test weight (g), seed yield per plant (g), biological yield and harvest index (%) for analysis of mean performance.

The mean values of each genotype were employed for statistical analysis. Genotypic (GCV) and phenotypic coefficient of variation (PCV) was computed as per formula given by Burton [57]; heritability in the broad sense (h^2) as proposed by Burton and De [58] and genetic advance as per the method designated by Johnson *et al.* [59]. Genotypic and phenotypic correlations were calculated by way of the formula described by Weber and Moorthy [60] and Miller [61]. The portion of direct and indirect donations of different traits to the total correlation coefficients with yield was evaluated through path coefficient analysis as suggested by Wright [62,63] and suggested by Dewey and Lu [64]. The genetic divergence was estimated using Mahalanobis D^2 statistic following Rao [55] Inter and intra-cluster distances were calculated by Tocher's method as suggested by Rao [55] to form the clusters.

RESULTS AND DISCUSSION

Prediction of genetic variability in a crop is a goal since hybrids between lines of diverse

backgrounds usually; exhibit better heterosis than those between closely related parents [12,13]. Selection of genetically dissimilar parents for hybridization is a main aim of any crop improvement programme to accomplish anticipated segregates. The analysis of variance revealed considerable genetic differences for each of the investigated traits advised that there was genetic divergence among genotypes (Table 2). This demonstrates that the available gene pool for yields and its components has enough selection space for promising lines. For most of the traits, for instance days to 50% flowering, days to maturity, plant height (cm), number(s) of primary and secondary branches per plant, length of main raceme(cm), number(s) of siliquae on main raceme, test weight (g), siliquae length(cm), seed yield per plant(g), harvest index and biological yield, a vast range of variation was documented. Thus, it suggests that there is a lot of scope for choosing various quantitative traits to improve Indian mustard. Estimates of PCV and GCV showed greater values for traits such as seed yield per plant(g), biological yield, number(s) of siliquae on main raceme and number(s) of secondary branches per plant (Table 3). Similar results for many characteristics of Indian mustard were also addressed by Singh [65] and Yadav *et al.* [66].

Like any crop, in Indian mustard, seed yield is highly variable and complex due to an array of interrelated contributing characters. Direct selection for yield may therefore be ineffective. When examining the genetic basis of the relationship between two qualities, Falconer [67] argued that pleiotropy or full linkage could be answerable for the linear association. When a gene has pleiotropy or linkage, it has a general influence on both aspects (positive correlation), but other genes tend to boost one feature while decreasing the other (negative correlation). In the present experimentation, the phenotypic and genotypic correlation coefficients between different traits were computed (Table 4; Table 5). Genotypic correlation coefficients were found to be higher than phenotypic correlation coefficients for most of the traits, signifying a substantial inherent link between several traits that was masked by environmental factors in terms of phenotypic representation. Comparable outcomes were reported by Larik and Rajput [68], Baghel *et al.* [40], Rajpoot *et al.* [46], Verma *et al.* [50] in Indian mustard.

The genotypic and phenotypic associations between seed yield per plant and the length of the main raceme and the numbers of siliquae on it were both significantly and positively ($r=0.575$ and $r=0.574$, respectively) correlated. Therefore, selection based on any of these traits, alone or in combination, will lead to the discovery of genotypes with higher yields. These results are in accordance with findings of Khan *et al.* [69], who found that plant height, dry matter yield,

numbers of seeds per siliquae, days to flowering, raceme length, and test weight were significantly and positively correlated with seed yield. The oil content, plant height, days to flowering, days to maturity, numbers of siliquae per plant, numbers of seeds per siliquae and 1000-seed weight were found to have positive and significant correlations with seed yield in the experiment of Sandhu and Gupta [70]. According to Kumar *et al.* [71], seed yield was positively connected with plant height, numbers of primary and secondary branches per plant, numbers of siliquae on the main shoot and the numbers of siliquae per plant with 1,000-seed weight and harvest index. Singh [65] found a positive link between seed yield per plant and numbers of branches per plant, height of the plant, numbers of siliquae on the main raceme, length of the main raceme, number of seeds per siliquae and weight of the seeds. Chaudhary *et al.* [72] found substantial and positive association between the numbers of seeds produced per plant and the numbers of siliquae per plant, biological yield, oil content and harvest index. Seed yield was significantly and positively correlated with days to 50% flowering, days to maturity, plant height, numbers of primary and secondary branches, length of main shoot, numbers of siliquae on main shoot, numbers of seeds per siliquae and 1000-seed weight in investigation of Kumar *et al.* [73]. Gupta *et al.* [74] observed that plant height, harvest index, days to maturity, numbers of siliquae on the main axis, 1000-seed weight and numbers of principal branches were all positively and substantially correlated with seed yield. Moreover, Baghel *et al.* [40] investigated that majority of the traits and the seed yield per plant had considerable and positive correlations with phenotypic correlation coefficients being greater than genotypic correlation coefficients. According to Singh *et al.* [35], there is a positive and significant correlation between plant height, numbers of seeds per siliquae, weight of 1000 seeds, oil content, numbers of days to flowering, numbers of days to maturity, the length of the major raceme, and biological yield.

The breeder would be able to select the breeding tactic to be employed based on the estimates of genotypic and phenotypic correlations so that the advantageous correlation could be utilized and the unfavorable ones adjusted by creating new variability to generate new recombinants [75]. Recurrent selection programmes have been revealed to break undesirable correlations by Miller [61], whereas a breeding programme that uses yield, plant or field as the selection method would be beneficial if physical features are strongly associated with yield. Thus, it became clear from correlation studies in the current study that seven traits *viz.*, days to 50% flowering, numbers of secondary branches per plant, length of the main raceme, numbers of

siliquae on the main raceme, test weight, biological yield and seed yield per plant are crucial for improving Indian mustard. Furthermore, two of these traits *i. e.*, length of the main raceme and numbers of siliquae on the main raceme have the highest correlation coefficient, making them the most significant characters.

Although correlation coefficients are helpful in identifying the relationships between several characters on complex attributes like yield, nevertheless, such investigations may not deliver a precise depiction of the relative significant of direct and indirect effects of each contributing characters. Path coefficient analysis splits (Fig. 1) the correlation coefficient into the direct and indirect effects of a set of independent variables on the dependent variables. The length of the main raceme, days to 50% flowering, thousand seed weight, numbers of secondary branches per plant and biological yield towards seed yield, all disclosed substantial positive direct donations from siliquae in the results of the genotypic and phenotypic path coefficient (Table 6; Table 7). Days to flowering, maturity, plant height, oil content, test weight, and major branches all have a direct impact on seed yield [76]. Numbers of primary branches, days to flowering, 1000-seed weight, and numbers of seeds per siliquae were found to have a direct impact on seed output. Comparable results were also documented by Gangapur *et al.* [77]. Badra [78] also investigated an indirect relationship between plant height, seeds per siliquae, primary branches, siliquae per plant and days to maturity. Indirect effects on siliquae per plant, plant height, branches per plant, seeds per siliquae, oil content, test weight, and days to flowering were noted by Akbar *et al.* [23] *via* biological yield. Most characteristics had a direct effect on seed yield and an indirect effect on the numbers of days to 50% blooming in the experiment. Biological yield per plant revealed the trait's highest indirect influence, while days to 50% flowering showed the most beneficial direct effect in the research of Devi [79].

The percent contribution of thirteen different characters towards the expression of genetic divergence revealed that numbers of siliquae on main raceme contributed maximum divergence (37.63%) tracked by days to maturity, biological yield, plant height (cm), length of main raceme (cm), seed yield per plant (g), days to 50% flowering, harvest index, test weight (g), numbers of seeds per siliquae and numbers of primary branches per plant. Whilst the numbers of secondary branches per plant (0.04%) and siliquae length (cm) (0.04%) donated lowest in the genetic divergence (Table 8). These findings are parallel with the results of Mahto [80].

Crop species use their understanding of genetic divergence to choose their parents since this idea facilitates the differentiation of well-defined populations [81]. A potent approach for assessing genetic divergence among the choices from the same geographic region is the Mahalanobis D^2 analysis of quantitative traits. Seventy-five genotypes, included in this investigation were grouped into 6 clusters. A total of 53 genotypes fell into cluster I, 14 genotypes in cluster II, 5 genotypes in cluster IV and 1 each in cluster III, V and VI (Table 9). Similar outcomes were also seen in studies of Ghosh and Gulati [82], Goswami and Behl [83], Malik *et al.* [84], Doddabhimappa *et al.* [85], Goyat *et al.* [86], Kumar *et al.* [87], Shekhawat *et al.* [88], Singh *et al.* [89] and Nandi *et al.* [90].

The intra cluster divergence was found to range between 22.17 for cluster I and 30.85 for cluster IV. It is ranged from 0.00 to 30.85. Cluster IV showed maximum intra cluster D^2 value ($D^2 = 30.85$), cluster II ($D^2 = 24.32$) and cluster I ($D^2 = 22.17$) whereas clusters III and IV showed zero value for Intra cluster distance (Table 9). The substantial genetic divergence of the genotypes led to the highest intra-cluster dispersion. There is virtually little probability of creating desirable types by removing genotypes from clusters that are like them and have low intra-cluster divergence values [89]. Therefore, it might make sense to test crossovers between types from clusters that are farther apart than usual. In order to further produce high yielding Indian mustard varieties, the little diversity and selection of parents within the cluster having a higher mean for a certain characteristic may be helpful.

Maximum inter cluster D^2 value (56.23) was recorded between clusters II and VI, whereas the minimum average inter cluster D^2 value (27.97) was recorded between cluster I and III (Fig. 2). The maximum inter cluster D^2 value indicated that varieties of cluster III and IV are not so closely related whereas the varieties of cluster I and III are closely related (Table 10). It is apparent therefore; the varieties of various clusters differ so significantly with regards to their relative genetic distance as indicated from the high variation of D^2 values. This makes it clear that the genotypes included in these clusters have a wide range of genetic diversity and may be used in a mustard hybridization programme to increase seed yield. Maximum cluster mean estimated for number of siliquae on main raceme cluster III showed maximum value (222.50) and cluster V showed minimum value (71.50), length of main raceme cluster IV showed maximum (117.40) and cluster VI had minimum (72.50) cluster mean values, while for seed yield per plant was found maximum in cluster IV (21.45) and while minimum value was noticed

for cluster I (9.03) (Table 11). These findings confirm in earlier study of Ghosh and Gulati [82]. Based on these traits superior genotypes are selected and used in hybridization programme as a donor parent. Inter-crossing of genotypes involved in these clusters could be practiced for inducing variability in the respective characters and their rationale improvement for increasing seed yield.

Conclusion

The study well highlighted the presence of the genetic variability and diversity among various Indian mustard genotypes for yield accrediting traits. The genotypes of various clusters differ significantly in terms of their relative genetic distance, as evidenced by the wide range of D2 values. This shows that the genotypes present in these clusters have a wide range of genetic variability and might be applied in a mustard hybridization programme to create cultivars with higher yields.

REFERENCES

1. Yang J, Liu D, Wang X. *et al.* The genome sequence of allopolyploid *Brassica juncea* and analysis of differential homoeolog gene expression influencing selection. *Nat Genet* 2016; 48: 1225–1232. <https://doi.org/10.1038/ng.3657>
2. Shyam C, Tripathi MK. Biochemical studies in Indian mustard [*Brassica juncea* (Linn) Czern & Coss] for fatty acid profiling. *Int. J. Chem. Stud.* 2019;7(4):338-343.
3. Thakur AK, Parmar N, Singh KH, Nanjundan J. Current achievements and future prospects of genetic engineering in Indian mustard (*Brassica juncea* L. Czern & Coss.). *Planta* 2020; 252:56.
4. Egorova KV, Sinyavina NG, Artemyeva AM, Kocherina NV, Chesnokov YV. QTL Analysis of the content of some bioactive compounds in *Brassica rapa* L. grown under light culture conditions. *Horticulturae* 2021, 7:583.
5. Treccarichi S, Rizzo GF, Branca F. In vitro culture for micropropagation, somatic embryogenesis, somatic mutation, and somatic hybridization in *Brassica juncea*. In: Kole C., Mohapatra T. (eds) *The Brassica juncea Genome. Compendium of Plant Genomes.* 2022; pp 159-169.

6. Shyam C, Tripathi M K, Tiwari S, Ahuja A, Tripathi N, Gupta N. *In vitro* regeneration from callus and cell suspension cultures in Indian mustard [*Brassica juncea* (Linn.) Czern & Coss] International Journal of Agricultural Technology. 2021a;17(3):1095-1112.
7. Shyam C, Tripathi MK, Tiwari S, Tripathi N, Solanki RS, Sapre S, Ahuja A, Tiwari S. *In vitro* production of somaclones with decreased erucic acid content in Indian mustard [*Brassica juncea* (Linn.) Czern & Coss. Plants. 2021b; 10:1297.
8. Shyam C, Tripathi M K, Tiwari S, Ahuja A, Tripathi N, Gupta N. Plant regeneration in Indian mustard [*Brassica juncea* (Linn.) Czern & Coss]: Experimental investigation. In book: Current Topics in Agricultural Sciences. 2021c; 3 :120-135.
9. Yadav DK, Yadav R, Vishwakarma H, Yashpal, Yadav S, Saini N, Vasudev S. Genetic diversity characterization and population structure in *Brassica juncea*. In: Kole C., Mohapatra T. (eds) The Brassica juncea Genome. Compendium of Plant Genomes. 2022; pp 73-84.
10. Central Organization for Oil Industry & Trade, 2022
11. Yadav LN, Tripathi MK, Sikarwar RS, Mishra AK. Heterosis in sesame. Sesame & Safflower Newsl. 2005; 20:6-10.
12. Tripathi MK, Tomar SS, Tiwari VK, Awasthi D, Gupta JC. Heterosis in Indian mustard [*Brassica juncea* (L) Czern and Coss]. Prog. Res.;2015; 10 (Special-VI):3376-3379.
13. Barfa D, Tripathi MK, Kandalkar VS, Gupta JC, Kumar G. Heterosis and combining ability analysis for seed yield in Indian mustard [*Brassica Juncea* (L) Czern & Coss]. Suppl 2017; 75-83.
14. Shyam C, Tripathi MK, Tiwari S, Tripathi N. Genetic components, and diversity analysis in Indian mustard [*Brassica juncea* (Linn.) Czern & Coss] based on different morpho-physiological traits. Current Journal of Applied Science and Technology. 2021d; 40 (20):34-57.
15. Shyam C, Tripathi MK, Tiwari S, Tripathi N, Sikarwar RS. Morpho-physiological variations and genetic components analysis in *Brassica juncea* (Linn.) Czern & Coss. In book: Research Developments in Science and Technology, 2022a;1:98-126.
16. Wu JG, Shi CH, Zhang HZ. Partitioning genetic effects due to embryo, cytoplasm and maternal parent for oil content in oilseed rape (*Brassica napus* L.). Genet. Mol. Biol. 2006; 29(3): 57-89.

17. Escobar M, Berti M, Matus I, Tapia M, Johnson B (2011). Genotype Environment Interaction in Canola (*Brassica napus* L.) Seed Yield in Chile. *Chilean J. Agric. Res.* 2011; 71
18. Jeromela AM, Nagl N, Varga JG, Hristov N, Spika AN, Vasicand MM, Marinkovic R. Genotype by environment interaction for seed yield per plant in rapeseed using AMMI model. *Pesq. Agropec. Bras.* 2011; 46 (2).
19. Zhang H, Berger JD, Milroy SP. Genotype \times environment interaction studies highlight the role of phenology in specific adaptation of canola (*Brassica napus*) to contrasting Mediterranean climates. *Field Crops Res.* 2013; 144: 77–88.
20. Shyam C, Tripathi MK, Tripathi N, Tiwari S, Sikarwar RS. Genetic variations in fatty acids and oil compositions among 188 Indian mustard *Brassica juncea* (Linn.) Czern & Coss genotypes. *Curr J Appl Sci Technol.* 2022b; 40(46):9-28.
21. Ali N, Javidfar F, Elmira JY, Mirza MY. Relationship among yield components and selection criteria for yield improvement in winter rapeseed (*Brassica napus* L.). *Pak. J. Bot.*, 2003; 35(2): 167-174.
22. Khan RSAK, Khan FA. Evaluation of genetic potential of some Brassica germplasm collections. *Int. J. Agri. Biol.*, 2003; 5(4): 630-631.
23. Akbar M, Mahmood T, Yaqub M, Anwar M, Ali M, Iqbal N. Variability, correlation and path coefficient studies in summer mustard (*B. juncea* L.). *Asian J. Plant Sci.*, 2003; 2(9): 696-698.
24. Akbar M, Saleem U, Tahira M, Yaqub, Iqbal N. Utilization of genetic variability, correlation and path analysis for seed yield improvement in mustard, *B. juncea* L. *J. Agric. Res.*, 2007; 45(1): 25-31.
25. Aytacı Z, Kinacı G. Genetic variability and association studies of some quantitative characters in winter rapeseed (*B. napus* L.). *Afr. J. Biotechnol.*, 2009; 8(15): 3547-3554.
26. Sadat HA, Nematzadeh GA, Jelodar NB, Chapi OG. Genetic evaluation of yield and yield components at advanced generations in rapeseed (*B. napus* L.). *Afr. J. Agric. Res.*, 2010; 5(15): 1958-1964.
27. Govindaraj R, Badra SR, Chaudhary L, Chaudhary L. Association of yield and its component in Indian mustard (*Brassica juncea* L. Czern and Coss). *Agriculture Science Digest*, 2015; 21(2): 83-86.

28. Sharma A, Sachan JN, Tewari SK. Genetic divergence for important quantitative traits in Indian mustard (*Brassica juncea* (L.) Czern and Coss). *Agricultural Science Digest*, 2020; 26, 269-272.
29. Tiwari N, Tiwari S, Tripathi N. Genetic characterization of Indian little millet (*Panicum sumatrense*) genotypes using Random Amplified Polymorphic DNA markers. *Agriculture and Natural Resources*. 2018; 52(4): 347-352.
30. Srivastava MK, Singh BP. Correlation and path analysis in Indian mustard (*Brassica juncea* L. Czern and Coss). *Crop Res. Hisar*, 2002; 23(3): 517-521.
31. Chauhan JS, Bhadauria VPS, Singh KH, Singh M, Kumar A. Genetic diversity analysis in rapeseed-mustard using quality characters. *Annals of Arid Zone*, 2008; 47(2): 145-149.
32. Banga R, Ranjan N, Chakraborty, Imam, Z. Divergence studies in Indian mustard. *Soil and Crops*, 2015; 14(2): 297- 304.
33. Vaishnava A, Sachan JN, Tewari SK. Genetic divergence for important quantitative traits in Indian mustard (*Brassica juncea* (L.) Czern and Coss). *Agricultural Science Digest*, 2006; 26, 269-272.
34. Alie FA, Singh T, Sharma PK. Genetic diversity analysis in Indian mustard [*Brassica juncea* (L.) Czern & Coss]. *Progressive Agriculture-An International Journal*, 2009; 9, 50-53.
35. Singh D, Arya RK, Chandra N, Niwas R, Salisbury P. Genetic diversity studies in relation to seed yield and its component traits in Indian mustard (*Brassica juncea* L. Czern & Coss.). *Journal of Oilseeds Brassica*, 2010; 1: 19-22.
36. Mishra N, Tripathi MK, Tiwari S, Tripathi N, Trivedi HK. Morphological and Molecular Screening of Soybean Genotypes against Yellow Mosaic Virus Disease. *LegumeResearch*;2020.DOI:10.18805/LR-4240
37. Sharma A, Tripathi MK, Tiwari S, Gupta N, Tripathi N, Mishra N. Evaluation of soybean (*Glycine max* L.) genotypes on the basis of biochemical contents and anti-oxidant enzyme activities. *Legume Res*. 2021; 44, LR-467.
38. Shyam C, Tripathi MK, Tripathi N, Tiwari S, Sikarwar RS. Analysis of genetic differences in fatty acids and oil contents among *Brassica juncea* (Linn.) Czern & Coss genotypes. In book: *Research Developments in Science and Technology*, 2022c; 1:127-149.

39. Mohammadi SA, Prasanna BM. Analysis of genetic diversity in crop plants salient statistical tools and considerations. Review & Interpretation. Crop Science, 2003; 43, 1235-1248.
40. Baghel R, Sharma AK, Tiwari S, Tripathi MK, Tripathi N. Genetic diversity analysis of Indian mustard (*Brassica spp.*) germplasm lines using SSR molecular markers. Int J Curr Microbiol App Sci. 2020; 9 (12): 137-143.
41. Rajpoot NS, Tripathi MK, Tiwari S, Tomar RS, Kandalkar VS. Characterization of Indian mustard germplasm on the basis of morphological traits and SSR markers. Curr J Appl Sci Technol. 2020; 39:300-311.
42. Shyam C, Tripathi MK, Tiwari S, Tripathi N, Ahuja A. Molecular characterization and identification of *Brassica* genotype(s) for low and high erucic acid content using SSR markers. Global J Biosci Biotechnol. 2020; 9(2):56-66.
43. Makwana K, Tiwari, S, Tripathi MK, Sharma A K, Pandya RK, Singh A K. Morphological characterization and DNA finger printing of pearl millet (*Pennisetum Glaucum* (L.) germplasms. Range Management and Agroforestry, 2021; 42(2):205-211
44. Pramanik A, Tiwari S, Tripathi MK, Mandloi S, Tomar RS. Identification of groundnut germplasm lines for foliar disease resistance and high oleic traits using SNP and gene-based markers and their morphological characterization. Legume Research; 2021.DOI: 10.18805/LR-4666
45. Singh L, Nanjundan J, Singh KH, Sharma D, Parmar N, Watts A, Jain R, Thakur AK. Development of a set of SSR markers for characterization of Indian mustard germplasm and varieties. Journal of Plant Biochemistry and Biotechnology. 2021; 9:1-1
46. Verma K, Tripathi MK, Tiwari S, Tripathi N. Analysis of genetic diversity among *Brassica juncea* genotypes using morpho-physiological and SSR markers. Int J Curr Microbiol App Sci. 2021; 10(01):1108-1117.
47. Asati R, Tripathi MK, Tiwari S, Yadav RK and Tripathi N. Molecular breeding and drought tolerance in chickpea. Life 2022;12, 1846. <https://doi.org/10.3390/life12111846>
48. Mishra N, Tripathi MK, Tripathi N, Tiwari S, Gupta N, Sharma, A. Screening of soybean genotypes against drought on the basis of gene-linked microsatellite markers. In book: Innovations in Science and Technology, 2022a; 3:49-61

49. Mishra N, Tripathi MK, Tiwari S, Tripathi N, Gupta N, Sharma A, Solanki RS. Characterization of soybean genotypes on the basis of yield attributing traits and SSR molecular markers. In book: Innovations in Science and Technology, 2022b;. 3:87-106.
50. Rajpoot NS, Tripathi MK, Tiwari S, Tomar RS, Tripathi N, Sikarwar RS, Tomar SS. Morphological and molecular characterization of Indian mustard germplasm lines. Research Developments in Science and Technology, 2022; 4: 151-165.
51. Shyam C, Tripathi MK, Tripathi N, Tiwari S, Sikarwar RS. Identification of low and high erucic acid containing genotype(S) in Indian mustard employing molecular markers. In book: Recent Progress in Plant and Soil Research, 2022d; 5: 18-36
52. Tomar YS, Tiwari S, Tripathi MK, Singh S, Gupta N. Genetic Diversity, Population Structure and Biochemical Parameters Estimations Driving Variations in Groundnut Germplasm. Legume Research, 2022; DOI: 10.18805/LR-4965.
53. Tripathi N, Tripathi MK, Tiwari S, Payasi DK. Molecular breeding to overcome biotic stresses in soybean: update. Plants (Basel). 2022; 11(15):1967.
54. Tripathi MK, Tripathi N, Tiwari S, Mishra N, Sharma A, Tiwari S, Singh S. Identification of Indian soybean (*Glycine max* [L.] Merr.) genotypes for drought tolerance and genetic diversity analysis using SSR markers SCI- 2023; 3(3);2023;31-46
55. Rao CR. Advanced Statistical Methods in Biometrics Research. John Willey and Sons, New York. 1952.
56. Mahalanobis PC. On the generalized distance in statistics. Proc. Natl. Acad. Sci. India, 1956; 2: 9-55.
57. Burton GW. Quantitative inheritance in grasses. Proced. 6th Int. Grassland Cong. 1952; 1:127-183.16.
58. Burton GW, De V. Estimating heritability in tall Fescue from replicated clonal material. Agron. J.1953; 45:475-481.17.
59. Johnson HW, Robinson HF, Comstock RE. Estimates of genetic and environmental variability in wheat. Agron. J. 1955; 47:314-318
60. Webber CR, Moorthy BR. Heritable and non-heritable relationship and variability of oil content and agronomic characteristics in the F generation of soybean crosses. J. Agron. 1952; 44:202-209.19.

61. Miller PA, Williams JE, Robinson HF, Comstock R E. Estimates of variance and covariance in upland cotton and their implications in selection. *Agron. J.* 1958; 50:126-131.20.
62. Wright S. Correlation and causation. *J. Agril. Res.* 1921; 20:557-587.21.
63. Wright S. The analysis of variance and the correlations between relatives with respect to deviations from an optimum. *J. Genet.* 1935; 30:243-256.22.
64. Dewey DR, Lu KHA. Correlation and path coefficient analysis of component in crested wheatgrass seed production. *Agron. J.* 1959; 51:515-518.
65. Singh B. Character association and path analysis under dryland condition in India mustard (*B. juncea*). *Cruciferae Newslet.*, 2004; 25: 99-100.
66. Yadav DK, Giri SC, Vignesh M, Sujata Vasudev, Yadav AK, Dass B, Singh R, Singh N, Mohapatra T, Prabhu KV. Genetic variability and trait association studies in Indian mustard (*Brassica juncea*). *Indian J. of Agric. Sci.*, 2011; 81(8): 712-716.
67. Falconer DS. Introduction to quantitative genetics. Oliver and Boyd Ltd., Edinburgh and London. 1960.
68. Larik AS, Rajput LS. Estimation of selection parameters in *Brassica juncea*. *J. of Botany*, 2000; 32(2): 323-330.
69. Khan MH, Ali MM, Vhuiyan SR, Mahmud F. Genetic divergence in rapeseed-mustard (*Brassica rapa* L.). *Bangladesh Journal of Agricultural Research*, 2013; 38(3):417-423.
70. Sandhu SK, Gupta NP. Genetic divergence and correlation studies in Brassica species. *Crop Improv.*, 1996; 23(2): 253-256.
71. Kumar S, Sangawan RS, Yadav IS, Kumar D. Correlation studies in Brassica species under dryland condition. *Annals Agric-Bio Res.*, 1999; 4(1) 63-65.
72. Chaudhary MA, Raheman LM, Rahem AK. Correlation and path coefficient analysis of nine important characters of mustard. *J. Agric.* 2003; 12(3): 149-154.
73. Kumar M, Singh KP, Singh D. Genetic divergence analysis for seed yield, its parameters and oil content in Indian mustard (*Brassica juncea* (L.) Czern & Coss). *Res. Crops*, 2007; 8: 183-185.
74. Gupta P, Chaudhary HB, Sirohi SPS, Lal SK, Gaurav SS. Correlation association and path coefficient analysis of different parameters affecting seed yield and its component in

- Indian mustard (*Brassica juncea* L. Czern and Coss). Progressive Agriculture, 2010; 10(3): 154-156.
75. Mishra N, Tripathi MK, Tiwari S, Tripathi N, Gupta N, Sharma A. Morphological and physiological performance of Indian soybean [*Glycine max* (L.) Merrill] genotypes in respect to drought. Legume Research; 2021. DOI: 10.18805/LR-4550
76. Kumar N, Shrivastava S. Path analysis in late sown Indian mustard (*Brassica juncea* (L) Czern and Coss]. J. Applied Biol., 2000; 10(2): 131-135.
77. Gangapur DR, Prakash BG, Salimath PM, Kumar R, Rao MS. Correlation and path analysis in Indian mustard (*Brassica juncea* L. Czern and Coss). Karnataka J. of Agri. Sci., 2009; 22(5): 971-977.
78. Badra SR, Chaudhary L, Chaudhary L. Association of yield and its component in Indian mustard (*Brassica juncea* L. Czern and Coss). Agriculture Science Digest, 2001; 21(2): 83-86.
79. Devi B. Correlation and path analysis in Indian mustard (*Brassica juncea* L.) in agro-climatic conditions of Jhansi (U.P.). J. Pharmaco. and Phytochem. 2008; 7(1): 1678-1681.
80. Mahto JL. Genetic divergence and stability in Indian mustard under rained conditions. J. Maharashtra Agril. Univ., 1966; 21(3): 334-337.
81. Aunwinithul, Patil S, Charjan SU, Thakare PG, Wankhade M. Genetic divergence studies in Indian mustard. Soil and Crops, 2004; 14(2): 297-304.
82. Ghosh SK, Gulati SC. Parental diversity to realize maximum heterosis in Indian mustard (*Brassica juncea* L. Czern & Coss). Indian J. of Gen. and Pl. Breed., 2002; 62(1): 25-28.
83. Goswami PK, Behl RK. Genetic divergence in Indian mustard. Annals of Agricultural Research, 2006; 27(2): 187-190.
84. Malik S, Sirohi SPS, Yadav R, Sandeep. Genetic divergence in Indian mustard (*Brassica juncea* L. Czern and Coss). Plant Archives, 2006; 6(1): 381-382.
85. Doddabhimappa R, Gangapur B, Prakash G, Hiremath CP. Genetic diversity analysis of Indian mustard (*Brassica juncea*). Electronic Journal of Plant Breeding, 2010; 1(4): 407-413.

86. Goyal B, Singh D, Avatar R, Singh A. Estimation of selection parameters in the elite gene pool of Indian mustard [*Brassica juncea* (L.) Czern & Coss.]. Res. on Crops, 2012; 13(3): 1098-1101.
87. Kumar V, Chand P, Bahuguna DK, Tyagi N, Kumar S. A study of character association and genetic divergence in Indian mustard (*Brassica juncea* (L.) Czern & Coss). Progressive Agriculture 2003; 13(1):187-192.
88. Shekhawat N, Jadeja GC, Singh J. Genetic variability for yield and its components in Indian mustard (*Brassica juncea* L. Czern & Coss). Electronic Journal of Plant Breeding, 2014; 5(1):117-119.
89. Singh L, Nanjundan J, Singh KH, Sharma D, Parmar N, Watts A, Jain R, Thakur AK. Development of a set of SSR markers for characterization of Indian mustard germplasm and varieties. Journal of Plant Biochemistry and Biotechnology. 2021 Nov 9:1-1
90. Nandi S, Rout S, Sur B, Chakraborty M, Hijam L, Roy SK, Gupta A. Assessment of Genetic Parameters, Trait Association and Diversity Analysis in Indian Mustard [*Brassica juncea* (L.) Czern and Coss]. Biological Forum – An International Journal, 2021; 13(2): 95-109.

Table 1 List of mustard genotypes with their parentage/ source

S. No.	Genotypes	Parentage/ Sources
1.	RB-50	Laxmi X RH-9617
2.	Pusa Bold	Varuna x BIC1780
3.	Varuna	Selection from Varansi Local
4.	Rohini	Selection from natural population of Varuna
5.	Kranti	Selection for Varuna
6.	RH- 725	CCSHAU Hisar
7.	Maya	Varuna x KRV 11
8.	Vardan	Derived through biparental mating involving Varuna, Keshari, CSU 10 and IB 1775, IB 1786, IB 1866
9.	Vasundhara	RH 839 x RH 30
10.	Swarn Jyoti	Selection from germplasm line RC 1670
11.	Pusa Jagannath	Varuna x Synthetic juncea
12.	Pusa Jai Kisan	Somaclone of Varuna
13.	Albeli	ZARS, Morena, RVSKVV, Gwalior
14.	Sej-2	Derived from a cross of B. juncea to a amphidiploid
15.	Shraddha	ZARS, Morena, RVSKVV, Gwalior
16.	DMH 1	CMS based hybrid
17.	L-4	Canada
18.	JMWR-908-1	ZARS, Morena, RVSKVV, Gwalior
19.	RGN-73	RGN 8 x Pusa Bold
20.	NRC-HB-101	BL 4 x Pusa Bold
21.	NRC-HB-506	(MJA 05 x MJR 1)
22.	RVM-3	ZARS, Morena
23.	RH-749	RH-781 xRH-9617
24.	NRC DR-2	MDOC 43 x NBPGR 36
25.	DRMR IJ-31	HB 9908 x HB 9916
26.	China	ZARS, Morena, RVSKVV, Gwalior
27.	GSL-1	Punjab agricultural university, Ludhiana
28.	GSC-7	Punjab agricultural university, Ludhiana
29.	PC-5	ZARS, Morena, RVSKVV, Gwalior
30.	PC-6	ZARS, Morena, RVSKVV, Gwalior
31.	RP-9	ZARS, Morena, RVSKVV, Gwalior
32.	Kiran	ZARS, Morena, RVSKVV, Gwalior
33.	JTC-1	ZARS, Morena, RVSKVV, Gwalior
34.	JM-1	Pusa Bold x L 6
35.	JM-2	MutantofRL9
36.	JM-3	Varuna x YRT 3
37.	RVM-1	ZARS, Morena
38.	RVM-2	Selection from Chambal growing region
39.	PM-25	Sej-8 x Pusa Jagannath
40.	PM-26	VEJ Open x PusaAgrani
41.	PM-27	Derived from the cross [(Divya x Pusa Bold) x (PR 666EPS) x PR 704EPS-2 x B85)]
42.	PM-28	SEJ8 x PUSA JAGANNATH
43.	PM-30	Bio 902 x ZEM 1
44.	Pusa Vijay	Synthetic <i>Brassica juncea</i> x VSL 5
45.	JMM-927	ZARS, Morena, RVSKVV, Gwalior
46.	JMM-991	ZARS, Morena, RVSKVV, Gwalior
47.	RMM-10-01-01	ZARS, Morena, RVSKVV, Gwalior

48.	RMM-12-01-18	ZARS, Morena, RVSKVV, Gwalior
49.	RMM-12-03-18	ZARS, Morena, RVSKVV, Gwalior
50.	WRR-5	White rust resistant advance breeding lines, DRMR, Bharatpur
51.	WRR-6	White rust resistant advance breeding lines, DRMR, Bharatpur
52.	WRR-7	White rust resistant advance breeding lines, DRMR, Bharatpur
53.	WRR-8	White rust resistant advance breeding lines, DRMR, Bharatpur
54.	WRR-9	White rust resistant advance breeding lines, DRMR, Bharatpur
55.	WRR-10	White rust resistant advance breeding lines, DRMR, Bharatpur
56.	WRR-11	White rust resistant advance breeding lines, DRMR, Bharatpur
57.	WRR-12	White rust resistant advance breeding lines, DRMR, Bharatpur
58.	WRR-13	White rust resistant advance breeding lines, DRMR, Bharatpur
59.	WRR-14	White rust resistant advance breeding lines, DRMR, Bharatpur
60.	WRR-15	White rust resistant advance breeding lines, DRMR, Bharatpur
61.	WRR-16	White rust resistant advance breeding lines, DRMR, Bharatpur
62.	WRR-17	White rust resistant advance breeding lines, DRMR, Bharatpur
63.	WRR-18	White rust resistant advance breeding lines, Bharatpur
64.	WRR-19	White rust resistant advance breeding lines, DRMR, Bharatpur
65.	WRR-20	White rust resistant advance breeding lines, DRMR, Bharatpur
66.	WRR-21	White rust resistant advance breeding lines, DRMR, Bharatpur
67.	WRR-22	White rust resistant advance breeding lines, DRMR, Bharatpur
68.	WRR-25	White rust resistant advance breeding lines, DRMR, Bharatpur
69.	WRR-26	White rust resistant advance breeding lines, DRMR, Bharatpur
70.	WRR-27	White rust resistant advance breeding lines, DRMR, Bharatpur
71.	WRR-28	White rust resistant advance breeding lines, DRMR, Bharatpur
72.	WRR-29	White rust resistant advance breeding lines, DRMR, Bharatpur
73.	WRR-30	White rust resistant advance breeding lines, DRMR, Bharatpur
74.	WRR-31	White rust resistant advance breeding lines, DRMR, Bharatpur
75.	WRR-32	White rust resistant advance breeding lines, DRMR, Bharatpur

Table 2 Analysis of variance for different traits in Indian mustard

Source of Var.	Df	DF	DM	PH	PB	SB	LMR	SMR	S/SL	SL	TSW	BY	HI	SYPP
Replication	1	26.45	23.20	235.35	12.90	6.82	6.82	20.16	0.3260	0.5400	1.6030	37.5000	62.39	0.272
Treatment	74	125.39**	774.85**	2291.60**	5.684**	31.801**	31.801**	7094.7**	4.004**	0.637**	0.971**	543.1**	156.21**	35.229**
Error	74	3.73	4.949	51.105	2.501	3.218	3.218	32.49	1.65	0.137	0.081	5.864	8.577	0.323

DF= Days to 50% Flowering, DM= Days to maturity, PH= Plant height (cm), PB= Numbers of primary branches/ plants, SB= Numbers of secondary branches/ plants, LMR= Length of main raceme (cm), SMR= Numbers of siliquae on main raceme, S/SL= Numbers of seed per siliquae, SL= Siliquae length (cm), TSW= Test weight (g), BY= Biological yield, HI= Harvest Index, SSYP= Seed yield per plant (g)

Table 3 Estimation of GCV, PCV, heritability and genetic advance for different quantitative characters

Observations	Range		Grand Mean	Coefficient of variation		h ² (bs)	Gen. Adv as 5% of Mean
	Min	Max		GCV	PCV		
DF	41	75.5	52.5533	14.841	15.066	0.97	30.113
DM	106	184	145.6867	13.467	13.511	0.994	27.654
PH	93.34	245.84	178.7997	18.719	18.932	0.978	38.13
PB	4.5	13.5	8.32	15.163	20.263	0.56	23.374
SB	6	23	11.1067	34.037	35.902	0.899	66.474
LMR	59	142.5	89.5733	21.292	21.517	0.979	43.401
SMR	65	294	144.42	41.146	41.241	0.995	84.566
S/SL	11	18	12.7667	8.497	11.084	0.588	13.419
SL	3.75	6.5	4.98	10.037	11.339	0.784	18.304
TSW	3.68	6.25	4.8086	13.871	14.492	0.916	27.351
BY	16	95	35.26	46.482	46.735	0.989	95.234
HI	15.07	55.62	30.8091	27.887	28.685	0.945	55.847
SSYP	4.85	27.68	10.1947	40.979	41.168	0.991	84.029

Table 4 Estimation of correlation coefficient at genotypic level in Indian mustard

Characters	DF	DM	PH	PB	SB	LMR	SMR	S/SL	SL	TSW	BY	HI	SSYP
DF	1.000	0.0936	0.0774	0.0721	-0.0240	-0.1130	-0.1080	0.1887	0.0779	-0.011	0.3182*	-0.2860	0.0939
DM		1.0000	-0.0310	0.1287	-0.0380	-0.0620	0.0459	0.1853	0.1971	-0.039	0.0693	-0.3628*	-0.16
PH			1.0000	0.0302	-0.1360	0.1414	0.0584	0.1154	-0.086	0.1511	0.1225	-0.0730	0.0888
PB				1.0000	0.5699**	0.0981	0.4166**	0.2086	-0.069	0.192	0.6593**	-0.1700	0.5908**
SB					1.0000	0.3024*	0.2661	-0.14	-0.135	-0.049	0.3757*	0.1124	0.5405**
LMR						1.0000	0.0569	-0.071	-0.018	0.1958	0.1247	0.0764	0.2247
SMR							1.0000	0.2608	0.1284	0.1047	0.4849*	-0.0720	0.5882**
S/SL								1.0000	0.5946**	-0.016	0.1158	-0.2130	0.0154
SL									1.0000	-0.005	-0.128	0.0230	-0.1140
TSW										1.0000	0.2469	0.1752	0.2926
BY											1.0000	-0.4476*	0.7303**
HI												1.0000	0.2182
SSYP													1.0000

Table 5 Estimation of correlation coefficient at phenotypic level in Indian mustard

Characters	DF	DM	PH	PB	SB	LMR	SMR	S/SL	SL	TSW	BY	HI	SYPP
DF	1.0000	0.0949	0.0765	0.0640	-0.0220	-0.1120	-0.1060	0.1356	0.0690	-0.0160	0.3155*	-0.2740*	0.0960
DM		1.0000	-0.0330	0.0957	-0.0330	-0.0600	0.0448	0.1417	0.1832	-0.0350	0.0686	-0.3510**	-0.1590
PH			1.0000	0.0121	-0.1360	0.1360	0.0581	0.0945	-0.0670	0.1462	0.1155	-0.0660	0.0858
PB				1.0000	0.4125**	0.0877	0.3085**	0.1691	-0.015	0.1354	0.5027**	-0.1370	0.4422**
SB					1.0000	0.2888*	0.2519*	-0.1120	-0.099	-0.0480	0.3587**	0.0949	0.5040**
LMR						1.0000	0.0569	-0.0530	-0.014	0.1852	0.1204	0.0789	0.2223
SMR							1.0000	0.2038	0.1115	0.0993	0.4823**	-0.0700	0.5848**
S/SL								1.0000	0.4444**	-0.0180	0.0874	-0.1610	0.0110
SL									1.0000	-0.0160	-0.1210	0.0235	-0.1080
TSW										1.0000	0.2398*	0.1651	0.2829*
BY											1.0000	-0.4480**	0.7247**
HI												1.0000	0.2218
SYPP													1.0000

DF= Days to 50% Flowering, DM= Days to maturity, PH= Plant height (cm), PB= Numbers of primary branches/ plants, SB= Numbers of secondary branches/ plants, LMR= Length of main raceme (cm), SMR= Numbers of siliquae on main raceme, S/SL= Numbers of seed per siliquae, SL= Siliquae length (cm), TSW= Test weight (g), BY= Biological yield, HI= Harvest Index, SSYP= Seed yield per plant (g)

Table 6 Path coefficient analysis showing the direct and indirect effect of different characters on the seed yield at genotypic level in Indian mustard

Character	DF	DM	PH	PB	SB	LMR	SMR	S/SL	SL	TSW	BY	HI
DF	0.0080	0.0008	0.0006	0.0006	-	-	-	0.0015	0.0006	-	0.0026	-
DM	0.0016	0.0166	-	0.0021	-	-	0.0008	0.0031	0.0033	-	0.0012	-
PH	0.0014	-	0.0176	0.0005	-	0.0025	0.0010	0.0020	-	0.0027	0.0022	-
PB	-	-	-	-	-	-	-	-	0.0023	-	-	0.0057
SB	-	-	-	0.0424	0.0744	0.0225	0.0198	-	-	-	0.0279	0.0084
LMR	-	-	0.0064	0.0044	0.0136	0.0450	0.0026	-	-	0.0088	0.0056	0.0034
SMR	-	0.0082	0.0104	0.0745	0.0476	0.0102	0.1787	0.0466	0.0230	0.0187	0.0867	-
S/SL	0.0094	0.0092	0.0057	0.0104	-	-	0.0129	0.0496	0.0295	-	0.0057	-
SL	-	-	0.0050	0.0040	0.0078	0.0010	-	-	-	0.0003	0.0074	-
TSW	0.0008	0.0028	-	-	0.0035	-	-	0.0011	0.0003	-	-	-
BY	0.2951	0.0642	0.1136	0.6115	0.3484	0.1156	0.4497	0.1074	-	0.2290	0.9274	-
HI	-	-	-	-	0.0745	0.0507	-	-	0.0153	0.1161	-	0.6628
SYPP	0.0939	-	0.0888	0.5908	0.5405	0.2247	0.5882	0.0154	-	0.2926	0.7303	0.2182

Table 7 Path coefficient analysis showing the direct and indirect effect of different characters on the seed yield at phenotypic level in Indian mustard

Character	DF	DM	PH	PB	SB	LMR	SMR	S/SL	SL	TSW	BY	HI
DF	0.0109	0.0010	0.0008	0.0007	-	-	-	0.001	0.000	-	0.003	-
DM	0.0006	0.0064	-	0.0006	-	-	0.000	0.000	0.001	-	0.000	-
PH	0.0015	-	0.0196	0.0002	-	0.002	0.001	0.001	-	0.002	0.002	-
PB	-	-	-	-	-	-	-	-	0.000	-	-	0.000
SB	-	-	-	0.0231	0.0559	0.016	0.014	-	-	-	0.020	0.005
LMR	-	-	0.0064	0.0042	0.0137	0.047	0.002	-	-	0.008	0.005	0.003
SMR	-	0.0082	0.0106	0.0565	0.0461	0.010	0.183	0.037	0.020	0.018	0.088	-

	0.0194					4	0	3	4	2	3	0.0129
S/SL	0.0028	0.0029	0.0020	0.0035	- 0.0023	- 0.0011	0.0042	0.0207	0.0092	- 0.0004	0.0018	- 0.0033
SL	- 0.0026	- 0.0070	0.0026	0.0006	0.0038	0.0005	0.0043	0.0170	0.0382	0.0006	0.0046	- 0.0009
TSW	0.0011	0.0024	- 0.0102	- 0.0095	0.0034	- 0.0130	- 0.0070	0.0012	0.0011	- 0.0701	- 0.0168	- 0.0116
BY	0.2876	0.0625	0.1053	0.4584	0.3271	0.1098	0.4398	0.0797	- 0.1107	0.2187	0.9117	- 0.4084
HI	- 0.1796	- 0.2298	- 0.0434	- 0.0895	0.0622	0.0517	- 0.0460	- 0.1053	0.0154	0.1082	- 0.2936	0.6554
SYPP	0.0960	- 0.1592	0.0858	0.4422	0.5040	0.2223	0.5848	0.0110	- 0.1082	0.2829	0.7247	0.2218

DF= Days to 50% Flowering, **DM**= Days to maturity, **PH**= Plant height (cm), **PB**= Number of primary branches/ plants, **SB**= Number of secondary branches/ plants, **LMR**= Length of main raceme (cm), **SMR**= Number of siliquae on main raceme, **S/SL**= Number of seed per siliquae, **SL**= Siliquae length (cm), **TSW**= Test weight (g), **BY**= Biological yield, **HI**= Harvest Index, **SSYP**= Seed yield per plant (g)

Table 8 Individual character's percentage contribution to genetic divergence

Source	Contribution %
Days to 50% flowering	2.36
Days to maturity	27.80
Plant height (cm)	7.64
Numbers of primary branches per plant	0.08
Numbers of secondary branches per plant	0.04
Length of main raceme (cm)	5.98
Numbers of siliquae on main raceme	37.63
Numbers of seeds per siliquae	0.09
Siliquae length(cm)	0.04
Test weight (g)	0.72
Biological Yield	12.43
Harvest Index	1.12
Seed Yield per Plant (g)	4.07
Total	100.00%

Table 9 Distribution of seventy-five Indian mustard genotypes by Tocher's method

Cluster No.	Name of genotypes	Numbers of genotypes
I	RB-50, Pusa Bold, Rohini, Kranti, RH-725, Maya, Vardan, Vasundhara, Swarn Jyoti, DMH1, JMWR-908-1, NRC-HB-101, NRC-HB-506, RVM-3, RH-749, NRC DR-2, DRMR IJ-31, CHINA, GSL-1, GSC-7, PC-5, PC-6, RP-9, JTC-1, JM-1, JM-2, RVM-1, RVM-2, PM-25, PM-26, PM-27, PM-30, Pusa Vijay, JMM-927, WRR-5, WRR-6, WRR-7, WRR-8, WRR-9, WRR-10, WRR-13, WRR-14, WRR-15, WRR-16, WRR-17, WRR-18, WRR-19, WRR-25, WRR-26, WRR-27, WRR-28, WRR-29, WRR-30, WRR-32	53
II	Pusa Jagannath, PusaJaiKisan, Albeli, Sej-2, Shraddha, RGN-73, JMM-991, RMM-12-03-18, WRR-11, WRR-12, WRR-20, WRR-21, WRR-22, WRR-31	14
III	Varuna	1
IV	L-4, JM-3, PM-28, RMM-10-01-01, RMM-12-01-18	5
V	KIRAN	1
VI	RP-9	1

Table10. Inter and intra-cluster distance in Indian mustard

Cluster	Cluster I	Cluster II	Cluster III	Cluster IV	Cluster V	Cluster VI
Cluster I	22.17	29.68	27.97	42.01	43.14	43.76
Cluster II		24.32	43.71	44.27	46.40	56.23
Cluster III			0.00	42.14	52.04	29.93
Cluster IV				30.85	51.73	38.74
Cluster V					0.00	51.63
Cluster VI						0.00

Table 11 Cluster mean in Indian mustard

Cluster	DF	DM	PH	PB	SB	LMR	SMR	S/SL	SL	TSW	BY	HI	SYP
Cluster I	51.7 2	153.0 4	179.7 0	8.10	10.3 9	88.83	139.2 2	12.9 7	5.1 0	4.76	31.4 3	30.2 3	9.03
Cluster II	52.7 5	118.7 9	171.5 1	8.07	10.9 3	84.79	117.2 9	12.1 8	4.7 3	4.82	30.4 6	34.4 7	9.66
Cluster III	43.0 0	178.5 0	162.9 9	8.00	10.5 0	89.00	222.5 0	11.0 0	4.7 5	5.10	38.5 0	25.9 4	9.98
Cluster IV	53.9 0	127.2 0	186.2 0	10.3 0	17.7 0	117.4 0	245.8 0	12.2 0	4.6 0	5.07	68.3 0	31.9 8	21.4 5
Cluster V	75.5 0	159.5 0	188.3 2	8.50	9.00	74.50	71.50	12.5 0	4.2 5	5.90	93.5 0	15.0 6	14.0 7
Cluster VI	74.0 0	178.5 0	202.3 4	13.5 0	21.5 0	72.50	288.0 0	15	4.7 5	4.41	78.5 0	24.8 3	19.4 7

DF= Days to 50% Flowering, DM= Days to maturity= Plant height (cm), PB= Number of primary branches/ plants, SB= Number of secondary branches/ plants, LMR= Length of main raceme (cm), SMR= Number of siliquae on main raceme, S/SL= Number of seed per siliquae, SL= Siliquae length (cm), TSW= Test weight (g), BY= Biological yield, HI= Harvest Index, SSYP= Seed yield per plant (g)




Fig. 2 Cluster Diagram in Indian mustard