

**Effect of Organic and Inorganic Fertilizer on Vegetative Growth Parameter of Lilium  
(*Lilium Longiflorum*) cv. Pavia Under Shade Net Condition**

**ABSTRACT**

The present investigation entitled, “**Effect of Organic and Inorganic Fertilizer on Vegetative Growth Parameter of Lilium (*Lilium Longiflorum*) cv. Pavia Under Shade Net Condition**” was under taken in the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, (*Allahabad*), during winter season of 2020-2021, 2021-2022 and Pooled. The experiment was layout in Randomized Block Design (RBD) with 13 treatments and each treatment replicated thrice. The treatments consist of different combinations of inorganic fertilizers (N, P and K) and organic fertilizers (FYM, Vermicompost and Poultry Manure) and control (No fertilizers and manures). The treatment T<sub>5</sub> 50% RDF + FYM 14 t ha<sup>-1</sup> was found the statistically significant compared to other treatment combination, which recorded highest plant height (23.68, 57.11 and 76.41 cm), spread area (8.19 cm<sup>2</sup>, 15.54 cm<sup>2</sup> and 18.47 cm<sup>2</sup>), Number of leaves (36.63, 57.11 and 76.41), Leaf area (15.03 cm<sup>2</sup>, 15.63 cm<sup>2</sup> and 15.33 cm<sup>2</sup>), Chlorophyll content (66.11 mg, 65.60 mg and 65.86 mg), followed by T<sub>9</sub> 25% RDF + Vermicompost 6.9 t ha<sup>-1</sup> and lowest yield was obtained from T<sub>0</sub> (control).

**Keywords:** Nitrogen, Phosphorus, Potash (Potassium), FYM, Vermicompost and Poultry Manure.

**INTRODUCTION**

*Lilium* sp. is an important genus for cut flowers, pot plants, and garden plants. It is widely cultivated across the world. “The genus *Lilium* belongs to the Liliaceae family, with over 100 species and over 9,400 cultivars divided into seven divisions”(Comber, 1949). The Northern Hemisphere, specifically Asia, North America, and Europe, particularly China, Nepal, Korea, and Japan, houses the majority of the world's gene centers for this species. It is a financially

significant genus for the production and sale of its cut flower in the global cut flower business **(Jimenez et. al., 2012)**.

Lilium produces large, appealing blooms in a variety of colours and forms, making them one of the best cut flowers. Lilium is considered as magnificent flowering potted plants which have a high decorative value in the landscape. Only three of the seven groups—Oriental hybrids, Asiatic hybrids, and cultivars of *Lilium longiflorum*—produced economically significant lily cultivars. At least 12 species from the *Sinomatagon* group underwent complex interspecific hybridization to produce Asiatic hybrids. Oriental hybrids are created when five species of the section *Archelirion*, to which *Lilium longiflorum* belongs, cross with one another. In addition to producing "LO" hybrids with "Asian" lilies, *L. longiflorum* also did so with "Oriental" lilies. The difficult cross-pollination of Asiatic and Oriental hybrids (also known as "OA" hybrids), two of the most significant lily clusters from a commercial standpoint, led to a breakthrough in lily breeding and hybridization. The availability of genetic variation in lilium for important characteristics has been greatly increased by intersectional crosses. The Netherlands produces lily bulbs on about 45% of its total farmland with Asian hybrids, 40% with Oriental hybrids, and 5% with cultivars of *L. longiflorum*. The *L. longiflorum* x Asiatic ('LA') hybrids, which are still relatively new, currently account for about 7% of the area used for lily production.

The quickest and clearest effect of nitrogen (N) on plant growth, which results in a high yield, is observed. "Due to the fact that it is a component of numerous amino acids, proteins, and chlorophyll, nitrogen is crucial for the growth of plants." Lack of nitrogen inhibits growth and lessens fruiting and flowering.

"Phosphorus is an essential component for plant growth and development. Phosphorus is necessary to maintain the quality of flowers". Phosphorus should be given as an initial dose, as  $P_2O_5$  is fundamentally required for optimal root development. Potassium is necessary for strong roots and stems and for plant respiration. Also known as potash. **(Parekh et.al., 2010)**.

"Lily plants may be cultivated in soil with low phosphorus levels without losing yield or quality. As a result, this study was structured out to determine the optimal nitrogen (N) and phosphorus (P) levels necessary for improved flower and bulb output and quality in Asiatic lily." **(Nair et al., 2000)**.

Vermicomposting is known as an environmentally beneficial process of decomposing organic wastes, and its result, Vermicopost, contains growth stimulating chemicals, which improve plant development and soil fertility, Microbial populations and water holding ability.

Poultry manure is an effective soil amendment that supplies nutrients for growing crops while also improving soil quality when applied correctly, as a result of its large organic matter concentration paired with accessible nutrients such as “nitrogen (N), phosphorous (P), and potassium (K) for plant growth”.

Farmyard manure is a degraded mixture of animal waste products including faeces and urine, as well as litter and scraps from hay roughages fed to livestock. “It typically contains 0.5% N, 0.2% P<sub>2</sub>O<sub>5</sub>, and 0.5% K<sub>2</sub>O”. Biological life of soil flora and fauna is stimulated by FYM, an excellent source of organic carbon. (Ghoshal and Singh, 1995).

#### **MATERIALS AND METHODS:-**

The present investigation entitled, “**Effect of Organic and Inorganic Fertilizer on Vegetative Growth Parameter of Lilium (*Lilium Longiflorum*) cv. Pavia Under Shade Net Condition**” was under taken in the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, (Allahabad). during winter season of 2020-2021, 2021-2022 and Pooled. The experiment was layout in Randomized Block Design (RBD) with 13 treatments and each treatment replicated thrice. The treatments consist of different combinations of inorganic fertilizers (N, P and K) and organic fertilizers (FYM, Vermicompost and Poultry Manure) and control (No fertilizers and manures). The treatments involved were T<sub>0</sub>. Control, T<sub>1</sub>. 100% RDF @ (NPK 140:100:80 kg ha<sup>-1</sup>), T<sub>2</sub>. 75% RDF + FYM 7 t ha<sup>-1</sup>, T<sub>3</sub>. 75% RDF + Vermicompost 2.3 t ha<sup>-1</sup>, T<sub>4</sub>. 75% RDF + Poultry manure 1.1 t ha<sup>-1</sup>, T<sub>5</sub>. 50% RDF + FYM 14 t ha<sup>-1</sup>, T<sub>6</sub>. 50% RDF + Vermicompost 4.6 t ha<sup>-1</sup>, T<sub>7</sub>. 50% RDF + Poultry manure 2.3 t ha<sup>-1</sup>, T<sub>8</sub>. 25% RDF + FYM 21 t ha<sup>-1</sup>, T<sub>9</sub>. 25% RDF + Vermicompost 6.9 t ha<sup>-1</sup>, T<sub>10</sub>. 25% RDF + Poultry manure 3.5 t ha<sup>-1</sup>, T<sub>11</sub>. 9.3 t FYM + 3.1 t Vermicompost + 1.5 t Poultry manure ha<sup>-1</sup> (33.33% FYM + 33.33% VC + 33.33% PM), T<sub>12</sub>. 25% RDF + 7 t FYM + 2.3 t Vermicompost + 1.1 Poultry manure ha<sup>-1</sup>.

## RESULTS AND DISCUSSION:-

### Growth parameters:-

The data revealed that the combination of different organic and inorganic fertilizers affected growth parameter during winter season of 2020-2021, 2021-2022 and Pooled. like Plant height, Plant spread, Number of leaves per plant, Leaf area and Chlorophyll content of Lilium as shown in (Table 1 and 2). Significant difference in the Plant height, Plant spread, Number of leaves per plant, Leaf area and Chlorophyll content was recorded due to application of different combinations of organic fertilizers and inorganic fertilizers. The treatment T<sub>5</sub> recorded the maximum plant height (73.42, 79.40 and 76.41 cm), followed by T<sub>9</sub> (72.33, 74.66 and 73.49 cm) and the maximum Plant spread T<sub>5</sub> (17.76 cm, 19.19 cm and 18.47 cm), followed by T<sub>9</sub> (17.10 cm, 18.99 cm and 18.05 cm) the maximum number of leaves per plant T<sub>5</sub> (73.76, 80.60 and 76.41), followed by T<sub>9</sub> (72.06, 77.06 and 73.49) and the maximum Leaf area was T<sub>5</sub> (15.03 cm<sup>2</sup>, 15.63 cm<sup>2</sup> and 15.33 cm<sup>2</sup>), followed by T<sub>9</sub> (14.90 cm<sup>2</sup>, 15.23 cm<sup>2</sup> and 15.06 cm<sup>2</sup>) the percentage of chlorophyll content T<sub>5</sub> (66.11mg, 65.60mg and 65.86mg), followed by T<sub>9</sub> (64.88mg, 65.23mg and 65.05mg) which differed significantly from each other as well from other treatments. Where in RDF: Recommended Dose of fertilizers, inorganic fertilizers : FYM, Vermicompost and Poultry manure. The plot size was 0.60 m x 0.40 m and spacing followed was 20 x 20 cm. the land was brought to a fine tilth by through ploughing and tillage. Irrigation channels and bunds were maintained properly and healthy and uniform bulb were collected from Van Zanten Flowerbulbs B.V 1e Loosterweg 1a, Hillegom, Netherlands (By The Plant Arena Nursery, Prayagraj-211007), & planted on 23<sup>rd</sup> November. Light irrigation was given after planting. The organic manures were applied 15-20 days before planting, for proper decomposition, 1/3 dose of nitrogen, and full dose phosphorus, potassium organic FYM, Vermicompost and Poultry manure, as per treatment were applied just before the planting. All cultural practices were followed regularly during crop growth and observations were recorded on growth characters i.e., plant height, plant spread, number of leaves per plant, leaf area and percentage of chlorophyll content were recorded from time to time. It was noticed that plant spread, number of leaves per plant, leaf area and percentage of chlorophyll content increased with increasing plant height successively with the increasing levels of inorganic fertilizer and organic fertilizer. Combination of inorganic fertilizer and organic fertilizer also recorded maximum plant height, plant spread, number of leaves per plant, leaf area and chlorophyll content also which helped the plants in better photosynthesis to attain vigor. The findings of the present investigation are in conformity with the reports of Foliar application thrice a

week increases fertilizer use efficiency of applied nitrogen, thus, increased nutrient contents might have accelerated the rates of various physiological and metabolic processes in the plant system that ultimately resulted in better plant growth. Similar findings were found by **Pahare *et al.*, (2020)** in Asiatic lily and **Singh *et al.*, (2013)** in carnation.

UNDER PEER REVIEW

**Table no. 1. Effect of organic and inorganic fertilizers on vegetative growth parameter of *Lilium cv. Pavia* under shade net condition.**

Treatments	2020-2021						2021-2022						Pooled					
	Plant height (cm)			Plant Spared (cm <sup>2</sup> )			Plant height (cm)			Plant Spared (cm <sup>2</sup> )			Plant height (cm)			Plant Spared (cm <sup>2</sup> )		
	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP
T <sub>0</sub> Control	16.29	43.16	67.07	5.13	10.60	14.10	18.93	44.66	61.13	5.80	11.20	14.05	17.61	43.91	64.10	5.46	10.90	14.07
T <sub>1</sub> 100% RDF @ (NPK 140:100:80 kg ha <sup>-1</sup> )	22.53	50.96	70.46	6.33	14.20	16.03	20.33	51.46	74.53	7.32	14.43	17.51	21.43	51.21	72.49	6.83	14.31	16.77
T <sub>2</sub> 75% RDF + FYM 7 t ha <sup>-1</sup>	22.55	51.76	71.10	6.50	14.46	16.80	21.40	51.13	73.33	7.93	15.07	18.01	21.97	51.45	72.21	7.21	14.77	17.40
T <sub>3</sub> 75% RDF + Vermicompost 2.3 t ha <sup>-1</sup>	19.70	52.90	68.96	7.26	14.90	17.03	21.30	52.13	72.33	8.49	14.78	18.15	20.50	52.51	70.65	7.88	14.84	17.59
T <sub>4</sub> 75% RDF + Poultry manure 1.1 t ha <sup>-1</sup>	19.80	51.06	70.58	6.53	14.46	17.16	20.86	50.80	71.60	8.03	15.45	18.63	20.33	50.93	71.09	7.28	14.95	17.90
<b>T<sub>5</sub> 50% RDF + FYM 14 t ha<sup>-1</sup></b>	<b>23.66</b>	<b>56.73</b>	<b>73.42</b>	<b>7.80</b>	<b>15.30</b>	<b>17.76</b>	<b>23.70</b>	<b>57.50</b>	<b>79.40</b>	<b>8.58</b>	<b>15.78</b>	<b>19.19</b>	<b>23.68</b>	<b>57.11</b>	<b>76.41</b>	<b>8.19</b>	<b>15.54</b>	<b>18.47</b>
T <sub>6</sub> 50% RDF + Vermicompost 4.6 t ha <sup>-1</sup>	19.76	54.16	68.44	6.56	14.16	16.66	22.60	50.73	73.80	7.80	14.90	18.36	21.18	52.45	71.12	7.18	14.53	17.51
T <sub>7</sub> 50% RDF + Poultry manure 2.3 t ha <sup>-1</sup>	20.90	52.13	71.37	6.50	14.43	15.56	22.73	52.76	73.40	8.39	15.37	17.58	21.81	52.45	72.38	7.44	14.90	16.57
T <sub>8</sub> 25% RDF + FYM 21 t ha <sup>-1</sup>	21.72	53.03	68.99	6.60	13.80	16.76	20.66	50.76	72.33	7.61	14.53	17.55	21.19	51.90	70.66	7.10	14.16	17.15
T <sub>9</sub> 25% RDF + Vermicompost 6.9 t ha <sup>-1</sup>	23.36	54.56	72.33	7.53	15.20	17.10	23.30	55.83	74.66	8.49	15.50	18.99	23.33	55.20	73.49	8.01	15.35	18.05
T <sub>10</sub> 25% RDF + Poultry manure 3.5 t ha <sup>-1</sup>	20.64	53.56	71.56	6.60	12.50	16.66	21.86	48.53	75.26	8.10	14.50	18.56	21.25	51.05	73.41	7.35	13.50	17.61
T <sub>11</sub> 9.3 t FYM + 3.1 t Vermicompost + 1.5 t Poultry manure ha <sup>-1</sup> (33.33% FYM +	20.16	52.16	71.94	7.20	13.83	15.93	22.73	51.60	70.73	8.03	15.09	18.48	21.45	51.88	71.33	7.61	14.46	17.20

33.33% VC + 33.33% PM)																		
T <sub>12</sub> 25% RDF + 7 t FYM + 2.3 t Vermicompost + 1.1 Poultry manure ha <sup>-1</sup>	20.16	51.83	71.35	7.03	14.20	16.06	21.83	52.66	72.60	8.27	14.62	18.85	21	52.25	71.97	7.65	14.41	17.45

**Table no. 2. Effect of organic and inorganic fertilizers on vegetative growth parameters of Lilium cv. Pavia under shade net condition.**

Treatments	2020-2021					2021-2022					Pooled				
	No. of leaves plant <sup>-1</sup>			Leaf area (cm <sup>2</sup> )	Chlorophyll (mg)	No. of leaves plant <sup>-1</sup>			Leaf area (cm <sup>2</sup> )	Chlorophyll (mg)	No. of leaves plant <sup>-1</sup>			Leaf area (cm <sup>2</sup> )	Chlorophyll (mg)
	25 DAP	50 DAP	75 DAP			25 DAP	50 DAP	75 DAP			25 DAP	50 DAP	75 DAP		
T <sub>0</sub> Control	19.93	48.30	60.16	12.66	55.60	21.33	49.16	64.73	12.76	56.84	20.63	43.91	64.10	12.71	56.22
T <sub>1</sub> 100% RDF @ (NPK 140:100:80 kg ha <sup>-1</sup> )	30.53	55.06	67.20	13.76	60.36	39.26	50.56	69.06	14.10	63.88	34.90	51.21	72.49	13.93	62.12
T <sub>2</sub> 75% RDF + FYM 7 t ha <sup>-1</sup>	30.86	53.00	67.20	13.63	60.51	40.13	53.63	73.80	13.96	64.15	35.50	51.45	72.21	13.79	62.33
T <sub>3</sub> 75% RDF + Vermicompost 2.3 t ha <sup>-1</sup>	28.93	51.86	65.46	14.16	63.05	38.06	49.83	75.00	14.83	63.40	33.50	52.51	70.65	14.49	63.23
T <sub>4</sub> 75% RDF + Poultry manure 1.1 t ha <sup>-1</sup>	28.06	53.20	68.60	13.83	62.05	37.36	52.46	74.46	14.23	63.81	32.71	50.93	71.09	14.03	62.93
<b>T<sub>5</sub> 50% RDF + FYM 14 t ha<sup>-1</sup></b>	<b>32.20</b>	<b>56.53</b>	<b>73.76</b>	<b>15.03</b>	<b>66.11</b>	<b>41.06</b>	<b>58.50</b>	<b>80.60</b>	<b>15.63</b>	<b>65.60</b>	<b>36.63</b>	<b>57.11</b>	<b>76.41</b>	<b>15.33</b>	<b>65.86</b>
T <sub>6</sub> 50% RDF + Vermicompost 4.6 t ha <sup>-1</sup>	29.73	54.93	69.20	13.63	62.16	34.63	56.56	78.60	14.30	64.62	32.18	52.45	71.12	13.96	63.39
T <sub>7</sub> 50% RDF + Poultry manure 2.3 t ha <sup>-1</sup>	26.66	54.00	69.23	13.36	60.86	38.80	56.13	76.53	13.83	64.92	32.73	52.45	72.38	13.59	62.89
T <sub>8</sub> 25% RDF + FYM 21 t ha <sup>-1</sup>	28.40	54.83	64.33	14.10	62.17	40.73	54.53	73.33	14.55	64.32	34.56	51.90	70.66	14.32	63.24
T <sub>9</sub> 25% RDF + Vermicompost 6.9 t ha <sup>-1</sup>	32.13	56.30	72.06	14.90	64.88	40.80	57.03	77.06	15.23	65.23	36.46	55.20	73.49	15.06	65.05

T <sub>10</sub> 25% RDF + Poultry manure 3.5 t ha <sup>-1</sup>	31.80	52.46	68.40	13.70	59.97	34.53	56.33	77.40	14.96	64.44	33.16	51.05	73.41	14.33	62.21
T <sub>11</sub> 9.3 t FYM + 3.1 t Vermicompost + 1.5 t Poultry manure ha <sup>-1</sup> (33.33% FYM + 33.33% VC + 33.33% PM)	27.33	52.63	65.76	14.16	61.46	38.00	53.76	73.33	15.00	63.81	32.66	51.88	71.33	14.58	62.63
T <sub>12</sub> 25% RDF + 7 t FYM + 2.3 t Vermicompost + 1.1 Poultry manure ha <sup>-1</sup>	30.13	53.30	65.40	14.23	61.81	40.73	56.83	78.00	15.23	63.74	35.43	52.25	71.97	14.73	62.77

UNDER PEER REVIEW

## **CONCLUSION:**

On the basis of present study, it is concluded that the application of T<sub>5</sub> 50% RDF + FYM 14 t ha<sup>-1</sup> resulted in maximum Plant height, Plant spread, Number of leaves per plant, Leaf area and chlorophyll content was found in maximum and the minimum was observed in T<sub>0</sub> (control).

UNDER PEER REVIEW

## REFERENCES:-

- Adhikari, Y.S., Bohra, M., Punetha, P., Upadhyay, S. and Nautiyal, B.P., (2018).**, Effect of Integrated Nutrient Management on Vegetative Growth, Floral Attributes, Corm Yield and Economics of Gladiolus cv. Arka Amar in Bharsar, Uttarakhand, Int. J. Pure App. Biosci. 6(4): 643-650.
- Abe H, Nakano M, Nakatsuka A, Nakayama M, Koshioka M, Yamagishi M., (2000),** Genetic analysis of floral anthocyanin pigmentation trait in Asiatic hybrid likely using molecular linkage maps, Theor Appl Genet, Vol.105; pp.1175-1182.
- Adnan, k., shashidhar, H. E. And Hittalani, S., (2013).**, Correlation and path analysis for root morphological traits in indica × indicia population of rice. Crop Research Hisar. 27(1):94-98
- Airadevi P. And Angadi.,(2014).**, Effect of integrated nutrient management on yield, economics and nutrient uptake of garland chrysanthemum (*Chrysanthemum coronarium* L.). Asian J. Hort., 9(1) pp.132-135.
- Anderson N, Plattes A, Opitz E, Younis A., (2011).**, Transgressive segregant, interspecific hybrids between *Lilium × formolongi* and, *L. Martagon* with unique Morphology , Acta Hort, Vol.900;pp.181-188.
- Anderson No, Younis A , Optiz e., (2009).**, Development of colored, non-vernalization-requiring seed propagated lilies , Acta Hort, Vol.836; pp.193-198,
- Anderson No , Younis A, Sun Y., (2010).**, Intersimple sequence repeats distinguish genetic differences in Easter Lily ‘Nellie White’ clonal ramets within and among bulb growers over years , J Amer Soc Hort Sci, Vol.135; pp.445-445.
- Anuburani A. And Gayathiri M.,(2008).**, Influence of integrated nutrient management on major nutrients in mullai (*Jasminum auriculatum*). Asian J. Hort., vol.3 No.2:323-326.
- Arya, G.R.J.K. and Gbalot, P.k., (2010).**, Effect of different levels of nitrogen, phosphorus and potassium on growth and flowering of African marigold cv. Pusa Narangi. Prog. Agri.10 (1) 8-83.
- Aryan A, Brader G, MCortel J, Paster M, Riedle-Bauer M., (2010).**, An abundant Stolbur tuft B phytoplasma strain is associated with grapevine, stinging nettle and *Hylesthe sobsoletus*. Eur J Plant Pathol 140:213-227.

**Asano Y., (1980),** Studies on crosses between distantly related species of lilies characteristics of newly obtained hybrids through embryo culture, J Japan Soc Hort. Sci., Vol.49; pp.241-250.

**Anuburani A., Arumugam S. And Gayathiri M., (2008),** Effect of organic in manures in combination with fertilizers on yield in Gundumalli (jasminum sambac Ait.) Asian J. Hort., Vol.3 No.2: 419-421.

**Balaji, S., Kulkarni, Reddy, B. S., Patil, B.C. and Divakara, A., (2006),** Influence of vermicompost and in situ vermiculture on the quality attributes and saleable yield in China asta. Scientific Hort., 10: 217-221.

**Bahr, L. And Compton,M.E., (2004),** Competence for in vitro bulb lets regeneration among eight liliium genotype, Hort. Science. 39(1): 127-129.

**Bankar, G.J. and Mukhopadhyay, A., (1990),** Effect of NPK on growth and flowering in tuberose cv “Double”. Indian J. Hort.,47 (1); 120-126.

**Bellubbi, S. Kulkarni, B.S and Patil, C.P., (2015),** Effect of integrated nutrient management on growth and flowering of gerbera (*Gerbera jamasonii* L.) var. Rosalin under naturally ventilated poly house condition. Int. J. Agril. Sc. & Vet. Med. 7(1): 93-97.

**Bhat M.R, Limaye S.r., (2010),** Nutrient status and plant growth promoting potential of prepared Vermicompost international Journal of environmental sciences Volume 3, No 1.

**Bouyocos (1952),** a hydrometer method for mating mechanical analysis of soil bull. Am Cerom.Soc 14.259.

**Comber H.F., (1949),** a new classification of the genus Liliium, RHS Lily Year Book, pp.86-105.

**Chaitra, R. And Patil, V.S., (2007),** Integrated nutrient management studies in China aster (*Callistephus chinensis* Ness) Var. Kamini. Darnataka J. Agril. Sci., 20(3): 689-690.

**Deivasigamani S.,Thanunathan K., Kathiresan R.M., sudhakar M., and Bharathi B., Karthikeyan., (2007),** Effect of bio-farming techniques on medicinal glory lily (*Gloriosa superb* L.) and Soil fertility status. Vol, 9(32), pp, 863-866.

**De jong PC (1974),** some notes on the evolution of lilies. Yearbook North North American Lily Society 27,23-28.

**Eidyan B., (2010).**, Effect of Iron and Citric Acid Foliar Applications in Combination with Nitrogen Fertigation on tuberose (*Polianthes tuberosa* L.), Horticulture. Daraj: Islamic Azad University-karaj Branch, 75.

**Gauhane, P.B., Kore V.N., Dixit, A.J.and Gondhali, B.V., (2004).**, Effect of graded doses of fertilizers and polythene mulches on growth, flower quality and yield of marigold (*Tagetes erects* L.) cv Pusa narangi gaudia. Orissa J. Horti.,32 (1):35-37.

**Grassotti A and Gimelli F. (2011).**, Bulb and cut flower production in the genus *Lilium* current status and the future, Acta Hort, Vol.900; pp.21-36.

**Ghule, A.D., Patil, P.v. and Kantharaju, K.T. (2003).**, Effect of different levels of nitrogen and phosphorus on growth and flowering of spider lily. J. Maharashtra Agri. Univ.28(2)128-130.

**Gupta P.,(2010).**, studies of the effect of integrated nutrient management on growth, flower production and seed vigour in African marigold (*Tagetes erecta* L.) cv. Giant African double orange, Thesis.

**Gupta L.M., Kumar S., Gupta M. And Sharma V., (2013).**, Integrated nutrient management for growth and yield in Glory Lily (*Gloriosa superb* L.) Journal of Medicinal Plants Research, Vol. 7(43), pp.3197-32201.

**Gupta M.L, Prasad A, Ram M, Kumar S., (2012).**, Effect of the vesicular-arbuscular mycorrhizal (VAM) fungus *Glomus fasciculatum* on the essential oil yield related charactes and nutrient acquisition in the crops of different cultivars under field conditions. Bioresource Technol 81:77-79.

**Hedden, p., Phillips, A.L., coles, J.P., Thomas, S., Appleford, N., Ward, D., Beale, M., and Lenton, J. (1999).**, Gibberellin biosynthesis: Genes, regulation and genetic manipulation. RIKEN Rev.21,29-30.

**Jackson, M.L., (1951).**, Soil organic carbon determination with fisher induction carbon apparatus. Soil Sci. Soc. Am. Proc. 16:370-371.

**Jefferson B., Howland H. (1995).**, The Gardener's Guide to Growing Lilies. Timber Press, Portland, Oregon, 160 pp.

**Jawaharlal, M. And Padmadevi, K., (2003).**, Effect of biofertilizers on growth and flowering of anthurium (*Anthurium andreanum* Linn.) cv. Temptation. Natl. Symp. Rec. Adv. Indian flort. Vellanikkar, India, Derala Agric, Univ., pp:32-34.

**Jambhelkar, H. 1994.** Bio-organic farming. Paper Presented in the seminar on development of Agriculture, Andhra Pradesh. 16.

**Jimenez, S., Plaza, B.M., Segura, M.L., Contreras, J.I. and Lao, T.M. (2012).**, Peat Substrate reuse in Liliium “Haveilia” crop. *Communications in Soil Science & Plant Analysis* **43**: 243-250.

**Kabir, A.k., Iman, m.R., Mondal, M.H. and Chowdhury, S., (2011).**, Response of Tuberose to INM, J. Environ. Sci. & natural Resource. **4**(2):55-59.

**Kejkar P.K. and polara N.D., (2015).**, Effect of n, p and k on growth, bulb yield and nutrient content in ratoon spider lily (*Hymenocallis littoralis* L.) cv. Local Hortflora Research Spectrum, **4**(1):22-27.

**Karimi M, Bleys A., Vanderhaeghen R, and Hilson P., (2007).**, Building blocks for plant gene assembly. *Plant Physiol* **145** 1183-1191.

**Kishore, G.R. and Singh, P.V., (2006).**, Effect of N.P.K. fertilization on vegetative growth of tuberose (*Polianthes tuberosa* L.) cv. Single. *Plant archives* Vol. No 6., pp.377-378.

**Khakha Ankita., Fatmi Urfi., N.N. Parihar., (2017).**, Effect of organic and inorganic fertilizers on plant growth and bulb yield of Asiatic Liliium hybrid courier under shadenet conditions. *Bioved* 2017 Vol.28 No.2 pp. 507-510.

**Kumar Madhur., Kasera Saurabh., Mishra Sanjay., Singh Nikhil V., and Singh Devi., (2018).**, Effect of Organic Manure and Inorganic Fertilizer on Growth and Yield Traits of Gladiolus (*Gladiolus grandiflora* L.) Cv. Plumtart. *Int.J.Curr.Microbiol.App.Sci* (2018) Special Issue-7: 1430-1435.

**Kishore, G.R., Arya J.K. and Ghalot P.K., (2010).**, Effect of different levels of nitrogen, phosphorus and potassium on growth and flowering of African marigold cv. ‘Pusa Narangi Gainda’. *Progressive Agriculture*, **82**(6): 941-945.

**Khan M.R. G., Ai X.-Y., Zhang J.-Z. (2014).**, Genetic regulation of flowering time in annual and perennial plants. *wileyInterdiscip. Rev. RNA* **5**,347-359.10.1002/wrna.1215.

**Lange M.J.P., Knop N. and Lange T., (2005).**, Stamen-derived bioactive gibberellins is essential for male flower development of *Cucurbita maxima* L. *J. Exp. Bot.* **63**: 2681-2691.

**Lin, W. C. and Wilkins H.F., (2009).**, Influence of bulb harvest date and temperature on growth and flowering of *Lilium longiflorum*. *J. Amer. Soc. Hort. Sci.* **100**: 6-9.

**Longchar A, and Kreditsu R., (2013).**, Flower yield and vase life of Gerbera in response to planting time and organic manures on Alfisol. *Sci. J. Agric.* 2(3): 124-128.

**Lucidos, J.G, Ryukh.A. Younis, C. K., Kim, Y., hwang, Son B.G. and Lim. K.B., (2013) .,** Defferent day and night temperatures responses in lilium hansonli Inrelation to growth and flower development. *Hort Environ Biotechnol.* 54:405-411.

**Matthews Victiria, (2007).**, International Lily Registrar Published in 2007 by the Royal Horticultural Society, 80 Vincent Square, London SW1P 2PE.

**Mohd. T. A., J.D. Desai, S.B. Parmar and B.R. Parmar., (2010).**, Effect of organic and inorganic fertilizers on growth, yield and quality of garlic cv GG.-1. *The Asian Journal of Horticulture*; Vol. 6 No. 1; (June, 2011) : 52-55.

**Manauwar A, Dheeraj K., and Mohammad, G., (2013).**, Effect of different nutrient management practices on engineering properties of soil and its response on marigold production. *Environment & Ecology.* 31 (1A): 289-292.

**McRae, E.A. (1998).**, Lilies: a guide for growers and collectors. 392pp., Timber press, Poetland, Oregon.

**Munikrishnappa, P.M., Katimani, K.N. and Ravikumar, M., (2004).**, Effect of vermicompost on growth and yield of tuberose (*Polianthus tuberosa* L.), Under semiarid tropics of north Karnataka. *National symposium on Recent Trends and Future Strategies in Ornamental Horticulture, Univesity of agricultural Sciences, Dharwal*, pp: 61.

**Mashaldi, A., (2000).**, Effect of organic and inorganic fertilizers on growth, yield and post harvest life of marigold (*Tagetes erecta* L.) cv. Double orange. *M.Sc.(Agri.) Thesis*, Univ. Agric. Sci., Bangalore.

**Mahanty, C. R., Misgra, N.K., and Padmaja P., (2013).**, Effect of nitrogen and phosphorous on flower production in marigold. *J. Orna. Hort.*, 5 (1): 67-68.

**Kumar, M., Kasera, S., Mishra, S., Singh, Nikhil V. and Singh, D., (2018).**, Effect of Organic Manure and Inorganic Fertilizer on Growth and Yield Traits of Gladiolus (*Gladiolus grandiflora* L.) Cv. Plumtart. *Int. J. Curr. Microbiol.App.Sci* (2018) Special Issue-7: 1430-1435

**Nair V. D., Jaleel C. A., Gopi R. Panneerselvam R. (2000),** Changes in growth and photosynthetic characteristics of *Ocimum sanctum* under growth regulator treatments. *Front. Biol, China* 4, 192-199, 10.1007/s1515-009-0001-1.

**Naseri, M. and Ebrahimi G.A., (2002),** Bulbous flowers physiology (translated) Jihad Mashhad University Press, pp 352.

**NHB, (2005),** National Horticulture Board, ministry of Agriculture, GOI. <http://www.nhb.gov.in>

**Olsen S. R. Cole C.V., Watanabe F.S., and Dean L. A. (1954),** Estimation of available phosphorus in soil by extraction with sodium bicarbonate. United States Department of Agriculture Circ.939.

**Parekh J, Jadeja D, Chanda S. (2010),** Efficacy of Aqueous and Methanol Extracts of Some Medicinal Plants for Potential Antibacterial Activity. *Turkish Journal of Biology*, 29: 203-210.

**Panse, V, G. and Sukhatme, P. V., (1984),** Statistical Methods for Agricultural Workers. ICAR, New Delhi.

**Parya, C.Pal, B. K. And Biswas, J., (2010),** Influence of integrated nutrient management on flower production efficiency, behaviour and quality of golden rod. *Joyrnal article Environment and Ecology*. 28(4): 2203-2205.

**Prakash, A., Sharma, S. K., Sindhu, S. S. and Prakash A., (2002),** Effect of phosphorus and FYM on NPK contents of marigold in chloride dominated saline soil. *Haryana J. Horti. Sci.*, 31 (1&2) : 47-49.

**Rao, K.D., Kameswari, P.L. and Rani, T.B., (2015),** Impact of integrated nutrient management on growth, flowering, yield and economics of tuberose. *Agric. Sci. Digrst*,35 (1): 66-69.

**Rout G.R., Mohapatra A. and Mohan J.S., (2006),** Tissue culture of ornamental pot plant: A critical review on present scenario and future prospect. *Biotechnological Advances* 24 (6) 531-560.

**Sathyanarayana, E., Patil, S., Bahubali, M. and Chawla, S. L., (2018),** Effect of INM on *Gladiolus (Gladiolus grandiflorus L.)* cv. American Beauty under Navsari and Tansa Conditions, *Int. J. Pure App. Biosci.* 6(4): 48-55 (2018).

**Singh, S.P., Kumar, N., Rizvi, S.E.H., and Sharma, P.K. (2014),** An economic analysis of gladiolus cultivation in Jammu district of J & K state. *Evonomic Affairs 2014*: 515-519.

**Sabah, S.S., (2014),** Effect of Different Organic and Inorganic Manure on Flower Yield and Tubers Yield of Dahlia (*Dahlia variabilis*) var. Glory of India as Intercropping with Damask Rose *European Academic Research – Vol.II, Issue 3 / June 2014.*

**Sahni, S., Sarma B. K., Singh D.P., Singh H. B. And Singh K.P., (2008),** Vermicompost enhances performance of plant growth-promoting rhizobacteria in *Cicerarietinum* rhizosphere against *Sclerotiumrolfsii* and quality of strawberry (*Fragaria x ananassa* Duch.) *Crop protection*, 27: 369-376.

**Subbaiah, B.V., Asija, G.L., (1956),** A rapid procedure for the estimation of available nitrogen in soils, *Current Sci.*, 25, 259. 32.

**Seyedi, M., Mohhammadi Torkashvand, A. and Allahyari. M.S., (2011),** Effect of medium, calcium concentration and nutrition on yield of lily M.Sc. Thesis, College of Agriculture, Islamic Azad University, Rasht Branch.

**Suseela, T., Chandrasekha, R., Bhaskar and Umakrishna, K. (2016),** Effect of organic manures, inorganic fertilizers and micronutrients on vegetative and floral characters of tuberose (*Polianthes tuberosa* L.) cv. Suvasini, *International Journal of Scientific and Research Publication*. 4 (2): 111-113.

**Van Tuyl, J.M., Van Dien, M.P., Van Creij, M.G.M., Van Kleinwee, T.C.M., Franken, J. and Bino, R.J.(1991),** Application of *in vitro* pollication, ovary culture, ovule culture and embryo rescue for overcoming incongruity barriers in interspecific liliium crosses. *Plant Science* **74**: 115-126.

**Wang J., Drayton M. C., George J., Cogan N. O., Baillie R. C., et al., (2010),** Identification of genetic factors influencing salt stress tolerance in white clover (*Trifolium repens* L.) by QTL analysis. *Theor. Appl. Genet.* 120: 607-619.

**Wu S.S., Chen L.N., Zhang Q.X., Lv Y. M., (1999),** Source and sink changes of lily bulb and the transportation role of the basal plate during the development of oriental hybrid lily 'Sorbonne'. *J. Food Agric, Environ.* 10 (2):213-219.