

**Effect of Organic and Inorganic Fertilizer on Vegetative Growth Parameter of Lilium
(*Lilium Longiflorum*) cv. Pavia Under Shade Net Condition**

ABSTRACT

The present investigation entitled, “**Effect of Organic and Inorganic Fertilizer on Vegetative Growth Parameter of Lilium (*Lilium Longiflorum*) cv. Pavia Under Shade Net Condition**” was under taken in the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, (*Allahabad*), during winter season of 2020-2021, 2021-2022 and Pooled. The experiment was layout in Randomized Block Design (RBD) with 13 treatments and each treatment replicated thrice. The treatments consist of different combinations of inorganic fertilizers (N, P and K) and organic fertilizers (FYM, Vermicompost and Poultry Manure) and control (No fertilizers and manures). The treatment T₅ 50% RDF + FYM 14 t ha⁻¹ was found the statistically significant compared to other treatment combination, which recorded highest plant height (23.68, 57.11 and 76.41 cm), spread area (8.19 cm², 15.54 cm² and 18.47 cm²), Number of leaves (36.63, 57.11 and 76.41), Leaf area (15.03 cm², 15.63 cm² and 15.33 cm²), Chlorophyll content (66.11 mg, 65.60 mg and 65.86 mg), followed by T₉ 25% RDF + Vermicompost 6.9 t ha⁻¹ and lowest yield was obtained from T₀ (control).

Keywords: Nitrogen, Phosphorus, Potash (Potassium), FYM, Vermicompost and Poultry Manure.

INTRODUCTION

Lilium sp. is an important genus for cut flowers, pot plants, and garden plants. It is widely cultivated across the world. “The genus *Lilium* belongs to the Liliaceae family, with over 100 species and over 9,400 cultivars divided into seven divisions”(Comber, 1949). The Northern Hemisphere, specifically Asia, North America, and Europe, particularly China, Nepal, Korea, and Japan, houses the majority of the world's gene centers for this species. It is a financially significant genus for the production and sale of its cut flower in the global cut flower business (Jimenez et. al., 2012).

Lilium produces large, appealing blooms in a variety of colours and forms, making them one of the best cut flowers. *Lilium* is considered as magnificent flowering potted plants which have a high decorative value in the landscape. Only three of the seven groups—Oriental hybrids, Asiatic hybrids, and cultivars of *Lilium longiflorum*—produced economically significant lily cultivars. At least 12 species from the *Sinomatagon* group underwent complex interspecific hybridization to produce Asiatic hybrids. Oriental hybrids are created when five species of the section *Archelirion*, to which *Lilium longiflorum* belongs, cross with one another. In addition to producing "LO" hybrids with "Asian" lilies, *L. longiflorum* also did so with "Oriental" lilies. The difficult cross-pollination of Asiatic and Oriental hybrids (also known as "OA" hybrids), two of the most significant lily clusters from a commercial standpoint, led to a breakthrough in lily breeding and hybridization. The availability of genetic variation in *lilium* for important characteristics has been greatly increased by intersectional crosses. The Netherlands produces lily bulbs on about 45% of its total farmland with Asian hybrids, 40% with Oriental hybrids, and 5% with cultivars of *L. longiflorum*. The *L. longiflorum* x Asiatic (LA') hybrids, which are still relatively new, currently account for about 7% of the area used for lily production.

The quickest and clearest effect of nitrogen (N) on plant growth, which results in a high yield, is observed. “Due to the fact that it is a component of numerous amino acids, proteins, and chlorophyll, nitrogen is crucial for the growth of plants.” Lack of nitrogen inhibits growth and lessens fruiting and flowering.

“Phosphorus is an essential component for plant growth and development. Phosphorus is necessary to maintain the quality of flowers”. Phosphorus should be given as an initial dose, as P_2O_5 is fundamentally required for optimal root development. Potassium is necessary for strong roots and stems and for plant respiration. Also known as potash. **(Parekh *et.al.*, 2010).**

“Lily plants may be cultivated in soil with low phosphorus levels without losing yield or quality. As a result, this study was structured out to determine the optimal nitrogen (N) and phosphorus (P) levels necessary for improved flower and bulb output and quality in Asiatic lily.” **(Nair *et al.*, 2000).**

Vermicomposting is known as an environmentally beneficial process of decomposing organic wastes, and its result, Vermicopost, contains growth stimulating chemicals, which improve plant development and soil fertility, Microbial populations and water holding ability.

Poultry manure is an effective soil amendment that supplies nutrients for growing crops while also improving soil quality when applied correctly, as a result of its large organic matter concentration paired with accessible nutrients such as “nitrogen (N), phosphorous (P), and potassium (K) for plant growth”.

Farmyard manure is a degraded mixture of animal waste products including faeces and urine, as well as litter and scraps from hay roughages fed to livestock. “It typically contains 0.5% N, 0.2% P_2O_5 , and 0.5% K_2O ”. Biological life of soil flora and fauna is stimulated by FYM, an excellent source of organic carbon. **(Ghoshal and Singh, 1995).**

MATERIALS AND METHODS:-

The present investigation was under taken in the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, (Allahabad). during winter season of 2020-2021, 2021-2022 and Pooled. The experiment was layout in Randomized Block Design (RBD) with 13 treatments and each treatment replicated thrice. The treatments consist of different combinations of inorganic fertilizers (N, P and K) and organic fertilizers (FYM, Vermicompost and Poultry Manure) and control (No fertilizers and manures). The treatments involved were T₀. Control, T₁. 100% RDF @ (NPK 140:100:80 kg ha⁻¹), T₂. 75% RDF + FYM 7 t ha⁻¹, T₃. 75% RDF + Vermicompost 2.3 t ha⁻¹, T₄. 75% RDF + Poultry manure 1.1 t ha⁻¹, T₅. 50% RDF + FYM 14 t ha⁻¹, T₆. 50% RDF + Vermicompost 4.6 t ha⁻¹, T₇. 50% RDF + Poultry manure 2.3 t ha⁻¹, T₈. 25% RDF + FYM 21 t ha⁻¹, T₉. 25% RDF + Vermicompost 6.9 t ha⁻¹, T₁₀. 25% RDF + Poultry manure 3.5 t ha⁻¹, T₁₁. 9.3 t FYM + 3.1 t Vermicompost + 1.5 t Poultry manure ha⁻¹ (33.33% FYM + 33.33% VC + 33.33% PM), T₁₂. 25% RDF + 7 t FYM + 2.3 t Vermicompost + 1.1 Poultry manure ha⁻¹.

RESULTS AND DISCUSSION:-

The present investigation concluded that treatment T₅. 50% RDF + FYM 14 t ha⁻¹ found the best results in terms of Growth parameters, is highly recommended for farmers for profitable cultivation.

Growth parameters:-

The data revealed that the combination of different organic and inorganic fertilizers affected growth parameter during winter season of 2020-2021, 2021-2022 and Pooled. like Plant height, Plant spread, Number of leaves per plant, Leaf area and Chlorophyll content of Lilium as shown in (Table 1 and 2). Significant difference in the Plant height, Plant spread, Number of leaves per plant, Leaf area and Chlorophyll content was recorded due to application of different combinations of organic fertilizers and inorganic fertilizers. The treatment T₅ recorded the maximum plant height (73.42, 79.40 and 76.41 cm), followed by T₉ (72.33, 74.66 and 73.49 cm) and the maximum Plant spread T₅ (17.76 cm, 19.19 cm and 18.47 cm), followed by T₉ (17.10

cm, 18.99 cm and 18.05 cm) the maximum number of leaves per plant T₅ (73.76, 80.60 and 76.41), followed by T₉ (72.06, 77.06 and 73.49) and the maximum Leaf area was T₅ (15.03 cm², 15.63 cm² and 15.33 cm²), followed by T₉ (14.90 cm², 15.23 cm² and 15.06 cm²) the percentage of chlorophyll content T₅ (66.11mg, 65.60mg and 65.86mg), followed by T₉ (64.88mg, 65.23mg and 65.05mg) which differed significantly from each other as well from other treatments. All cultural practices were followed regularly during crop growth and observations were recorded on growth characters i.e., plant height, plant spread, number of leaves per plant, leaf area and percentage of chlorophyll content were recorded from time to time. It was noticed that plant spread, number of leaves per plant, leaf area and percentage of chlorophyll content increased with increasing plant height successively with the increasing levels of inorganic fertilizer and organic fertilizer. Combination of inorganic fertilizer and organic fertilizer also recorded maximum plant height, plant spread, number of leaves per plant, leaf area and chlorophyll content also which helped the plants in better photosynthesis to attain vigor. Similar findings were found by **Pahare et al., (2020)** in Asiatic lily and **Singh et al., (2013)** in carnation.

Table no. 1. Effect of organic and inorganic fertilizers on vegetative growth parameter of *Lilium cv. Pavia* under shade net condition.

Treatments	2020-2021						2021-2022						Pooled					
	Plant height (cm)			Plant Spared (cm ²)			Plant height (cm)			Plant Spared (cm ²)			Plant height (cm)			Plant Spared (cm ²)		
	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP	25 DAP	50 DAP	75 DAP
T ₀ Control	16.29	43.16	67.07	5.13	10.60	14.10	18.93	44.66	61.13	5.80	11.20	14.05	17.61	43.91	64.10	5.46	10.90	14.07
T ₁ 100% RDF @ (NPK 140:100:80 kg ha ⁻¹)	22.53	50.96	70.46	6.33	14.20	16.03	20.33	51.46	74.53	7.32	14.43	17.51	21.43	51.21	72.49	6.83	14.31	16.77
T ₂ 75% RDF + FYM 7 t ha ⁻¹	22.55	51.76	71.10	6.50	14.46	16.80	21.40	51.13	73.33	7.93	15.07	18.01	21.97	51.45	72.21	7.21	14.77	17.40
T ₃ 75% RDF + Vermicompost 2.3 t ha ⁻¹	19.70	52.90	68.96	7.26	14.90	17.03	21.30	52.13	72.33	8.49	14.78	18.15	20.50	52.51	70.65	7.88	14.84	17.59
T ₄ 75% RDF + Poultry manure 1.1 t ha ⁻¹	19.80	51.06	70.58	6.53	14.46	17.16	20.86	50.80	71.60	8.03	15.45	18.63	20.33	50.93	71.09	7.28	14.95	17.90
T₅ 50% RDF + FYM 14 t ha⁻¹	23.66	56.73	73.42	7.80	15.30	17.76	23.70	57.50	79.40	8.58	15.78	19.19	23.68	57.11	76.41	8.19	15.54	18.47
T ₆ 50% RDF + Vermicompost 4.6 t ha ⁻¹	19.76	54.16	68.44	6.56	14.16	16.66	22.60	50.73	73.80	7.80	14.90	18.36	21.18	52.45	71.12	7.18	14.53	17.51
T ₇ 50% RDF + Poultry manure 2.3 t ha ⁻¹	20.90	52.13	71.37	6.50	14.43	15.56	22.73	52.76	73.40	8.39	15.37	17.58	21.81	52.45	72.38	7.44	14.90	16.57
T ₈ 25% RDF + FYM 21 t ha ⁻¹	21.72	53.03	68.99	6.60	13.80	16.76	20.66	50.76	72.33	7.61	14.53	17.55	21.19	51.90	70.66	7.10	14.16	17.15
T ₉ 25% RDF + Vermicompost 6.9 t ha ⁻¹	23.36	54.56	72.33	7.53	15.20	17.10	23.30	55.83	74.66	8.49	15.50	18.99	23.33	55.20	73.49	8.01	15.35	18.05
T ₁₀ 25% RDF + Poultry manure 3.5 t ha ⁻¹	20.64	53.56	71.56	6.60	12.50	16.66	21.86	48.53	75.26	8.10	14.50	18.56	21.25	51.05	73.41	7.35	13.50	17.61
T ₁₁ 9.3 t FYM + 3.1 t Vermicompost + 1.5 t Poultry manure ha ⁻¹ (33.33% FYM + 33.33% VC + 33.33% PM)	20.16	52.16	71.94	7.20	13.83	15.93	22.73	51.60	70.73	8.03	15.09	18.48	21.45	51.88	71.33	7.61	14.46	17.20
T ₁₂ 25% RDF + 7 t FYM + 2.3 t Vermicompost + 1.1 Poultry manure ha ⁻¹	20.16	51.83	71.35	7.03	14.20	16.06	21.83	52.66	72.60	8.27	14.62	18.85	21	52.25	71.97	7.65	14.41	17.45

Table no. 2. Effect of organic and inorganic fertilizers on vegetative growth parameters of Lilium cv. Pavia under shade net condition.

Treatments	2020-2021					2021-2022					Pooled				
	No. of leaves plant ⁻¹			Leaf area (cm ²)	Chlorophyll (mg)	No. of leaves plant ⁻¹			Leaf area (cm ²)	Chlorophyll (mg)	No. of leaves plant ⁻¹			Leaf area (cm ²)	Chlorophyll (mg)
	25 DAP	50 DAP	75 DAP			25 DAP	50 DAP	75 DAP			25 DAP	50 DAP	75 DAP		
T ₀ Control	19.93	48.30	60.16	12.66	55.60	21.33	49.16	64.73	12.76	56.84	20.63	43.91	64.10	12.71	56.22
T ₁ 100% RDF @ (NPK 140:100:80 kg ha ⁻¹)	30.53	55.06	67.20	13.76	60.36	39.26	50.56	69.06	14.10	63.88	34.90	51.21	72.49	13.93	62.12
T ₂ 75% RDF + FYM 7 t ha ⁻¹	30.86	53.00	67.20	13.63	60.51	40.13	53.63	73.80	13.96	64.15	35.50	51.45	72.21	13.79	62.33
T ₃ 75% RDF + Vermicompost 2.3 t ha ⁻¹	28.93	51.86	65.46	14.16	63.05	38.06	49.83	75.00	14.83	63.40	33.50	52.51	70.65	14.49	63.23
T ₄ 75% RDF + Poultry manure 1.1 t ha ⁻¹	28.06	53.20	68.60	13.83	62.05	37.36	52.46	74.46	14.23	63.81	32.71	50.93	71.09	14.03	62.93
T₅ 50% RDF + FYM 14 t ha⁻¹	32.20	56.53	73.76	15.03	66.11	41.06	58.50	80.60	15.63	65.60	36.63	57.11	76.41	15.33	65.86
T ₆ 50% RDF + Vermicompost 4.6 t ha ⁻¹	29.73	54.93	69.20	13.63	62.16	34.63	56.56	78.60	14.30	64.62	32.18	52.45	71.12	13.96	63.39
T ₇ 50% RDF + Poultry manure 2.3 t ha ⁻¹	26.66	54.00	69.23	13.36	60.86	38.80	56.13	76.53	13.83	64.92	32.73	52.45	72.38	13.59	62.89
T ₈ 25% RDF + FYM 21 t ha ⁻¹	28.40	54.83	64.33	14.10	62.17	40.73	54.53	73.33	14.55	64.32	34.56	51.90	70.66	14.32	63.24
T ₉ 25% RDF + Vermicompost 6.9 t ha ⁻¹	32.13	56.30	72.06	14.90	64.88	40.80	57.03	77.06	15.23	65.23	36.46	55.20	73.49	15.06	65.05
T ₁₀ 25% RDF + Poultry manure 3.5 t ha ⁻¹	31.80	52.46	68.40	13.70	59.97	34.53	56.33	77.40	14.96	64.44	33.16	51.05	73.41	14.33	62.21
T ₁₁ 9.3 t FYM + 3.1 t Vermicompost + 1.5 t Poultry manure ha ⁻¹ (33.33% FYM + 33.33% VC + 33.33% PM)	27.33	52.63	65.76	14.16	61.46	38.00	53.76	73.33	15.00	63.81	32.66	51.88	71.33	14.58	62.63
T ₁₂ 25% RDF + 7 t FYM + 2.3 t Vermicompost + 1.1 Poultry manure ha ⁻¹	30.13	53.30	65.40	14.23	61.81	40.73	56.83	78.00	15.23	63.74	35.43	52.25	71.97	14.73	62.77

CONCLUSION:

On the basis of present study, it is concluded that the application of T₅ 50% RDF + FYM 14 t ha⁻¹ resulted in maximum Plant height, Plant spread, Number of leaves per plant, Leaf area and chlorophyll content was found in maximum and the minimum was observed in T₀ (**control**).

Authors' contribution: The contribution of each author is to be mentioned in the following heads.

Conceptualization of research (The research was conceptualized by supervisor- Dr. V.M. Prasad, Dr. Anita Kerketta and Dr. Kanchan S. Das)

Contribution of experimental materials (Department of Horticulture, SHUATS resources were utilized for experimentation)

Execution of field/lab experiments and data collection (Narendra Kumar executed the field experiments and comprehensive data collection was done which was further curated by supervisors)

Analysis of data and interpretation (Analysis was performed by Narendra Kumar and interpretations were further curated by Dr. V.M. Prasad)

Preparation of the manuscript (The manuscript was prepared by collaborative efforts of Narendra Kumar and Dr. V.M. Prasad).

Declaration: None of the authors have any conflict of interest. This manuscript is being shared with the consent of each authors associated.

REFERENCES:-

1. **Anuburani A., Arumugam S. And Gayathiri M., (2008)**, Effect of organic in manures in combination with fertilizers on yield in Gundumalli (*Jasminum sambac* Ait.) Asian J. Hort., Vol.3 No.2: 419-421.
2. **Balaji, S., Kulkarni, Reddy, B. S., Patil, B.C. and Divakara, A., (2006)**., Influence of vermicompost and in situ vermiculture on the quality attributes and saleable yield in China aster. Scientific Hort., 10: 217-221.
3. **Kumar, M., Kasera, S., Mishra, S., Singh, Nikhil V. and Singh, D., (2018)**., Effect of Organic Manure and Inorganic Fertilizer on Growth and Yield Traits of *Gladiolus (Gladiolus grandiflora* L.) Cv. Plumtart. *Int. J. Curr. Microbiol.App.Sci* (2018) Special Issue-7: 1430-1435
4. **Kishore, G.R., Arya J.K. and Ghalot P.K., (2010)**., Effect of different levels of nitrogen, phosphorus and potassium on growth and flowering of African marigold cv. 'Pusa Narangi Gainda'. Progressive Agriculture, 82(6): 941-945.
5. **Mohd. T. A., J.D. Desai, S.B. Parmar and B.R. Parmar., (2010)**., Effect of organic and inorganic fertilizers on growth, yield and quality of garlic cv GG.-1. The Asian Journal of Horticulture; Vol. 6 No. 1; (June, 2011) : 52-55.
6. **Munikrishnappa, P.M., Katimani, K.N. and Ravikumar, M., (2004)**., Effect of vermicompost on growth and yield of tuberose (*Polianthus tuberosa* L.), Under semiarid tropics of north Karnataka. *National symposium on Recent Trends and Future Strategies in Ornamental Horticulture, Univesity of agricultural Sciences, Dharwal*, pp: 61.
7. **Mashaldi, A., (2000)**., Effect of organic and inorganic fertilizers on growth, yield and post harvest life of marigold (*Tagetes erecta* L.) cv. Double orange. *M.Sc.(Agri.) Thesis*, Univ. Agric. Sci., Bangalore.
8. **Nair V. D., Jaleel C. A., Gopi R. Panneerselvam R. (2000)**., Changes in growth and photosynthetic characteristics of *Ocimum sanctum* under growth regulator treatments. *Front. Biol, China* 4, 192-199, 10.1007/s1515-009-0001-1.
9. **Parekh J, Jadeja D, Chanda S. (2010)**., Efficacy of Aqueous and Methanol Extracts of Some Medicinal Plants for Potential Antibacterial Activity. *Turkish Journal of Biology*, 29: 203-210.

10. **Prakash, A., Sharma, S. K., Sindhu, S. S. and Prakash A., (2002),** Effect of phosphorus and FYM on NPK contents of marigold in chloride dominated saline soil. *Haryana J. Horti. Sci.*, 31 (1&2) : 47-49.
11. **Sabah, S.S., (2014),** Effect of Different Organic and Inorganic Manure on Flower Yield and Tubers Yield of Dahlia (*Dahlia variabilis*) var. Glory of India as Intercropping with Damask Rose *European Academic Research – Vol.II, Issue 3 / June 2014.*
12. **Suseela, T., Chandrasekha, R., Bhaskar and Umakrishna, K. (2016),** Effect of organic manures, inorganic fertilizers and micronutrients on vegetative and floral characters of tuberose (*Polianthes tuberosa* L.) cv. Suvasini, *International Journal of Scientific and Research Publication.* 4 (2): 111-113.

UNDER PEER REVIEW