

Biochemical and oxidative stress responses in *Clarias gariepinus* exposed to sublethal concentrations of benzo[a]pyrene.

ABSTRACT

The effect of benzo[a]pyrene (BaP) on selected plasma biochemical parameters of the tropical African catfish *C. gariepinus* was investigated. Apparently healthy juvenile fish (n = 90; mass = 19.7 ± 1.8 g) were exposed to sublethal concentrations of BaP over a period of 35 days after which haematological and plasma biochemical analysis were carried out on whole blood and plasma respectively. While there were significant declines in red blood cell count (RBC), haemoglobin, haematocrit and platelet count, significant elevations were observed in mean cell volume (MCV), mean cell haemoglobin concentration (MCHC) and white blood cell (WBC) count. Significant increases were also observed in the activities of the liver enzymes, alanine transaminase (ALT), aspartate transaminase (AST) and alkaline phosphatase (ALP). There were significant increases in the activities of the oxidative stress enzymes, catalase (CAT), superoxide dismutase (SOD) and glutathione s-transferase (GST). Findings from this study reveal that benzo[a]pyrene causes changes in haematological and plasma biochemical profiles in exposed aquatic organisms.

Key words: benzo[a]pyrene, haematological, biochemical, plasma, aquatic, pollution

1. INTRODUCTION

Aquatic pollution is a cause for concern because of its impact on aquatic organisms which serve as food for humans and also maintain the ecological balance. One of the causes of aquatic pollution are polycyclic aromatic hydrocarbons (PAHs). PAHs are a collection of carbon-based organic compounds which consist of two or more benzene rings [1]. PAHs are found widespread in the environment. These compounds are formed from either natural or anthropogenic activities [2]. The dominant sources of PAHs in the environment are from human activity. Such activities include but are not limited to wood-burning, combustion of fossil fuels, mining activities etc. One of the most studied PAHs in the terrestrial but least studied in the aquatic environment is benzo[a]pyrene (BaP). BaP is a five-ringed PAH that is formed by the incomplete combustion of organic compounds just like other PAHs. In recent times, BaP has been confirmed as having properties capable of causing cancer [3]

The aim of this study was to evaluate the effect of BaP on haematological parameters, some enzymes involved in amino acid metabolism and possible oxidative response of the tropical catfish to BaP. Our intention was to provide data that will help in the prediction of the effects of sublethal concentrations of BaP on aquatic organisms. The tropical catfish was chosen for this work because of its abundance in tropical climes together with its hardiness and relative ease of handling. This study will contribute immensely to the toxicological database and also enhance the ecological risk assessment of BaP in aquatic environments.

2. MATERIALS AND METHODS

2.1 Chemicals

All reagents used were of analytical grade. Benzo[a]pyrene (BaP) was obtained from Sigma Aldrich (Germany). Acetone was obtained from BDH chemicals (UK). Kits for alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP) were obtained from Randox laboratories ltd. (Antrim, UK). Adrenaline, 1-chloro-2,4-

dinitrobenzene (CDNB), 5,5-dithiobis-2-nitrobenzoic acid (DTNB), and hydrogen peroxide (H₂O₂) were all purchased from Sigma Chemical Company (London, UK).

2.2 Animals

Juvenile catfish, *C. gariepinus* (n = 90) weighing 19.7±1.8 g were obtained from a commercial fish farm in Aba, South east Nigeria. The fish were acclimatized for 2 weeks in dechlorinated tap water prior to experimentation. The fish were fed twice daily *ad libitum* with commercial fish feed. Fish fecal matter and uneaten food was removed daily to prevent organic pollution with the attendant algal growth.

2.3 Sublethal toxicity tests

Stock solutions of BaP were prepared by dissolving BaP in distilled water taking acetone as solvent carrier. Test solutions were prepared by dilution of stock solutions in tap water. The acute LC₅₀ value of BaP was determined in a pilot study using a semi-static method. During sublethal studies, fish were exposed to 1/2 and 1/4 of the LC₅₀ value (corresponding to treatment levels 1 and 2). A solvent control was included in the experimental design. Fish were kept in groups of 10 in 30L plastic tanks containing the test solutions. Experiments were performed in triplicates. Period of exposure lasted 35 days

2.4 Assays

At the end of the exposure period, fish were anaesthetized using non-chemical method by hypothermia. Blood was then collected from the immobilized fish by caudal vein puncture method as described by Argungu *et al.* (2015) [4] using a 5ml sterile disposable syringe with a 22 gauge needle. The blood was transferred to EDTA tubes. The blood samples were spun with a bucket centrifuge at 4000 rpm for 5 minutes so as to separate the plasma from the packed cells. The plasma obtained were kept in plain blood containers and used for biochemical analysis.

2.4.1 Haematology

Haematological parameters were determined using automated haematology analyzer machine (Mindray BC 2300, USA).

2.4.2 Liver enzymes

Randox diagnostic kits were used to assay the activities of plasma ALT, AST and ALP. Assay of AST and ALT activities was based on the principle as modified by Tietz *et al.* (1994) [5]. AST activity was assayed by monitoring the concentration level of oxaloacetate hydrazone formed with 2,4-dinitrophenylhydrazine at 546 nm, while ALT activity was assayed by monitoring levels of pyruvate hydrazone formed with 2,4-dinitrophenylhydrazine at 546 nm. ALP was assayed in accordance with the principles of Tietz (1995) [6]. The p-nitrophenol formed by the hydrolysis of p-nitrophenyl phosphate confers a yellowish colour on the reaction mixture. Its intensity is monitored at 405 nm to measure enzyme activity.

2.4.3 Oxidative stress enzymes

The procedure of Misra and Fridovich (1972) [7] as described by Magwere *et al.* (1996) [8] was used to determine plasma superoxide dismutase (SOD) activity by measuring the inhibition of autooxidation of adrenaline at pH 10.2 and 30°C. SOD activity was expressed in U/mL. Plasma catalase activity was determined according to the method of Sinha (1972) [9] by measuring the reduction of dichromate in acetic acid to chromic acetate at 570 nm. Catalase activity was expressed in kU/L. Plasma glutathione S-transferase (GST) activity was

determined by the method described by Habig *et al.* (1974) [10] using 1-chloro-2,4-dinitrobenzene (CDNB) as substrate. GST activity was expressed in U/L.

2.5 Statistical analysis

Results were expressed as mean \pm standard error. Data from the different treatment groups were compared by a one-way analysis of variance (ANOVA) followed by a Scheffes test to determine statistically different groups. All differences were considered significant at $p < 0.05$. Statistical analysis was performed using Microsoft Excel and the SPSS statistical package (ver. 24.0 SPSS Company, Chicago, IL, USA).

3. RESULTS

The results showed that exposure to sublethal concentrations of BaP affected some of the haematological parameters. There was a statistically significant ($p < 0.05$) reduction in the red blood cell count of exposed fish. The observed reduction appeared to be dose-dependent with a higher dose of the chemical (8 $\mu\text{g/L}$) causing a greater reduction in the reduction compared to the lower dose (4 $\mu\text{g/L}$). There was also an observed reduction in haemoglobin concentration and haematocrit. The impact of BaP on the erythrocyte indices showed mixed results. While there were increases in MCV and MCHC, there were no observed changes in MCH. There was a dose-dependent increase in WBC count. Though there was a decline in platelet count, the reduction was not statistically significant.

Fish exposed to 8 $\mu\text{g/L}$ BaP showed significant ($p < 0.05$) increases in the activities of the liver enzymes with AST showing the highest activity. BaP induced the expression of the oxidative stress enzymes. All three enzymes CAT, SOD and GST showed statistically significant ($p < 0.05$) increases in their activities compared to the control.

Table 1: Haematological parameters of *C. gariepinus* exposed to sublethal concentrations of benzo[a]pyrene

Parameter	Control group	Benzo[a]pyrene concentration	
		4 $\mu\text{g/L}$	8 $\mu\text{g/L}$
RBC ($\times 10^9/\text{L}$)	2.86 \pm 0.09 ^a	2.28 \pm 0.05 ^b	2.23 \pm 0.13 ^b
Hb (g/L)	100.25 \pm 3.9 ^a	88.75 \pm 0.63 ^b	91.5 \pm 0.67 ^b
Hct (%)	37.52 \pm 2.38 ^a	30.25 \pm 0.32 ^b	29.68 \pm 0.32 ^b
MCV (fL)	132.8 \pm 1.71 ^a	132.55 \pm 1.36 ^a	120.25 \pm 1.07 ^b
MCH (pg)	38.75 \pm 0.19	36.9 \pm 1.15	38.98 \pm 0.49
MCHC (g/L)	272 \pm 3.4 ^a	294.25 \pm 0.85 ^b	313.25 \pm 3.5 ^b
WBC ($\times 10^9/\text{L}$)	90.87 \pm 4.2 ^a	125.83 \pm 3.32 ^b	132.2 \pm 2.5 ^b
PLT ($\times 10^9/\text{L}$)	20 \pm 1.2	18 \pm 2.2	16 \pm 1.6




Figure 1: Plasma ALT, AST and ALP activities of *C. gariepinus* exposed to sublethal concentrations of benzo[a]pyrene. Means not sharing the same letter (a, b or c) are statistically different at $p < 0.05$




Figure 2: Plasma CAT activity of *C. gariepinus* exposed to sublethal concentrations of benzo[a]pyrene. Means not sharing the same letter (a or b) are statistically different at $p < 0.05$




Figure 3: Plasma SOD activity of *C. gariepinus* exposed to sublethal concentrations of benzo[a]pyrene. Means not sharing the same letter (a, b or c) are statistically different at $p < 0.05$




Figure 4: Plasma GST activity of *C. gariepinus* exposed to sublethal concentrations of benzo[a]pyrene. Means not sharing the same letter (a or b) are statistically different at $p < 0.05$

4. DISCUSSION

BaP had adverse effects on some of the studied haematological parameters. At the end of our experiment (35 days). Fish exposed to BaP showed significant reductions in the RBC count compared to the control. The potential of BaP to cause anaemia is likely due to changes in the metabolism of iron. BaP may also inhibit the synthesis of red blood cells in the exposed fish. Many inorganic and organic contaminants have been shown to minimize the concentrations of red blood cells in circulation [11]. Findings by Kim *et al.* (2007) [12] show that subchronic dietary B[a]P exposure caused a significant reduction in erythrocytes. Similar

results were also observed in common carp exposed to ethinylestradiol [13] and rainbow trout exposed to heavy metals [14]. Both studies demonstrated dose-dependent reductions in the red blood cell counts in fish exposed to the contaminants.

Haemoglobin (Hb) is a polypeptide containing heme and globin as prosthetic group and apoprotein respectively [15]. BaP exposure caused a significant reduction in the observed Hb concentration. As in red blood cell count, the reduction in Hb concentration can be attributed to the inhibition of erythropoiesis in the exposed fish. A similar finding was made by Kim *et al.* (2007) [16] where BaP exposure led to a significant decline in Hb concentration. Recently, Dey *et al.* (2021) [17] showed that naphthalene caused a significant reduction in Hb concentration. The authors attributed the observed reduction in Hb concentration to disruptions in oxygen supply to the tissues. This impairment could have caused a decline in metabolism and subsequently led to a reduction in energy generation.

Hematocrit (HCT) also known as packed cell volume (PCV) assays the volume of packed red blood cells compared to the whole blood [18]. Fish exposed to BaP showed significant reductions in the HCT value. The low HCT values in the present study could be attributed to the combined effects of destruction of erythrocytes, inhibition of erythropoiesis and destruction of erythrocytes [19]. Gluszczak *et al.* (2006) [20] reported the reduced HCT value in *L. obtusidens* to glyphosate exposure. Similarly, Barcellos *et al.* (2003) [21] showed that lower HCT could indicate the anaemic status of the fish in contact with contaminants.

While there were significant increases in MCV and MCHC, there were no observed changes in MCH on exposure to BaP. Red blood cell parameters are very useful in understanding the nature of anemias. Anemias can be classified according to the size of the erythrocyte, as being normocytic (normal MCV), macrocytic (increased MCV), or microcytic (decreased MCV) [22].

The present work showed that BaP was immunotoxic to *C. gariiepinus*. There was an observed increase in the levels of circulating white blood cells in exposed fish. Leukocytes which are white blood cells are a crucial part of the immune system in most vertebrates and take part in immune responses. They circulate in the blood and mount inflammatory response to toxicants or pathogens [23]. Ramesh and Saravanan (2008) [24] reported an elevation in WBCs in Indian carp *Labeo rohito* exposed to deltamethrin at sublethal concentrations. More recently, Parma and Shar (2021) [25] demonstrated that reactive red (RR) azo dye 120 caused an increase in WBC count in Indian catfish exposed to the insecticide, malathion.

Platelets perform important roles in wound healing and inflammation. It is proposed that they perform these functions via interaction with cells of the immune system [26]. Platelets are however not cells, but rather fragments of cytoplasmic origin derived from megakaryocytes²⁶. In the present study, though there was a decline in platelet count, though the observed change was not significant. The results of Dey *et al.* (2021) [17] also disclosed a reduction in platelet count in *A. testudineus* under naphthalene exposure.

ALT, AST and ALP are enzymes involved in amino acid metabolism. Changes in the activities of these enzymes allow the identification of tissue damage in organs like the liver and kidney [27]. In the present study, there were observed increases in the activity of the enzymes in fish exposed to BaP. These increases could be indicative of liver damage and possible disruptions in the permeability of the hepatic membrane. This would have subsequently led to the release of these enzymes to the blood [28]. Derakhshesh *et al.* (2019) [3] examined the effect of BaP on liver cell culture of *E. coioides*. The authors reported increases in activities of all three enzymes in varying concentrations of BaP. Increases in all three enzymes have also been reported in *S. schlegelii* [29] and in *C. carpio* on exposure to pyrene [30].

Oxidative stress is known to be caused as a result of an imbalance between the production of reactive oxygen species (ROS) and the capacity of the biological system to

mop up the oxidizing agents [31]. In the present study, BaP was found to induce the expression of the antioxidant enzyme systems. The scientific literature is rich in publications which show that toxicants lead to the production of free radicals and that these free radicals induced the expression of CAT, SOD and GST to ameliorate the effects of oxidative stress on the exposed biological system. Jiffa *et al.* (2006) [32] observed significant increases in SOD and CAT activities in a dose-effect related pattern in *L. japonix* exposed to benzo[a]pyrene. Palanikumar *et al.* (2011) [33] also reported increases in GST activity of BaP-exposed fish. These results are in agreement with the findings of the present work.

5. CONCLUSION

In conclusion, BaP causes changes in haematological parameters in *C. gariepinus* and has a damaging effect on the liver tissue via oxidative stress. These effects could predispose aquatic organisms to disease and parasite infestation. There is need for further research to study the impact of BaP in aquatic organisms at the genetic level and also to study metabolism and biotransformation pathways of BaP in *C. gariepinus*. The information obtained from this study will help to assess the sublethal effects of PAHs in general and BaP in particular and to establish water quality criteria for control policies and conservation strategies in tropical region

ETHICAL APPROVAL

Animal ethic committee approval was obtained prior to the commencement of experiment

References

1. Byeong-Kyu L, Sources, Distribution and Toxicity of Polyaromatic Hydrocarbons (PAHs). 2016; 17: 4. Accessed 29 December, 2019.
Available:
<http://www.intechopen.com/books/air-pollution/sources-distribution-and-toxicity-of-polyaromatic-hydrocarbonspahs>
2. Sørensen A, Wichert B. Asphalt and Bitumen. In: *Ullmann's Encyclopedia of Industrial Chemistry*. Weinheim: Wiley-VCH; 2018.
3. Derakhshesh N, Negin S, AbdolAli M, Mahmoud H. Exposure of liver cell culture from the orange-spotted grouper, *Epinephelus coioides*, to benzo[a]pyrene and light results in oxidative damage as measured by antioxidant enzymes. *Chemosphere*. 2019; 22(6) 54 – 58.
4. Argungu LA, Siraj S, Christanus A. A simple and rapid method for blood collection from walking catfish. *Iran. J. Fish. Sci.* 2015; 16(3) 934 – 944.
5. Tietz NW, Pruden EL, Siggaard-Andersen O. Liver function. In: Burtis AC, Ashwood ER, editors. *Tietz Textbook of Clinical Chemistry*. London: Saunders; 1994.
6. Tietz NW. *Clinical Guide to Laboratory Tests*. Philadelphia: Saunders; 1995.
7. Misra HP, Fridovich I. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J. Biol. Chem.* 1972; 247, 3170 – 3175.

8. Magwere T, Naik YS, Hasler JA. Effects of chloroquine treatment on antioxidant enzymes in rat liver and kidney. *Free Radic. Biol. Med.* 1996; 41: 321–327.
9. Sinha AK. Colorimetric assay of catalase. *Anal. Biochem.* 1997; 47: 389–394.
10. Habig WH, Pabst MJ, Jakoby WB. Glutathione transferases, the first enzymatic step in mercapturic acid formation. *J. Biol. Chem.* 1974; 249: 7130–7139.
11. Dey S, Palas S, Niladri S. Dose specific responses of *Anabas testudineus* (Bloch) to anthracene (PAH): Haematological and biochemical manifestation. *Emerg. Contam.* 2019; 5: 232 – 239.
13. Schwaiger J, Spieser OH, Bauer C, Ferling H, Mallow U. Chronic toxicity of nonylphenol and ethinylestradiol: Haematological and histopathological effects in juvenile common carp (*Cyprinus carpio*). *Aquat. Toxicol.* 2000; 51: 69 – 78.
14. Haux C, Larsson A. Long-term sublethal physiological effects in rainbow trout, *Salmo gairdneri*, during exposure to cadmium and after subsequent recovery. *Aquat. Toxicol.* 1984; 5: 129–142.
15. Chhabra, N. Structure of Hemoglobin-An Overview. 2013; 13:7. Accessed 30 December 2020.
Available:
<http://www.ourbiochemistry.com/knowledge-base/category/proteins>
16. Kim SG, Park DK, Jang, SW, Lee JS, Kim SS, Chung MH. Effects of dietary Bezo[a]pyrene on growth and hematological parameters in juvenile rockfish, *Sebastes schlegeli* (Hilgendorf). *Bull. Environ. Contam. Toxicol.* 2007; 81: 470 – 474.
17. Dey S, Puspita B, Arghya M, Palas S. Blood Biochemical and Erythrocytic Morphopathological Consequences of Naphthalene Intoxication in Indian Teleost, *Anabas testudineus* (Bloch), *Environ. Toxicol. Pharm.* 2020; 80: 103 – 109.
18. Mondal H. Hematocrit. 2020; 16:4. Accessed 27 October 2021.
Available:
<http://www.statpearls.com/articlelibrary/viewarticle/36935>.
19. Eriegha OJ, Omitoyin BO, Jani EK. Evaluation of Haematological and Biochemical Parameters of Juvenile *Oreochromis niloticus* after Exposure to Water Soluble Fractions of Crude Oil. *J. Appl. Sci. Environ. Manag.* 2017; 21(6): 1041 – 1045.
20. Gluszcak L, Denis D, Crestani M. Effects of glyphosate herbicide on acetylcholinesterase activity and metabolic and haematological parameters in Piava,. *Ecotoxicol. Environ. Saf.* 2006; 65(2): 214 – 231.
21. Barcellos L, Carlos K, Rodriguez L, Hematological and biochemical characteristics of male jumdia: changes after acute stress. *Aqua. Res.* 34. 2003; 21: 45 – 52.
22. Walker HK, Hall WD, Hurst JW. *Clinical Methods: The History, Physical, and Laboratory Examinations.* Boston: Butterworths; 1990.

23. Tigner A, Sherif A, Ian M. Histology, White Blood Cell. 2021; 4: 3. Accessed 11 August 2021.
Available:
<https://www.ncbi.nlm.nih.gov/books/NBK563148/>
24. Ramesh M, Saravanan M. Haematological and biochemical responses in a fresh water fish *Cyprinus carpio* exposed to chlorpyrifos. *Int. J. Integr. Biol.* 2008; 3(1): 80 – 84.
25. Parmar A, Shah I. Acute toxicity, behavioural response and haematological alterations of *Catla catla* exposed to Reactive Red 120 textile dye. *Ind. J. Exp. Biol.* 2021; 6(59): 275 – 279.
26. Valli VEO. Hematopoietic system. In: Maxie, M. G. editor. *Kennedy and Palmer's Pathology of Domestic Animals*. Elsevier, St. Louis: Elsevier; 2007.
27. Begum G. In vivo biochemical changes in liver and gill of *clarias batrachus* during cypermethrin exposure and following cessation of exposure. *Pest. Biochem. Physio.* 2005; 82(3): 185 – 196.
28. Yousef MI, Awad TI, Mohamed EH. Deltamethrin-induced oxidative damage and biochemical alterations in rat and its attenuation by vitamin E. *Toxicology.* 2006; 227: 240 – 247.
29. Kim SG, Park DK, Jang SW, Lee JS, Kim SS, Chung MH. Effects of dietary benzo[a]pyrene on growth and hematological parameters in juvenile Rockfish, *Seabastes schlegeli* (Hilgendorf). *Bull. Environ. Contam. Toxicol.* 2008; 81: 470 – 474.
30. Shirdel I, Mohammad RK, Milad S, Roya O. The response of thyroid hormones, biochemical and enzymological biomarkers to pyrene exposure in common carp (*Cyprinus carpio*). *Ecotox. Environ. Saf.* 2016; 130: 207 – 213.
31. Sawyer DB, Siwik DA, Xiao L, Pimental D. Role of oxidative stress in myocardial hypertrophy and failure. *J. Mol. Cell. Cardiol.* 2002; 34(4): 379 – 388.
32. Jifaa W, Yu Zhiminga Y, Xiuxiana S, Youa W. Response of integrated biomarkers of fish (*Lateolabrax japonicus*) exposed to benzo[a]pyrene and sodium dodecylbenzene sulfonate. *Ecotox. Environ. Saf.* 2006; 65: 230 – 237.
33. Palanikumar N, Kumaraguru CM, Ramakritinan. Biochemical response of anthracene and benzo[a]pyrene in milkfish *Chanos chanos* L. *Ecotox. Environ. Saf.* 2012; 75: 187–197.