

Common Fixed Point Theorem for (ϕ, \mathfrak{F}) -Multi-Valued Mappings in Cone b -Metric Spaces Over Banach Algebra

Author's contributions

This work was carried out in collaboration between both authors. Author RJ designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author RJ and KM managed the analysis of the study. Author KM managed the literature searches. Both authors read and approved the final manuscript.

Original Research Article

Abstract

In the present paper, we introduced the concept of generalized multi-valued contraction mappings, via the class functions Φ and Ψ . Also we proved some fixed point results for (ϕ, \mathfrak{F}) -multi-valued mappings on cone b -metric spaces over Banach algebra \mathfrak{A} . The conditions for existence and uniqueness of the fixed point are investigated. We give an example to support our main result.

Keywords: Cone b -Metric space, Multi-valued mappings, Banach algebra.

MSC: 47H10, 54H25, 54M20.

1 Introduction

In 1922, Stefan Banach [3] proved a fixed point theorem for contractive mappings in complete metric spaces. In 1969, Nadler [12] Introduce the concept of Multi-valued function. In 1989, Bakhtin [4] introduced the concept of b -metric space opens a gate to further generalization of metric space. Later, Czerwik [6,7] initiate the concept of b -metrics which generalized usual metric spaces. After his contribution, many results were presented in β -generalized weak contractive multifunction's and b -metric spaces. In 2007, Huang et al. [15] introduced cone metric space with normal cone, as a generalization of metric space. In 2012, Aydi et al [2]. Reformulate the b -metric space. Many researcher work in this area of research of multivalued function and b -metric spaces [1,5,8,9,10,11,12,20]. Liu and Xu [17] introduced the notion of cone metric space over Banach algebras, replacing Banach spaces by Banach algebras as the underlying spaces of cone metric space. They improved the fixed point theorems for (ϕ, \mathfrak{F}) -multi-valued mappings on cone b -metric spaces over Banach algebra, via the class functions Φ and Ψ

Let \mathfrak{A} be a real Banach algebra, i.e. \mathfrak{A} is a real Banach space in which an operation of multiplication is defined, subject to the following properties:

For all $\varrho, \varsigma, \nu \in \mathfrak{A}, \delta \in \mathfrak{R}$

- (i) $\varrho(\varsigma\varepsilon) = (\varrho\varsigma)\varepsilon;$
- (ii) $\varrho(\varsigma + \varepsilon) = \varrho\varsigma + \varrho\varepsilon$ and $(\varrho + \varsigma)\varepsilon = \varrho\varepsilon + \varsigma\varepsilon;$
- (iii) $\delta(\varrho\varsigma) = (\delta\varrho)\varsigma = \varrho(\delta\varsigma);$
- (iv) $\|\varrho\varsigma\| \leq \|\varrho\| \|\varsigma\|.$

We shall assume that the Banach algebra \mathfrak{A} has a unit, i.e., a multiplicative identity e such that $e\varrho = \varrho e = \varrho$ for all $\varrho \in \mathfrak{A}$. An element $\varrho \in \mathfrak{A}$ is said to be invertible if there is an inverse element $\varsigma \in \mathfrak{A}$ such that $\varrho\varsigma = \varsigma\varrho = e$. The inverse of ϱ is denoted by ϱ^{-1} .

Let \mathfrak{A} be a real Banach algebra with a unit e and $\varrho \in \mathfrak{A}$. If the spectral radius $\rho(\varrho)$ of ϱ is less than 1, that is

$$\rho(\varrho) = \lim_{n \rightarrow \infty} \|\varrho^n\|^{\frac{1}{n}} = \inf_{n \geq 1} \|\varrho^n\|^{\frac{1}{n}} < 1$$

then $e - \varrho$ is invertible. Actually,

$$(e - \varrho)^{-1} = \sum_{i=0}^{\infty} \varrho^i.$$

A subset \mathfrak{P} of \mathfrak{A} is called a cone of \mathfrak{A} if

- i. $\{\theta, e\} \subset \mathfrak{P},$
- ii. $\mathfrak{P}^2 = \mathfrak{P}\mathfrak{P} \subset \mathfrak{P}, \mathfrak{P} \cap (-\mathfrak{P}) = \{\theta\},$
- iii. $\mathfrak{h}\mathfrak{P} + \beta\mathfrak{P} \subset \mathfrak{P}, \forall \mathfrak{h}, \beta \in \mathfrak{R}.$

For a given cone $\mathfrak{P} \subset \mathfrak{A}$, we define a partial ordering \preceq with respect to \mathfrak{P} by $\varrho \preceq \varsigma$ if and only if $\varsigma - \varrho \in \mathfrak{P}$; $\varrho \prec \varsigma$ will stand for $\varrho \preceq \varsigma$ and $\varrho \neq \varsigma$, while $\varrho \ll \varsigma$ stand for $\varsigma - \varrho \in \text{int}\mathfrak{P}$, where $\text{int}\mathfrak{P}$ denotes the interior of \mathfrak{P} . If $\text{int}\mathfrak{P} \neq \emptyset$, then \mathfrak{P} is called a solid cone. Write $\|\cdot\|$ as the norm of \mathfrak{A} . A cone \mathfrak{P} is called normal if there is a number $\mathfrak{M} > 0$ such that $\forall \varrho, \varsigma \in \mathfrak{A}$, we have

$$\theta \preceq \varrho \preceq \varsigma \implies \|\varrho\| \leq \mathfrak{M} \|\varsigma\|.$$

The least positive number satisfying above is called the normal constant of \mathfrak{P} . Note that, for any normal cone \mathfrak{P} we have $\mathfrak{M} \geq 1$.

In the following we suppose that \mathfrak{A} is a real Banach algebra with a unit e , \mathfrak{P} is a solid cone and \preceq with respect to \mathfrak{P} .

2 Preliminaries

Lemma 2.1

(see [17]) If \mathfrak{C} is a real Banach space with a cone \mathfrak{P} and if $\delta \preceq \delta\delta$ with $\delta \in \mathfrak{P}$ and $0 \leq \delta < 1$, then $\delta = \theta$.

Lemma 2.2

(see [17]) If \mathfrak{C} is a real Banach space with a solid cone \mathfrak{P} and if $\theta \preceq u \ll c$ for each $\theta \ll c$, then $u = \theta$.

Lemma 2.3

(see [17]) Let \mathfrak{B} be a cone in a Banach algebra \mathfrak{A} and $\mathcal{K} \in \mathfrak{B}$ be a given vector. Let $\{u_n\}$ be a sequence in \mathfrak{B} . If $\forall c_1 \gg \theta, \exists \mathfrak{N}_1 \ni u_n \ll c_1 \forall n > \mathfrak{N}_1$, then $\forall c_2 \gg \theta, \exists \mathfrak{N}_2 \ni \mathcal{K}u_n \ll c_2 \forall n > \mathfrak{N}_2$.

Lemma 2.4.

(see [17]) If \mathfrak{E} is a real Banach space with a solid cone \mathfrak{B} and $\{\varrho_n\} \subset \mathfrak{B}$ is a sequence with $\|\varrho_n\| \rightarrow 0 (n \rightarrow \infty)$, then $\forall \theta \ll c, \exists \mathfrak{N} \in \mathbb{N} \ni n > \mathfrak{N}$ we have, $\varrho_n \ll c$ i. e., $\{\varrho_n\}$ is a c -sequence.

Lemma 2.5

(see [19]) Let \mathfrak{A} be a Banach algebra with a unit $e, i, j \in \mathfrak{A}$. If i commutes with j , then $\rho(i + j) \leq \rho(i) + \rho(j), \rho(ij) \leq \rho(i)\rho(j)$.

Remark 2.6

(see [19]) If $\rho(\varrho) < 1$, then $\|\varrho_n\| \rightarrow 0$ as $n \rightarrow \infty$.

Definition 2.7

(see [16]) Let X be a non-empty set, $\omega \geq 1$ be a constant and \mathfrak{A} be a Banach algebra. A function $D_b: X \times X \rightarrow \mathfrak{A}$ is said to be a cone b -metric provide that, for all $\varrho, \varsigma, \varepsilon \in X$,

(d1) $D_b(\varrho, \varsigma) = 0$ if and only if $\varrho = \varsigma$;

(d2) $D_b(\varrho, \varsigma) = D_b(\varsigma, \varrho)$;

(d3) $D_b(\varrho, \varepsilon) \leq \omega[D_b(\varrho, \varsigma) + D_b(\varsigma, \varepsilon)]$.

A pair (X, D_b) is called a cone b -metric space over Banach algebra \mathfrak{A} .

Example 2.8.

Let $\mathfrak{A} = C[\mathfrak{d}, \mathfrak{b}]$ be the set of continuous functions on the interval $[\mathfrak{d}, \mathfrak{b}]$ with the supremum norm. Define multiplication in the usual way. Then \mathfrak{A} is a Banach algebra with a unit 1. Set $\mathfrak{B} = \{\varrho \in \mathfrak{A}: \varrho(t) \geq 0, t \in [\mathfrak{d}, \mathfrak{b}]\}$ and $X = \mathfrak{A}$. Defined a mapping $D_b: X \times X \rightarrow \mathfrak{A}$ by $D_b(\varrho, \varsigma)(t) = |\varrho - \varsigma|^p e^t \forall \varrho, \varsigma \in X$, where $p > 1$ is a constant. This makes (X, D_b) into a cone b -metric space over Banach algebra \mathfrak{A} with the coefficient $b = 2^{p-1}$, but it is not a cone metric space over Banach algebra since the triangle inequality is not satisfied.

Definition 2.9

(see [16]) Let (X, D_b) be a cone b -metric space over Banach algebra $\mathfrak{A}, \varrho \in X$, let $\{\varrho_n\}$ be a sequence in X . Then

- i. $\{\varrho_n\}$ converges to ϱ whenever for every $c \in \mathfrak{A}$ with $\theta \ll c$ there is natural number $n_0 \ni D_b(\varrho_n, \varrho) \ll c, \forall n \geq n_0$. We denote this by $\lim_{n \rightarrow \infty} \varrho_n = \varrho$.
- ii. $\{\varrho_n\}$ is a Cauchy sequence whenever for every $c \in \mathfrak{A}$ with $\theta \ll c$ there is natural number $n_0 \ni D_b(\varrho_n, \varrho_m) \ll c, \forall n, m \geq n_0$.
- iii. $\{\varrho, D_b\}$ is complete cone b -metric if every Cauchy sequence in X is convergent.

Remark 2.10

(see [18]) Let (X, D_b) be a cone b -metric space over Banach algebra \mathfrak{A} with the coefficient $\omega \geq 1$, denote

$$\mathfrak{N}(X) = \{\mathcal{A}: \mathcal{A} \text{ is non empty subset of } X\} \text{ and } s(p) = \{q \in \mathfrak{C}: p \preceq q\} \text{ for } q \in \mathfrak{A},$$

$$s(\mathfrak{d}, \mathfrak{B}) = \bigcup_{\mathfrak{b} \in \mathfrak{B}} s(D_b(\mathfrak{d}, \mathfrak{b})) = \bigcup_{\mathfrak{b} \in \mathfrak{B}} \{q \in \mathfrak{A}: D_b(\mathfrak{d}, \mathfrak{b}) \preceq q\} \text{ for } \mathfrak{d} \in X \text{ and } \mathfrak{B} \in \mathfrak{N}(X).$$

For $\mathcal{A}, \mathfrak{B} \in \mathfrak{N}(X)$ we denote

$$s(\mathcal{A}, \mathfrak{B}) = (\bigcap_{\mathfrak{d} \in \mathcal{A}} s(\mathfrak{d}, \mathfrak{B})) \cap (\bigcap_{\mathfrak{b} \in \mathfrak{B}} s(\mathfrak{b}, \mathcal{A})).$$

Remark 2.11

(see [18]) Let (X, D_b) be a cone b -metric space over Banach algebra \mathfrak{A} , with the coefficient $\omega \geq 1$. If $\mathfrak{A} = \mathfrak{R}$ and $\mathfrak{B} = \mathfrak{R}_0^+$ then (X, D_b) is a metric space. Moreover, for $\mathcal{A}, \mathfrak{B} \in \mathfrak{CB}(X)$, $\mathcal{H}(\mathcal{A}, \mathfrak{B}) = \inf s(\mathcal{A}, \mathfrak{B})$ is the Hausdorff distance induced by D_b .

Definition 2.12

(see [18]) Let (X, D_b) be a cone b -metric space over Banach algebra \mathfrak{A} . A map $\mathfrak{S}: X \rightarrow \mathfrak{CB}(X)$ is said to be multi-valued contraction if $\exists 0 \leq s < 1$ such that $\mathcal{H}(\mathfrak{S}q, \mathfrak{S}\zeta) \leq s D_b(q, \zeta)$, for all $q, \zeta \in X$.

Lemma 2.13

(see [18]) If $\mathcal{A}, \mathfrak{B} \in \mathfrak{CB}(X)$ and $\mathfrak{d} \in \mathcal{A}$, then for each $\epsilon > 0$, there exists $\mathfrak{b} \in \mathfrak{B}$ such that $D_b(\mathfrak{d}, \mathfrak{b}) \leq \mathcal{H}(\mathcal{A}, \mathfrak{B}) + \epsilon$.

Lemma 2.14

(see [16]) Let \mathfrak{C} be a real Banach space with a solid cone \mathfrak{P}

- 1) If $\mathfrak{d}_1, \mathfrak{d}_2, \mathfrak{d}_3 \in \mathfrak{C}$ and $\mathfrak{d}_1 \preceq \mathfrak{d}_2 \ll \mathfrak{d}_3$, then $\mathfrak{d}_1 \ll \mathfrak{d}_3$.
- 2) If $\mathfrak{d}_1 \in \mathfrak{P}$ and $\mathfrak{d}_1 \ll \mathfrak{d}_3$ for each $\mathfrak{d}_3 \gg \theta$, then $\mathfrak{d}_1 = \theta$.

Lemma 2.15

(see [16]) Let \mathfrak{P} be a solid cone in a Banach algebra \mathfrak{A} . Suppose that $\mathfrak{h} \in \mathfrak{P}$ and $\{\varrho_n\} \subset \mathfrak{P}$ is a c -sequence. Then $\{\mathfrak{h}\varrho_n\}$ is a c -sequence.

Lemma 2.16

(see [16]) Let \mathfrak{A} be a Banach algebra with a unit $e, \mathfrak{h} \in \mathfrak{A}$, then $\lim_{n \rightarrow \infty} \|\mathfrak{h}^n\|^{\frac{1}{n}}$ exists and the spectral radius $\rho(\mathfrak{h})$ satisfies

$$\rho(\mathfrak{h}) = \lim_{n \rightarrow \infty} \|\mathfrak{h}^n\|^{\frac{1}{n}} = \inf \|\mathfrak{h}^n\|^{\frac{1}{n}}.$$

If, then $(\delta e - \mathfrak{h})$ is invertible in \mathfrak{A} , moreover,

$$(\delta e - \mathfrak{h})^{-1} = \sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{\delta^{i+1}},$$

where δ is a complex constant.

Definition 2.18

(see [8]) Let $\mathfrak{S}, \mathfrak{Q}: X \rightarrow X$ be a mappings on set X .

- 1) If $\mathfrak{w} = \mathfrak{S}\varrho = \mathfrak{Q}\varrho$ for some $\varrho \in X$, then ϱ is called a coincidence point of \mathfrak{S} and \mathfrak{Q} , and \mathfrak{w} is called a point of coincidence of \mathfrak{S} and \mathfrak{Q} .
- 2) The pair $(\mathfrak{S}, \mathfrak{Q})$ is called weakly compatible if \mathfrak{S} and \mathfrak{Q} commute at all of their coincidence points, that is, $\mathfrak{S}\mathfrak{Q}\varrho = \mathfrak{Q}\mathfrak{S}\varrho, \forall \varrho \in \mathcal{C}(\mathfrak{S}, \mathfrak{Q}) = \{\varrho \in X : \mathfrak{S}\varrho = \mathfrak{Q}\varrho\}$.

Lemma 2.19

(see [8]) Let \mathfrak{S} and \mathfrak{Q} be weakly compatible self-maps of a set X . If \mathfrak{S} and \mathfrak{Q} have a unique point of coincidence $\mathfrak{w} = \mathfrak{S}\varrho = \mathfrak{Q}\varrho$, then \mathfrak{w} is the unique common fixed point of \mathfrak{S} and \mathfrak{Q} .

3 MAIN RESULTS

We prove a unique fixed point for generalized (ϕ, \mathfrak{S}) - multi-valued mappings via the class functions Φ and Ψ .

Lemma 3.1

Let \mathfrak{A} be a Banach algebra with a unit $e, \mathfrak{h} \in \mathfrak{A}$, if δ is a complex constant and $\rho(\mathfrak{h}) < |\delta|$, then $\rho((\delta e - \mathfrak{h})^{-1}) \leq \frac{1}{|\delta| - \rho(\mathfrak{h})}$.

Proof. Since $\rho(\mathfrak{h}) < |\delta|$, it follows by lemma 2.16 that $(\delta e - \mathfrak{h})$ is invertible and

$$(\delta e - \mathfrak{h})^{-1} = \sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{\delta^{i+1}}.$$

Set $\omega = \sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{\delta^{i+1}}, \omega_n = \sum_{i=0}^n \frac{\mathfrak{h}^i}{\delta^{i+1}}$, then $\omega_n \rightarrow \omega (n \rightarrow \infty)$ and ω_n commutes with $\omega \forall n$. It follows immediately from lemma 2.5 that

$$\begin{aligned} \rho(\omega_n) &= \rho(\omega_n - \omega + \omega) \leq \rho(\omega - \omega_n) + \rho(\omega) \Rightarrow \rho(\omega_n) - \rho(\omega) \leq \rho(\omega - \omega_n), \\ \rho(\omega) &= \rho(\omega - \omega_n + \omega) \leq \rho(\omega - \omega_n) + \rho(\omega_n) \Rightarrow \rho(\omega) - \rho(\omega_n) \leq \rho(\omega - \omega_n), \end{aligned}$$

Which imply that

$$|\rho(\omega_n) - \rho(\omega)| \leq \rho(\omega - \omega_n) \leq \|\omega - \omega_n\| \Rightarrow \rho(\omega_n) \rightarrow \rho(\omega) (n \rightarrow \infty).$$

Thus again by lemma 2.5,

$$\begin{aligned} \rho((\delta e - \mathfrak{h})^{-1}) &= \rho\left(\sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{\delta^{i+1}}\right) = \rho(\omega) = \lim_{n \rightarrow \infty} \rho(\omega_n) \\ &= \lim_{n \rightarrow \infty} \rho\left(\sum_{i=0}^n \frac{\mathfrak{h}^i}{\delta^{i+1}}\right) \\ &\leq \lim_{n \rightarrow \infty} \sum_{i=0}^n \frac{[\rho(\mathfrak{h})]^i}{\delta^{i+1}} \\ &= \lim_{n \rightarrow \infty} \sum_{i=0}^{\infty} \frac{[\rho(\mathfrak{h})]^i}{\delta^{i+1}} = \frac{1}{|\delta| - \rho(\mathfrak{h})}. \end{aligned}$$

Lemma 3.2

Let \mathfrak{A} be a Banach algebra with a unit e and \mathfrak{P} be a solid cone in \mathfrak{A} . Let $\gamma \in \mathfrak{A}$ and $\varrho_n = \gamma^n$. If $\rho(\gamma) < 1$, then $\{\varrho_n\}$ is a c -sequence.

Proof. Since $\rho(\gamma) = \lim_{n \rightarrow \infty} \|\gamma^n\|^{\frac{1}{n}} < 1$, then $\exists \tau > 0 \ni \lim_{n \rightarrow \infty} \|\gamma^n\|^{\frac{1}{n}} < \tau < 1$. Letting n be big enough, we obtain $\|\gamma^n\|^{\frac{1}{n}} \leq \tau$, which implies that $\|\gamma^n\|^{\frac{1}{n}} \leq \tau^n \rightarrow 0 (n \rightarrow \infty)$. So $\|\gamma^n\| \rightarrow 0$, i.e., $\|\varrho_n\| \rightarrow 0 (n \rightarrow \infty)$. Note that $\forall c \gg \theta$, there is $\beta > 0$ such that

$$U(c, \beta) = \{\varrho \in E : \|\varrho - c\| < \beta\} \subset \mathfrak{P}.$$

In view of $\|\varrho_n\| \rightarrow 0 (n \rightarrow \infty)$, $\exists \mathfrak{N} \ni \|\varrho_n\| < \beta \forall n > \mathfrak{N}$. Consequently, $\|(c - \varrho_n) - c\| = \|\varrho_n\| < \beta$, this leads to $c - \varrho_n \in U(c, \beta) \subset \mathfrak{P}$, that is, $c - \varrho_n \in \text{int}\mathfrak{P}$, thus $\varrho_n \ll c \forall n > \mathfrak{N}$.

Definition 3.3

Let \mathfrak{A} be a Banach algebra and $\mathfrak{P} = \mathfrak{R}_0^+$ be a cone in \mathfrak{A} . A mapping $\mathfrak{F}: \mathfrak{P} \rightarrow \mathfrak{P}$ such that

- 1) \mathfrak{F} is non-decreasing and continuous;
- 2) $\lim_{n \rightarrow \infty} \mathfrak{F}^n(t) = \theta$ for all $(t \in \mathfrak{P}), t \geq 0$, where \mathfrak{F}^n stands for the n^{th} iterate of \mathfrak{F} ;
- 3) $\mathfrak{F}(t) < t$ for each $t > 0$;
- 4) $\mathfrak{F}(\theta) = \theta$.

Definition 3.4

Let \mathfrak{A} be a Banach algebra and $\mathfrak{P} = \mathfrak{R}_0^+$ be a cone in \mathfrak{A} . A mapping $\phi: \mathfrak{P} \rightarrow \mathfrak{P}$ such that:

- 1) ϕ is monotone non-decreasing and continuous;
- 2) $\{\phi^n(t)\} (t > 0)$ is a c -sequence in \mathfrak{P} ;
- 3) If $\{u_n\}$ is a c -sequence in \mathfrak{P} , then $\{\phi(u_n)\}$ is also a c -sequence in \mathfrak{P} ;
- 4) $\phi(t) = \mathcal{K}t$, for some $(\mathcal{K} \in \mathfrak{P}), \mathcal{K} > 0$.

Theorem 3.5

Let (X, D_b) be a cone b -metric space over Banach algebra \mathfrak{A} and \mathfrak{P} be a solid cone in \mathfrak{A} with the coefficient $\omega \geq 1$. $h_i \in \mathfrak{P} (i = 1, 2, \dots, 5)$ be a generalized Lipschitz constant with $2\omega\rho(h_5) + (\omega + 1)\rho(h_1 + h_2 + \omega h_3 + \omega h_4) < 2$. Suppose that h_5 commutes with $h_1 + h_2 + \omega h_3 + \omega h_4$ and the mappings $\mathfrak{S}, \mathfrak{L}: X \rightarrow \mathfrak{CB}(X)$ be generalized multi-valued (ϕ, \mathfrak{F}) -contraction mapping, satisfies that

$$\phi(\mathcal{H}(\mathfrak{S}\varrho, \mathfrak{L}\varsigma)) \leq \mathfrak{F}(\phi(\mathcal{M}(\varrho, \varsigma))) \dots \dots \dots (1.1)$$

where,

$$\mathcal{M}(\varrho, \varsigma) = h_1 \frac{D_b(\varrho, \mathfrak{S}\varrho)}{1 + D_b(\varrho, \mathfrak{S}\varrho)} + h_2 \frac{D_b(\varsigma, \mathfrak{L}\varsigma)}{1 + D_b(\varsigma, \mathfrak{L}\varsigma)} + h_3 \frac{D_b(\varrho, \mathfrak{L}\varsigma)}{1 + D_b(\varrho, \mathfrak{L}\varsigma)} + h_4 \frac{D_b(\varsigma, \mathfrak{S}\varrho)}{1 + D_b(\varsigma, \mathfrak{S}\varrho)} + h_5 D_b(\varrho, \varsigma)$$

where, $\mathfrak{F} \in \Psi$, $\phi \in \Phi$ such that $\forall \varrho, \varsigma \in X$. Moreover, if \mathfrak{S} and \mathfrak{L} are weakly compatible, then \mathfrak{S} and \mathfrak{L} have a unique common fixed point.

Proof. Fix any $\varrho \in X$. Define $\varrho_0 = \varrho$ and let $\varrho_1 \in \mathfrak{S}\varrho_0$, $\varrho_2 \in \mathfrak{L}\varrho_1$ such that $\varrho_{2n+1} = \mathfrak{S}\varrho_{2n}$, $\varrho_{2n+2} = \mathfrak{L}\varrho_{2n+1}$, by lemma 2.13, we may choose $\varrho_2 \in \mathfrak{L}\varrho_1$ such that $\phi(D_b(\varrho_1, \varrho_2)) \leq \phi(H(\mathfrak{S}\varrho_0, \mathfrak{L}\varrho_1)) + (h_1 + h_5 + \omega h_3)$

$$\begin{aligned}
\varphi(D_b(\varrho_1, \varrho_2)) &\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_0, \mathfrak{S}\varrho_0)}{1+D_b(\varrho_0, \mathfrak{S}\varrho_0)} + \mathfrak{h}_2 \frac{D_b(\varrho_1, \mathfrak{S}\varrho_1)}{1+D_b(\varrho_1, \mathfrak{S}\varrho_1)} + \mathfrak{h}_3 \frac{D_b(\varrho_0, \mathfrak{S}\varrho_1)}{1+D_b(\varrho_0, \mathfrak{S}\varrho_1)} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_1, \mathfrak{S}\varrho_0)}{1+D_b(\varrho_1, \mathfrak{S}\varrho_0)} + \mathfrak{h}_5 D_b(\varrho_0, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3) \\
&= \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_0, \varrho_1)}{1+D_b(\varrho_0, \varrho_1)} + \mathfrak{h}_2 \frac{D_b(\varrho_1, \varrho_2)}{1+D_b(\varrho_1, \varrho_2)} + \mathfrak{h}_3 \frac{D_b(\varrho_0, \varrho_2)}{1+D_b(\varrho_0, \varrho_2)} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_1, \varrho_1)}{1+D_b(\varrho_1, \varrho_1)} + \mathfrak{h}_5 D_b(\varrho_0, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3) \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(\varrho_0, \varrho_1) + \mathfrak{h}_2 D_b(\varrho_1, \varrho_2) + \mathfrak{h}_3 D_b(\varrho_0, \varrho_2) \\ &+ \mathfrak{h}_4 D_b(\varrho_1, \varrho_1) + \mathfrak{h}_5 D_b(\varrho_0, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3) \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(\varrho_0, \varrho_1) + \mathfrak{h}_2 D_b(\varrho_1, \varrho_2) \\ &+ \mathfrak{h}_3 \omega [D_b(\varrho_0, \varrho_1) + D_b(\varrho_1, \varrho_2)] \\ &+ \mathfrak{h}_5 D_b(\varrho_0, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3) \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &(\mathfrak{h}_1 + \omega \mathfrak{h}_3 + \mathfrak{h}_5) D_b(\varrho_0, \varrho_1) \\ &+ (\mathfrak{h}_2 + \omega \mathfrak{h}_3) D_b(\varrho_1, \varrho_2) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3)
\end{aligned}$$

which implies that

$$(e - \mathfrak{h}_2 - \omega \mathfrak{h}_3) \varphi(D_b(\varrho_1, \varrho_2)) \leq \left(\begin{aligned} &(\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3) \psi \left(\varphi(D_b(\varrho_0, \varrho_1)) \right) \\ &+ (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3) \end{aligned} \right) \quad (1.2)$$

Then,

$$\begin{aligned}
\varphi(D_b(\varrho_2, \varrho_1)) &\leq \varphi(H(\mathfrak{S}\varrho_1, \mathfrak{S}\varrho_0)) + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) \\
\varphi(D_b(\varrho_2, \varrho_1)) &\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_1, \mathfrak{S}\varrho_1)}{1+D_b(\varrho_1, \mathfrak{S}\varrho_1)} + \mathfrak{h}_2 \frac{D_b(\varrho_0, \mathfrak{S}\varrho_0)}{1+D_b(\varrho_0, \mathfrak{S}\varrho_0)} + \mathfrak{h}_3 \frac{D_b(\varrho_1, \mathfrak{S}\varrho_0)}{1+D_b(\varrho_1, \mathfrak{S}\varrho_0)} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_0, \mathfrak{S}\varrho_1)}{1+D_b(\varrho_0, \mathfrak{S}\varrho_1)} + \mathfrak{h}_5 D_b(\varrho_1, \varrho_0) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) \\
&= \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_1, \varrho_2)}{1+D_b(\varrho_1, \varrho_2)} + \mathfrak{h}_2 \frac{D_b(\varrho_0, \varrho_1)}{1+D_b(\varrho_0, \varrho_1)} + \mathfrak{h}_3 \frac{D_b(\varrho_1, \varrho_1)}{1+D_b(\varrho_1, \varrho_1)} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_0, \varrho_2)}{1+D_b(\varrho_0, \varrho_2)} + \mathfrak{h}_5 D_b(\varrho_0, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(\varrho_1, \varrho_2) + \mathfrak{h}_2 D_b(\varrho_0, \varrho_1) + \mathfrak{h}_3 D_b(\varrho_1, \varrho_1) \\ &+ \mathfrak{h}_4 D_b(\varrho_0, \varrho_2) + \mathfrak{h}_5 D_b(\varrho_0, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(\varrho_1, \varrho_2) + \mathfrak{h}_2 D_b(\varrho_0, \varrho_1) \\ &+ \mathfrak{h}_4 \omega [D_b(\varrho_0, \varrho_1) + D_b(\varrho_1, \varrho_2)] \\ &+ \mathfrak{h}_5 D_b(\varrho_0, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)
\end{aligned}$$

$$= \left(\psi \left(\varphi \left(\begin{array}{c} (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) D_b(\varrho_0, \varrho_1) \\ + (\mathfrak{h}_1 + \omega \mathfrak{h}_4) D_b(\varrho_2, \varrho_1) \\ + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) \end{array} \right) \right) \right)$$

which implies that

$$(e - \mathfrak{h}_1 - \omega \mathfrak{h}_4) \varphi(D_b(\varrho_2, \varrho_1)) \leq \left(\begin{array}{c} (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) \psi \left(\varphi(D_b(\varrho_0, \varrho_1)) \right) \\ + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) \end{array} \right) \quad (1.3)$$

Adding inequalities (1.2) and (1.3), we obtain $\varphi(D_b(\varrho_1, \varrho_2))$ where,

$$(2e - \mathfrak{h}_1 - \mathfrak{h}_2 - \omega \mathfrak{h}_3 - \omega \mathfrak{h}_4) \varphi(D_b(\varrho_1, \varrho_2)) \leq \left(\begin{array}{c} (2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2) \psi \left(\varphi(D_b(\varrho_0, \varrho_1)) \right) \\ + \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4 \\ + (2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2 + \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4) \end{array} \right) \quad (1.4)$$

Denote $\mathfrak{h}_1 + \mathfrak{h}_2 + \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4 = \mathfrak{h}$, then (1.4) yields that

$$(2e - \mathfrak{h}) \varphi(D_b(\varrho_1, \varrho_2)) \leq (2\mathfrak{h}_5 + \mathfrak{h}) \psi \left(\varphi(D_b(\varrho_0, \varrho_1)) \right) + (2\mathfrak{h}_5 + \mathfrak{h}) \quad (1.5)$$

Similarly, it can be shown that, there exists $\varrho_2 \in \mathfrak{S}\varrho_1$, $\varrho_3 \in \mathfrak{Q}\varrho_2$ such that

$$\begin{aligned} \varphi(D_b(\varrho_2, \varrho_3)) &\leq \varphi(H(\mathfrak{S}\varrho_1, \mathfrak{Q}\varrho_2)) + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2 \\ \varphi(D_b(\varrho_2, \varrho_3)) &\leq \left(\psi \left(\varphi \left(\begin{array}{c} \mathfrak{h}_1 \frac{D_b(\varrho_1, \mathfrak{S}\varrho_1)}{1+D_b(\varrho_1, \mathfrak{S}\varrho_1)} + \mathfrak{h}_2 \frac{D_b(\varrho_2, \mathfrak{Q}\varrho_2)}{1+D_b(\varrho_2, \mathfrak{Q}\varrho_2)} + \mathfrak{h}_3 \frac{D_b(\varrho_1, \mathfrak{Q}\varrho_2)}{1+D_b(\varrho_1, \mathfrak{Q}\varrho_2)} \\ + \mathfrak{h}_4 \frac{D_b(\varrho_2, \mathfrak{S}\varrho_1)}{1+D_b(\varrho_2, \mathfrak{S}\varrho_1)} + \mathfrak{h}_5 D_b(\varrho_1, \varrho_2) \\ + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3)^2 \end{array} \right) \right) \right) \\ &= \left(\psi \left(\varphi \left(\begin{array}{c} \mathfrak{h}_1 \frac{D_b(\varrho_1, \varrho_2)}{1+D_b(\varrho_1, \varrho_2)} + \mathfrak{h}_2 \frac{D_b(\varrho_2, \varrho_3)}{1+D_b(\varrho_2, \varrho_3)} + \mathfrak{h}_3 \frac{D_b(\varrho_1, \varrho_3)}{1+D_b(\varrho_1, \varrho_3)} \\ + \mathfrak{h}_4 \frac{D_b(\varrho_2, \varrho_2)}{1+D_b(\varrho_2, \varrho_2)} + \mathfrak{h}_5 D_b(\varrho_1, \varrho_2) \\ + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3)^2 \end{array} \right) \right) \right) \\ &\leq \left(\psi \left(\varphi \left(\begin{array}{c} \mathfrak{h}_1 D_b(\varrho_1, \varrho_2) + \mathfrak{h}_2 D_b(\varrho_2, \varrho_3) + \mathfrak{h}_3 D_b(\varrho_1, \varrho_3) \\ + \mathfrak{h}_4 D_b(\varrho_2, \varrho_2) + \mathfrak{h}_5 D_b(\varrho_1, \varrho_2) \\ + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3)^2 \end{array} \right) \right) \right) \\ &\leq \left(\psi \left(\varphi \left(\begin{array}{c} \mathfrak{h}_1 D_b(\varrho_1, \varrho_2) + \mathfrak{h}_2 D_b(\varrho_2, \varrho_3) \\ + \mathfrak{h}_3 \omega [D_b(\varrho_1, \varrho_2) + D_b(\varrho_2, \varrho_3)] \\ + \mathfrak{h}_5 D_b(\varrho_1, \varrho_2) \\ + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3)^2 \end{array} \right) \right) \right) \\ &\leq \left(\psi \left(\varphi \left(\begin{array}{c} (\mathfrak{h}_1 + \omega \mathfrak{h}_3 + \mathfrak{h}_5) D_b(\varrho_1, \varrho_2) \\ + (\mathfrak{h}_2 + \omega \mathfrak{h}_3) D_b(\varrho_2, \varrho_3) \\ + (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3)^2 \end{array} \right) \right) \right) \end{aligned}$$

which implies that

$$(e - \mathfrak{h}_2 - \omega \mathfrak{h}_3) \varphi(D_b(\varrho_1, \varrho_2)) \leq \left(\begin{array}{c} (\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3)^2 \psi \left(\varphi(D_b(\varrho_0, \varrho_1)) \right) \\ + 2(\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3)^2 \end{array} \right) \quad (1.6)$$

Then,

$$\varphi(D_b(\varrho_3, \varrho_2)) \leq \varphi(H(\mathfrak{Q}\varrho_2, \mathfrak{S}\varrho_1)) + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2$$

$$\begin{aligned}
\varphi(D_b(\varrho_3, \varrho_2)) &\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_2, \mathfrak{L}\varrho_2)}{1+D_b(\varrho_2, \mathfrak{L}\varrho_2)} + \mathfrak{h}_2 \frac{D_b(\varrho_1, \mathfrak{S}\varrho_1)}{1+D_b(\varrho_1, \mathfrak{S}\varrho_1)} + \mathfrak{h}_3 \frac{D_b(\varrho_2, \mathfrak{S}\varrho_1)}{1+D_b(\varrho_2, \mathfrak{S}\varrho_1)} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_1, \mathfrak{L}\varrho_2)}{1+D_b(\varrho_1, \mathfrak{L}\varrho_2)} + \mathfrak{h}_5 D_b(\varrho_2, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2 \\
&= \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_2, \varrho_3)}{1+D_b(\varrho_2, \varrho_3)} + \mathfrak{h}_2 \frac{D_b(\varrho_1, \varrho_2)}{1+D_b(\varrho_1, \varrho_2)} + \mathfrak{h}_3 \frac{D_b(\varrho_2, \varrho_2)}{1+D_b(\varrho_2, \varrho_2)} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_1, \varrho_3)}{1+D_b(\varrho_1, \varrho_3)} + \mathfrak{h}_5 D_b(\varrho_2, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2 \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(\varrho_2, \varrho_3) + \mathfrak{h}_2 D_b(\varrho_1, \varrho_2) + \mathfrak{h}_3 D_b(\varrho_2, \varrho_2) \\ &+ \mathfrak{h}_4 D_b(\varrho_1, \varrho_3) + \mathfrak{h}_5 D_b(\varrho_2, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2 \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(\varrho_2, \varrho_3) + \mathfrak{h}_2 D_b(\varrho_1, \varrho_2) \\ &+ \mathfrak{h}_4 \omega [D_b(\varrho_1, \varrho_2) + D_b(\varrho_2, \varrho_3)] \\ &+ \mathfrak{h}_5 D_b(\varrho_2, \varrho_1) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2 \\
&= \left(\psi \left(\varphi \left(\begin{aligned} &(\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) D_b(\varrho_1, \varrho_2) \\ &+ (\mathfrak{h}_1 + \omega \mathfrak{h}_4) D_b(\varrho_2, \varrho_3) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2
\end{aligned}$$

which implies that

$$(e - \mathfrak{h}_1 - \omega \mathfrak{h}_4) \varphi(D_b(\varrho_3, \varrho_2)) \leq \left(\begin{aligned} &(\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2 \psi(\varphi(D_b(\varrho_0, \varrho_1))) \\ &+ 2(\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^2 \end{aligned} \right) \quad (1.7)$$

Adding inequalities (1.6) and (1.3), we obtain $\varphi(D_b(\varrho_1, \varrho_2))$ where,

$$\left(\begin{aligned} &2e - \mathfrak{h}_1 - \mathfrak{h}_2 \\ &-\omega \mathfrak{h}_3 - \omega \mathfrak{h}_4 \end{aligned} \right) \varphi(D_b(\varrho_2, \varrho_3)) \leq \left(\begin{aligned} &\left(\begin{aligned} &(2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2)^2 \\ &+ \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4 \end{aligned} \right) \psi(\varphi(D_b(\varrho_0, \varrho_1))) \\ &+ 2(2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2 + \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4)^2 \end{aligned} \right) \quad (1.8)$$

Denote $\mathfrak{h}_1 + \mathfrak{h}_2 + \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4 = \mathfrak{h}$, then (1.8) yields that

$$(2e - \mathfrak{h}) \varphi(D_b(\varrho_1, \varrho_2)) \leq (2\mathfrak{h}_5 + \mathfrak{h})^2 \psi(\varphi(D_b(\varrho_0, \varrho_1))) + 2(2\mathfrak{h}_5 + \mathfrak{h})^2 \quad (1.9)$$

Continuing this process, we obtain by induction a sequence $\{\varrho_n\}$ such that $\varrho_{2n+1} \in \mathfrak{S}\varrho_{2n}$, $\varrho_{2n+2} \in \mathfrak{L}\varrho_{2n+1}$ such that

$$\begin{aligned}
\varphi(D_b(\varrho_{2n+1}, \varrho_{2n+2})) &\leq \varphi(H(\mathfrak{S}\varrho_{2n}, \mathfrak{L}\varrho_{2n+1})) + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_{2n}, \mathfrak{S}\varrho_{2n})}{1+D_b(\varrho_{2n}, \mathfrak{S}\varrho_{2n})} + \mathfrak{h}_2 \frac{D_b(\varrho_{2n+1}, \mathfrak{L}\varrho_{2n+1})}{1+D_b(\varrho_{2n+1}, \mathfrak{L}\varrho_{2n+1})} + \mathfrak{h}_3 \frac{D_b(\varrho_{2n}, \mathfrak{L}\varrho_{2n+1})}{1+D_b(\varrho_{2n}, \mathfrak{L}\varrho_{2n+1})} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_{2n+1}, \mathfrak{S}\varrho_{2n})}{1+D_b(\varrho_{2n+1}, \mathfrak{S}\varrho_{2n})} + \mathfrak{h}_5 D_b(\varrho_{2n}, \varrho_{2n+1}) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \\
&= \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_{2n}, \varrho_{2n+1})}{1+D_b(\varrho_{2n}, \varrho_{2n+1})} + \mathfrak{h}_2 \frac{D_b(\varrho_{2n+1}, \varrho_{2n+2})}{1+D_b(\varrho_{2n+1}, \varrho_{2n+2})} + \mathfrak{h}_3 \frac{D_b(\varrho_{2n}, \varrho_{2n+2})}{1+D_b(\varrho_{2n}, \varrho_{2n+2})} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_{2n+1}, \varrho_{2n+1})}{1+D_b(\varrho_{2n+1}, \varrho_{2n+1})} + \mathfrak{h}_5 D_b(\varrho_{2n}, \varrho_{2n+1}) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1}
\end{aligned}$$

$$\begin{aligned}
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(\varrho_{2n}, \varrho_{2n+1}) + \mathfrak{h}_2 D_b(\varrho_{2n+1}, \varrho_{2n+2}) \\ &+ \mathfrak{h}_3 \omega [D_b(\varrho_{2n}, \varrho_{2n+1}) + D_b(\varrho_{2n+1}, \varrho_{2n+2})] \\ &+ \mathfrak{h}_5 D_b(\varrho_{2n}, \varrho_{2n+1}) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \\
&= \left(\psi \left(\varphi \left(\begin{aligned} &(\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3) D_b(\varrho_{2n}, \varrho_{2n+1}) \\ &+ (\mathfrak{h}_2 + \omega \mathfrak{h}_3) D_b(\varrho_{2n+1}, \varrho_{2n+2}) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1}
\end{aligned}$$

which implies that

$$(e - \mathfrak{h}_2 - \omega \mathfrak{h}_3) \varphi(D_b(\varrho_{2n+1}, \varrho_{2n+2})) \leq \left(\begin{aligned} &(\mathfrak{h}_1 + \mathfrak{h}_5 + \omega \mathfrak{h}_3) \psi \left(\varphi(D_b(\varrho_{2n}, \varrho_{2n+1})) \right) \\ &+ (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \end{aligned} \right) \quad (1.10)$$

Also,

$$\begin{aligned}
\varphi(D_b(\varrho_{2n+2}, \varrho_{2n+1})) &\leq \varphi(H(\mathfrak{L}\varrho_{2n+1}, \mathfrak{S}\varrho_{2n})) + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_{2n+1}, \mathfrak{L}\varrho_{2n+1})}{1+D_b(\varrho_{2n+1}, \mathfrak{L}\varrho_{2n+1})} + \mathfrak{h}_2 \frac{D_b(\varrho_{2n}, \mathfrak{S}\varrho_{2n})}{1+D_b(\varrho_{2n}, \mathfrak{S}\varrho_{2n})} + \mathfrak{h}_3 \frac{D_b(\varrho_{2n+1}, \mathfrak{S}\varrho_{2n})}{1+D_b(\varrho_{2n+1}, \mathfrak{S}\varrho_{2n})} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_{2n}, \mathfrak{L}\varrho_{2n+1})}{1+D_b(\varrho_{2n}, \mathfrak{L}\varrho_{2n+1})} + \mathfrak{h}_5 D_b(\varrho_{2n+1}, \varrho_{2n}) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \\
&= \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_{2n+1}, \varrho_{2n+2})}{1+D_b(\varrho_{2n+1}, \varrho_{2n+2})} + \mathfrak{h}_2 \frac{D_b(\varrho_{2n}, \varrho_{2n+1})}{1+D_b(\varrho_{2n}, \varrho_{2n+1})} + \mathfrak{h}_3 \frac{D_b(\varrho_{2n+1}, \varrho_{2n+1})}{1+D_b(\varrho_{2n+1}, \varrho_{2n+1})} \\ &+ \mathfrak{h}_4 \frac{D_b(\varrho_{2n}, \varrho_{2n+2})}{1+D_b(\varrho_{2n}, \varrho_{2n+2})} + \mathfrak{h}_5 D_b(\varrho_{2n+1}, \varrho_{2n}) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \\
&\leq \left(\psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(\varrho_{2n+1}, \varrho_{2n+2}) + \mathfrak{h}_2 D_b(\varrho_{2n}, \varrho_{2n+1}) \\ &+ \mathfrak{h}_4 \omega [D_b(\varrho_{2n}, \varrho_{2n+1}) + D_b(\varrho_{2n+1}, \varrho_{2n+2})] \\ &+ \mathfrak{h}_5 D_b(\varrho_{2n+1}, \varrho_{2n}) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \\
&= \left(\psi \left(\varphi \left(\begin{aligned} &(\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) D_b(\varrho_{2n}, \varrho_{2n+1}) \\ &+ (\mathfrak{h}_1 + \omega \mathfrak{h}_4) D_b(\varrho_{2n+1}, \varrho_{2n+2}) \end{aligned} \right) \right) \right) \\
&\quad + (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1}
\end{aligned}$$

$$(e - \mathfrak{h}_1 + \omega \mathfrak{h}_4) \varphi(D_b(\varrho_{2n+2}, \varrho_{2n+1})) \leq \left(\begin{aligned} &(\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4) \psi \left(\varphi(D_b(\varrho_{2n}, \varrho_{2n+1})) \right) \\ &+ (\mathfrak{h}_2 + \mathfrak{h}_5 + \omega \mathfrak{h}_4)^{2n+1} \end{aligned} \right) \quad (1.11)$$

Add up (1.10) and (1.11) yields that

$$\left(\begin{aligned} &(2e - \mathfrak{h}_1 - \mathfrak{h}_2) \\ &(-\omega \mathfrak{h}_3 - \omega \mathfrak{h}_4) \end{aligned} \right) \varphi(D_b(\varrho_{2n+1}, \varrho_{2n+2})) \leq \left(\begin{aligned} &((2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2) \\ &+ \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4) \psi \left(\varphi(D_b(\varrho_{2n}, \varrho_{2n+1})) \right) \\ &+ (2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2 + \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4)^{2n+1} \end{aligned} \right) \quad (1.12)$$

Denote $\mathfrak{h}_1 + \mathfrak{h}_2 + \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4 = \mathfrak{h}$, then (1.12) yields that

$$(2e - \mathfrak{h}) \varphi(D_b(\varrho_{2n+1}, \varrho_{2n+2})) \leq \left(\begin{aligned} &(2\mathfrak{h}_5 + \mathfrak{h}) \psi \left(\varphi(D_b(\varrho_{2n}, \varrho_{2n+1})) \right) \\ &+ (2\mathfrak{h}_5 + \mathfrak{h})^{2n+1} \end{aligned} \right) \quad (1.13)$$

Therefore,

$$\varphi(D_b(\varrho_n, \varrho_{n+1})) \leq \varphi(H(\mathfrak{S}\varrho_{n-1}, \mathfrak{L}\varrho_n)) + (2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2 + \omega \mathfrak{h}_3 + \omega \mathfrak{h}_4)^n$$

$$\begin{pmatrix} (2e - \mathfrak{h}_1 - \mathfrak{h}_2) \\ -\omega\mathfrak{h}_3 - \omega\mathfrak{h}_4 \end{pmatrix} \varphi(D_b(\varrho_n, \varrho_{n+1})) \leq \begin{pmatrix} (2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2) \psi(\varphi(D_b(\varrho_{n-1}, \varrho_n))) \\ +\omega\mathfrak{h}_3 + \omega\mathfrak{h}_4 \\ + (2\mathfrak{h}_5 + \mathfrak{h}_1 + \mathfrak{h}_2)^n \\ +\omega\mathfrak{h}_3 + \omega\mathfrak{h}_4 \end{pmatrix} \quad (1.14)$$

Denote $\mathfrak{h}_1 + \mathfrak{h}_2 + \omega\mathfrak{h}_3 + \omega\mathfrak{h}_4 = \mathfrak{h}$, then (1.14) yields that

$$(2e - \mathfrak{h})\varphi(D_b(\varrho_n, \varrho_{n+1})) \leq (2\mathfrak{h}_5 + \mathfrak{h})\psi(\varphi(D_b(\varrho_{n-1}, \varrho_n))) + (2\mathfrak{h}_5 + \mathfrak{h})^n \quad (1.15)$$

Note that

$2\rho(\mathfrak{h}) \leq (\omega + 1)\rho(\mathfrak{h}) \leq 2\omega\rho(\mathfrak{h}_5) + (\omega + 1)\rho(\mathfrak{h}) < 2$
 $\rho(\mathfrak{h}) < 1 < 2$, then by lemma 2.16 it follows that $(2e - \mathfrak{h})$ is invertible. Furthermore,

$$(2e - \mathfrak{h})^{-1} = \sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{2^{i+1}}$$

By multiplying in both sides of (2.15) by $(2e - \mathfrak{h})^{-1}$, we get

$$\varphi(D_b(\varrho_n, \varrho_{n+1})) \leq \begin{pmatrix} (2e - \mathfrak{h})^{-1}(2\mathfrak{h}_5 + \mathfrak{h})\psi(\varphi(D_b(\varrho_{n-1}, \varrho_n))) \\ + (2e - \mathfrak{h})^{-1}(2\mathfrak{h}_5 + \mathfrak{h})^n \end{pmatrix} \quad (1.16)$$

Let $\gamma = (2e - \mathfrak{h})^{-1}(2\mathfrak{h}_5 + \mathfrak{h})$, then by (1.16) we get

$$\begin{aligned} \varphi(D_b(\varrho_n, \varrho_{n+1})) &\leq \psi(\gamma\varphi(D_b(\varrho_{n-1}, \varrho_n))) + \gamma^n \\ &\leq \gamma\psi(\gamma\varphi(D_b(\varrho_{n-2}, \varrho_{n-1}))) + 2\gamma^n \\ &= \psi(\gamma^2\varphi(D_b(\varrho_{n-2}, \varrho_{n-1}))) + 2\gamma^n \\ &\vdots \\ &\leq \gamma^n\psi(\varphi(D_b(\varrho_0, \varrho_1))) + n\gamma^n. \end{aligned}$$

Since \mathfrak{h}_5 commutes with \mathfrak{h} , it follows that

$$\begin{aligned} (2e - \mathfrak{h})^{-1}(2\mathfrak{h}_5 + \mathfrak{h}) &= \left(\sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{2^{i+1}}\right)(2\mathfrak{h}_5 + \mathfrak{h}) \\ &= 2\left(\sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{2^{i+1}}\right)\mathfrak{h}_5 + \left(\sum_{i=0}^{\infty} \frac{\mathfrak{h}^{i+1}}{2^{i+1}}\right) \\ &= 2\mathfrak{h}\left(\sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{2^{i+1}}\right) + \mathfrak{h}\left(\sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{2^{i+1}}\right) \\ &= (2\mathfrak{h}_5 + \mathfrak{h})\left(\sum_{i=0}^{\infty} \frac{\mathfrak{h}^i}{2^{i+1}}\right) = (2\mathfrak{h}_5 + \mathfrak{h})(2e - \mathfrak{h})^{-1} \end{aligned}$$

then, $(2e - \mathfrak{h})^{-1}$ commutes with $(2\mathfrak{h}_5 + \mathfrak{h})$. Note by Lemma 2.5 and Lemma 3.1 that

$$\begin{aligned} \rho(\gamma) &= \rho((2\mathfrak{h}_5 + \mathfrak{h})(2e - \mathfrak{h})^{-1}) \\ &\leq \rho((2e - \mathfrak{h})^{-1})\rho((2\mathfrak{h}_5 + \mathfrak{h})) \\ &\leq \frac{1}{2 - \rho(\mathfrak{h})}[2\rho(\mathfrak{h}_5) + \rho(\mathfrak{h})] \\ &= \frac{1}{2 - \rho(\mathfrak{h}_1 + \mathfrak{h}_2 + \omega\mathfrak{h}_3 + \omega\mathfrak{h}_4)}[2\rho(\mathfrak{h}_5) + \rho(\mathfrak{h}_1 + \mathfrak{h}_2 + \omega\mathfrak{h}_3 + \omega\mathfrak{h}_4)] \\ &< \frac{1}{\omega}, \quad [\text{since } 2\omega\rho(\mathfrak{h}_5) + (\omega + 1)(\rho(\mathfrak{h}_1) + \rho(\mathfrak{h}_2) + \omega\rho(\mathfrak{h}_3) + \omega\rho(\mathfrak{h}_4)) < 2] \end{aligned}$$

which establishes that $(e - \omega\gamma)$ is invertible and $\|\gamma^m\| \rightarrow 0 (m \rightarrow \infty)$. Hence, for any $m \geq 1$,

$p \geq 1$ and $\gamma \in \mathfrak{F}$ with $\rho(\gamma) < \frac{1}{\omega}$, we have that

$$\phi(D_b(\varrho_m, \varrho_{m+p})) \leq \psi(\phi\omega[D_b(\varrho_m, \varrho_{m+1}) + D_b(\varrho_{m+1}, \varrho_{m+p})])$$

$$\begin{aligned}
&\leq \omega\psi(\phi D_b(\varrho_m, \varrho_{m+1})) + \omega^2\psi\left(\phi \begin{bmatrix} D_b(\varrho_{m+1}, \varrho_{m+2}) \\ +D_b(\varrho_{m+2}, \varrho_{m+p}) \end{bmatrix}\right) \\
&\leq \left(\begin{aligned} &\omega\psi(\phi D_b(\varrho_m, \varrho_{m+1})) + \omega^2\psi(\phi D_b(\varrho_{m+1}, \varrho_{m+2})) \\ &+ \omega^3\psi(\phi D_b(\varrho_{m+2}, \varrho_{m+3})) + \dots \\ &+ \omega^{p-1}\psi(\phi D_b(\varrho_{m+p-2}, \varrho_{m+p-1})) \\ &+ \omega^{p-1}\psi(\phi D_b(\varrho_{m+p-1}, \varrho_{m+p})) \end{aligned} \right) \\
&\leq \left(\begin{aligned} &\omega\gamma^m\psi(\phi D_b(\varrho_0, \varrho_1)) + \omega^2\gamma^{m+1}\psi(\phi D_b(\varrho_0, \varrho_1)) \\ &+ \omega^3\gamma^{m+2}\psi(\phi D_b(\varrho_0, \varrho_1)) + \dots \\ &+ \omega^{p-1}\gamma^{m+p-2}\psi(\phi D_b(\varrho_0, \varrho_1)) \\ &+ \omega^p\gamma^{m+p-1}\psi(\phi D_b(\varrho_0, \varrho_1)) \end{aligned} \right) \\
&= \omega\gamma^m[e + \omega\gamma + \omega^2\gamma^2 + \dots + (\omega\gamma)^{p-1}]\psi(\phi D_b(\varrho_0, \varrho_1)) \\
&\leq \omega\gamma^m(e - \omega\gamma)^{-1}\psi(\phi D_b(\varrho_0, \varrho_1)).
\end{aligned}$$

In view, $\|\omega\gamma^m\psi(\phi D_b(\varrho_0, \varrho_1))\| \leq \|\omega\gamma^m\| \|\psi(\phi D_b(\varrho_0, \varrho_1))\| \rightarrow 0$ ($m \rightarrow \infty$), by lemma 2.4, we have $\{\omega\gamma^m\psi(\phi D_b(\varrho_0, \varrho_1))\}$ is a \mathfrak{c} -sequence. Next by using Lemma 3.2 and lemma 2.15, we conclude that $\{\varrho_n\}$ is a Cauchy sequence. Since (X, D_b) is complete, there exists $\mathfrak{s} \in X$ such that $\varrho_n \rightarrow \mathfrak{s}$. We shall prove that \mathfrak{s} is a common fixed point of \mathfrak{S} and \mathfrak{Q} .

$$\begin{aligned}
\varphi(D_b(\mathfrak{s}, \mathfrak{S}\mathfrak{s})) &\leq \psi(\varphi(\omega [D_b(\mathfrak{s}, \varrho_{2n+1}) + D_b(\varrho_{2n+1}, \mathfrak{S}\mathfrak{s})])) \\
&\leq \psi(\varphi(\omega [D_b(\mathfrak{s}, \varrho_{2n+1}) + \mathcal{H}(\varrho_{2n+1}, \mathfrak{S}\mathfrak{s})])) \\
\varphi(D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})) &\leq \psi(\varphi(\omega [D_b(\mathfrak{s}, \varrho_{2n+1}) + D_b(\varrho_{2n+1}, \mathfrak{Q}\mathfrak{s})])) \quad (1.17) \\
&\leq \psi(\varphi(\omega [D_b(\mathfrak{s}, \varrho_{2n+1}) + \mathcal{H}(\varrho_{2n}, \mathfrak{Q}\mathfrak{s})]))
\end{aligned}$$

Where,

$$\varphi(\mathcal{H}(\varrho_{2n}, \mathfrak{Q}\mathfrak{s})) \leq \psi\left(\varphi\left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\varrho_{2n}, \mathfrak{S}\varrho_{2n})}{1+D_b(\varrho_{2n}, \mathfrak{S}\varrho_{2n})} + \mathfrak{h}_2 \frac{D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})}{1+D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})} \\ &+ \mathfrak{h}_3 \frac{D_b(\varrho_{2n}, \mathfrak{Q}\mathfrak{s})}{1+D_b(\varrho_{2n}, \mathfrak{Q}\mathfrak{s})} + \mathfrak{h}_4 \frac{D_b(\mathfrak{s}, \mathfrak{S}\varrho_{2n})}{1+D_b(\mathfrak{s}, \mathfrak{S}\varrho_{2n})} + \mathfrak{h}_5 D_b(\varrho_{2n}, \mathfrak{s}) \end{aligned}\right)\right) \quad (1.18)$$

Using (1.18) in (1.17) and letting as $n \rightarrow \infty$, we obtain,

$$\begin{aligned}
\varphi(D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})) &\leq \psi(\varphi(\omega D_b(\mathfrak{s}, \mathfrak{s}))) \\
&\leq \psi\left(\varphi\left(\omega \left[\begin{aligned} &\mathfrak{h}_1 \frac{D_b(\mathfrak{s}, \mathfrak{s})}{1+D_b(\mathfrak{s}, \mathfrak{s})} + \mathfrak{h}_2 \frac{D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})}{1+D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})} + \mathfrak{h}_3 \frac{D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})}{1+D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})} \\ &+ \mathfrak{h}_4 \frac{D_b(\mathfrak{s}, \mathfrak{s})}{1+D_b(\mathfrak{s}, \mathfrak{s})} + \mathfrak{h}_5 D_b(\mathfrak{s}, \mathfrak{s}) \end{aligned}\right]\right)\right) \\
&= \psi\left(\varphi\left(\omega \left[\begin{aligned} &\mathfrak{h}_1 D_b(\mathfrak{s}, \mathfrak{s}) + \mathfrak{h}_2 D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s}) + \mathfrak{h}_3 D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s}) \\ &+ \mathfrak{h}_4 D_b(\mathfrak{s}, \mathfrak{s}) + \mathfrak{h}_5 D_b(\mathfrak{s}, \mathfrak{s}) \end{aligned}\right]\right)\right) \\
&= \psi(\varphi(\omega [\mathfrak{h}_2 D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s}) + \mathfrak{h}_3 D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})])) \\
&\leq \psi\left(\varphi(\omega (\mathfrak{h}_2 + \mathfrak{h}_3) D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s}))\right)
\end{aligned}$$

Which implies that

$$\psi(\phi D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})) \leq 0$$

Then by Lemma 2.2, we deduce that $\psi(\phi D_b(\mathfrak{s}, \mathfrak{Q}\mathfrak{s})) = 0$, that is $\mathfrak{Q}(\mathfrak{s}) = \mathfrak{s}$. Similarly, $\mathfrak{S}(\mathfrak{s}) = \mathfrak{s}$.

Hence $\mathfrak{S}s = \mathfrak{L}s = r$. In the following we shall prove \mathfrak{L} and \mathfrak{S} have a unique point of coincidence. Such that $s \neq s^*$ then from (1.1) we have

$$\begin{aligned}
\varphi(D_b(s, s^*)) &\leq \varphi(\mathcal{H}(\mathfrak{S}s, \mathfrak{L}s^*)) \\
&\leq \psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{D_b(s, \mathfrak{S}s)}{1 + D_b(s, \mathfrak{S}s)} + \mathfrak{h}_2 \frac{D_b(s^*, \mathfrak{L}s^*)}{1 + D_b(s^*, \mathfrak{L}s^*)} + \mathfrak{h}_3 \frac{D_b(s, \mathfrak{L}s^*)}{1 + D_b(s, \mathfrak{L}s^*)} \\ &+ \mathfrak{h}_4 \frac{D_b(s^*, \mathfrak{S}s)}{1 + D_b(s^*, \mathfrak{S}s)} + \mathfrak{h}_5 D_b(s, s^*) \end{aligned} \right) \right) \\
&\leq \psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 D_b(s, \mathfrak{S}s) + \mathfrak{h}_2 D_b(s^*, \mathfrak{L}s^*) + \mathfrak{h}_3 D_b(s, \mathfrak{L}s^*) \\ &+ \mathfrak{h}_4 D_b(s^*, \mathfrak{S}s) + \mathfrak{h}_5 D_b(s, s^*) \end{aligned} \right) \right) \\
&\leq \psi \left(\varphi(\mathfrak{h}_3 D_b(s, s^*) + \mathfrak{h}_4 D_b(s, s^*) + \mathfrak{h}_5 D_b(s, s^*)) \right) \\
&= (\mathfrak{h}_3 + \mathfrak{h}_4 + \mathfrak{h}_5) \psi(\phi D_b(s, s^*))
\end{aligned}$$

Set $(\mathfrak{h}_3 + \mathfrak{h}_4 + \mathfrak{h}_5) = \zeta$, then it follows that

$$\varphi(D_b(s, s^*)) \leq \zeta \psi(\varphi(D_b(s, s^*))) \leq \dots \leq \zeta^n \psi(\varphi(D_b(s, s^*))) \quad (1.19)$$

Because of

$$2\rho(\mathfrak{h}_5) + 2\rho(\mathfrak{h}) \leq 2\omega\rho(\mathfrak{h}_5) + (\omega + 1)\rho(\mathfrak{h}) < 2,$$

It follows that $\rho(\mathfrak{h}_5) + \rho(\mathfrak{h}) < 1$. Since \mathfrak{h}_5 commutes with \mathfrak{h} , then by Lemma 2.5,

$$\rho(\mathfrak{h}_5 + \mathfrak{h}) \leq \rho(\mathfrak{h}_5) + \rho(\mathfrak{h}) < 1.$$

Accordingly, by Lemma 2.5, we speculate that $\{(\mathfrak{h}_5 + \mathfrak{h})^n\}$ is a c -sequence. Noticing that $\zeta \leq \mathfrak{h}_5 + \mathfrak{h}$ leads to $\zeta^n \leq (\mathfrak{h}_5 + \mathfrak{h})^n$, we claim that $\{\zeta^n\}$ is a c -sequence. Consequently, in view (1.19), it easy to see $\psi(\phi D_b(s, s^*)) = 0$, that is $s = s^*$.

Finally, if $(\mathfrak{S}, \mathfrak{L})$ is weakly compatible, then Lemma 2.19, we claim that \mathfrak{L} and \mathfrak{S} have a unique common fixed point.

Corollary.3.6

Let (X, D_b) be a cone b -metric space over Banach algebra \mathfrak{A} and \mathfrak{B} be a solid cone in \mathfrak{A} with the coefficient $\omega \geq 1$. $\mathfrak{h}_i \in \mathfrak{B}$ ($i = 1, 2, \dots, 4$) be a generalized Lipschitz constant with $2\omega\rho(\mathfrak{h}_4) + (\omega + 1)\rho(\mathfrak{h}_1 + \omega\mathfrak{h}_2 + \omega\mathfrak{h}_3) < 2$. Suppose that \mathfrak{h}_4 commutes with $\mathfrak{h}_1 + \omega\mathfrak{h}_2 + \omega\mathfrak{h}_3$ and the mappings $\mathfrak{S}, \mathfrak{L} : X \rightarrow \mathfrak{CB}(X)$ be generalized multi-valued (ϕ, \mathfrak{F}) -contraction mapping, satisfies that

$$\phi(\mathcal{H}(\mathfrak{S}\varrho, \mathfrak{L}\varsigma)) \leq \mathfrak{F}(\phi(\mathcal{M}(\varrho, \varsigma)))$$

where,

$$\mathcal{M}(\varrho, \varsigma) = \mathfrak{h}_1 \frac{D_b(\varrho, \mathfrak{S}\varrho)}{1 + D_b(\varrho, \mathfrak{S}\varrho)} + \mathfrak{h}_2 \frac{D_b(\varrho, \mathfrak{L}\varsigma)}{1 + D_b(\varrho, \mathfrak{L}\varsigma)} + \mathfrak{h}_3 \frac{D_b(\varsigma, \mathfrak{S}\varrho)}{1 + D_b(\varsigma, \mathfrak{S}\varrho)} + \mathfrak{h}_4 D_b(\varrho, \varsigma)$$

where, $\mathfrak{F} \in \Psi$, $\phi \in \Phi$ such that $\forall \varrho, \varsigma \in X$. Moreover, if \mathfrak{S} and \mathfrak{L} are weakly compatible, then \mathfrak{S} and \mathfrak{L} have a unique common fixed point.

Proof. Choose $\mathfrak{h}_1 = \mathfrak{h}_3 = \mathfrak{h}_4 = \mathfrak{h}_5 = \mathfrak{h}$ and $\mathfrak{h}_2 = 0$ in theorem 3.5, the proof is valid.

Corollary.3.7

Let (X, D_b) be a cone b -metric space over Banach algebra \mathfrak{A} and \mathfrak{B} be a solid cone in \mathfrak{A} with the coefficient $\omega \geq 1$. $\mathfrak{h}_i \in \mathfrak{B}$ ($i = 1, 2, \dots, 4$) be a generalized nonnegative real constant with

$2\omega(\mathfrak{h}_4) + (\omega + 1)(\mathfrak{h}_1 + \omega \mathfrak{h}_2 + \omega \mathfrak{h}_3) < 2$. Let mappings $\mathfrak{S}, \mathfrak{L} : X \rightarrow \mathfrak{CB}(X)$ be generalized multi-valued (ϕ, \mathfrak{F}) -contraction mapping, satisfies that

$$\phi(\mathcal{H}(\mathfrak{S}\varrho, \mathfrak{L}\varsigma)) \leq \mathfrak{F}(\phi(\mathfrak{h}_1 D_b(\varrho, \mathfrak{S}\varrho) + \mathfrak{h}_2 D_b(\varrho, \mathfrak{L}\varsigma) + \mathfrak{h}_3 D_b(\varsigma, \mathfrak{S}\varrho) + \mathfrak{h}_4 D_b(\varrho, \varsigma)))$$

where, $\mathfrak{F} \in \Psi$, $\phi \in \Phi$ such that $\forall \varrho, \varsigma \in X$. Moreover, if \mathfrak{S} and \mathfrak{L} are weakly compatible, then \mathfrak{S} and \mathfrak{L} have a unique common fixed point.

Proof. Taking $\mathfrak{h}_1, \mathfrak{h}_3, \mathfrak{h}_4, \mathfrak{h}_5 \in \mathbb{R}^+$ in theorem 3.5, we obtain the desired result.

Corollary.3.8

Let (X, D_b) be a cone b -metric space over Banach algebra \mathfrak{A} and \mathfrak{B} be a solid cone in \mathfrak{A} with the coefficient $\omega \geq 1$. $\mathfrak{h}_i \in \mathfrak{B}$ be a generalized Lipschitz constant with $\rho(\mathfrak{h}) < \frac{1}{\omega^2 + \omega}$. Suppose that the mappings $\mathfrak{S}, \mathfrak{L} : X \rightarrow \mathfrak{CB}(X)$ be generalized multi-valued (ϕ, \mathfrak{F}) -contraction mapping, satisfies that

$$\phi(\mathcal{H}(\mathfrak{S}\varrho, \mathfrak{L}\varsigma)) \leq \mathfrak{F}(\phi(\mathfrak{h}(D_b(\varrho, \mathfrak{L}\varsigma) + D_b(\varsigma, \mathfrak{S}\varrho)))$$

where, $\mathfrak{F} \in \Psi$, $\phi \in \Phi$ such that $\forall \varrho, \varsigma \in X$. Moreover, if \mathfrak{S} and \mathfrak{L} are weakly compatible, then \mathfrak{S} and \mathfrak{L} have a unique common fixed point.

Proof. Putting $\mathfrak{h}_1 = \mathfrak{h}_2 = \mathfrak{h}_5 = 0$ and $\mathfrak{h}_3 = \mathfrak{h}_4 = \mathfrak{h}$ in theorem 3.5, we complete the proof.

Example 3.9

Let $X = [0,1]$ and \mathfrak{A} be the set of all real valued functions on X which also have continuous derivatives on X with the norm $\|\varrho\| = \|\varrho\|_\infty + \|\varrho'\|_\infty$ and the usual multiplication. Let $\mathfrak{B} = \{\varrho \in \mathfrak{A} : \varrho(t) \geq 0, t \in X\}$. It is clear that \mathfrak{B} is a non normal cone and \mathfrak{A} is a Banach algebra with a unit $e = 1$. Define a mapping $D_b : X \times X \rightarrow \mathfrak{A}$ by

$$D_b(\varrho, \varsigma)(t) = |\varrho - \varsigma|^2 e^t$$

we make a conclusion that (X, D_b) is a complete cone b -metric space over Banach algebra \mathfrak{A} with the coefficient $\omega = 2$. Now define the mappings $\mathfrak{S}, \mathfrak{L} : X \rightarrow X$ by

$$\mathfrak{S}(\varrho) = \frac{\varrho}{8}, \quad \mathfrak{L}(\varsigma) = \frac{\varsigma}{2}$$

Taking $\mathfrak{h}_1 = \frac{1}{12} + \frac{1}{12}t$, $\mathfrak{h}_2 = \frac{1}{16} + \frac{1}{16}t$ and $\mathfrak{h}_3 = \mathfrak{h}_4 = 0$, $\mathfrak{h}_5 = \frac{1}{8} + \frac{1}{8}t$. Show that all conditions of Theorem 3.5 are satisfied. Theorem, 0 is the unique common fixed point of \mathfrak{S} and \mathfrak{L} .

Example 3.10

Let $X = [0,1]$. Define a function $D_b : X \times X \rightarrow \mathfrak{A}$ by $D_b(\varrho, \varsigma) = |\varrho - \varsigma|$. Clearly, (X, D_b) is a complete cone b -metric space over Banach algebra \mathfrak{A} with coefficient $\omega = 2$. Now define $\psi : \mathfrak{B} \rightarrow \mathfrak{B}$ by $\psi(t) = t$ for all $t > 0$. Then $\psi \in \Psi$. Also define $\phi : \mathfrak{B} \rightarrow \mathfrak{B}$ by $\phi(t) = \mathcal{K}t$ for all $t > 0$. Then ψ is a continuous comparison function.

Define the mapping $\mathfrak{S}, \mathfrak{L} : X \rightarrow \mathfrak{CB}(X)$ by $\mathfrak{S}(\varrho) = \frac{\varrho}{8}$, $\mathfrak{L}(\varsigma) = \frac{\varsigma}{2}$ for all $\varrho, \varsigma \in X$. Then,

$$\phi(\mathcal{H}(\mathfrak{S}\varrho, \mathfrak{L}\varsigma)) \leq \mathfrak{F}(\phi(\mathcal{M}(\varrho, \varsigma)))$$

where,

$$\mathcal{M}(\varrho, \varsigma) = \mathfrak{h}_1 \frac{D_b(\varrho, \mathfrak{S}\varrho)}{1 + D_b(\varrho, \mathfrak{S}\varrho)} + \mathfrak{h}_2 \frac{D_b(\varsigma, \mathfrak{L}\varsigma)}{1 + D_b(\varsigma, \mathfrak{L}\varsigma)} + \mathfrak{h}_3 \frac{D_b(\varrho, \mathfrak{L}\varsigma)}{1 + D_b(\varrho, \mathfrak{L}\varsigma)} + \mathfrak{h}_4 \frac{D_b(\varsigma, \mathfrak{S}\varrho)}{1 + D_b(\varsigma, \mathfrak{S}\varrho)} + \mathfrak{h}_5 D_b(\varrho, \varsigma)$$

$$\begin{aligned}
\phi(\mathcal{H}(\mathfrak{S}\varrho, \mathfrak{L}\varsigma)) &\leq \psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{|\varrho - \mathfrak{S}\varrho|}{1+|\varrho - \mathfrak{S}\varrho|} + \mathfrak{h}_2 \frac{|\varsigma - \mathfrak{L}\varsigma|}{1+|\varsigma - \mathfrak{L}\varsigma|} + \mathfrak{h}_3 \frac{|\varrho - \mathfrak{L}\varsigma|}{1+|\varrho - \mathfrak{L}\varsigma|} \\ &+ \mathfrak{h}_4 \frac{|\varsigma - \mathfrak{S}\varrho|}{1+|\varsigma - \mathfrak{S}\varrho|} + \mathfrak{h}_5 |\varrho - \varsigma| \end{aligned} \right) \right) \\
&\leq \psi \left(\varphi \left(\begin{aligned} &\mathfrak{h}_1 \frac{|\varrho - \frac{\varrho}{8}|}{1+|\varrho - \frac{\varrho}{8}|} + \mathfrak{h}_2 \frac{|\varsigma - \frac{\varsigma}{2}|}{1+|\varsigma - \frac{\varsigma}{2}|} + \mathfrak{h}_3 \frac{|\varrho - \frac{\varsigma}{2}|}{1+|\varrho - \frac{\varsigma}{2}|} \\ &+ \mathfrak{h}_4 \frac{|\varsigma - \frac{\varrho}{8}|}{1+|\varsigma - \frac{\varrho}{8}|} + \mathfrak{h}_5 |\varrho - \varsigma| \end{aligned} \right) \right) \\
&\leq \psi \left(\phi \left(\frac{1}{6} |\varrho - \varsigma| \right) \right), \\
&\leq \psi \left(\frac{1}{6} |\varrho - \varsigma| \right) \\
&\leq \frac{\mu}{6} |\varrho - \varsigma|, \quad \text{for } 0 < \mu < 1 \\
&\leq \frac{\mu}{6} M(\varrho, \varsigma) = \frac{\mu}{6} \phi(M(\varrho, \varsigma)) \\
&\leq \psi(\phi(M(\varrho, \varsigma))).
\end{aligned}$$

Choose $\mathfrak{h}_1 = \mathfrak{h}_2 = \mathfrak{h}_3 = \mathfrak{h}_4 = 0, \mathfrak{h}_5 = \frac{1}{6}$. Note that \mathfrak{S} and \mathfrak{L} commute at the coincidence point $\varrho = 0$. The pair $(\mathfrak{S}, \mathfrak{L})$ is weakly compatible, it is easy to see that all the conditions of theorem 3.5 holds trivially good and 0 is the unique common fixed point of \mathfrak{S} and \mathfrak{L} .

4 Conclusions

In Theorem 3.5 we have formulated a new contractive conditions to modify and extend some common fixed point theorem (ϕ, \mathfrak{F}) -multi-valued mapping in cone b -metric space over Banach algebra. The existence and uniqueness of the result is presented in this article. We have also given some example which satisfies the condition of our main result. Our result may be the vision for other authors to extend and improve several results in such spaces and applications to other related areas.

5 Acknowledgements

The authors thanks the management, Asian Research Journal of Mathematics for their constant support towards the successful completion of this work. We wish to thank the anonymous reviewers for a careful reading of manuscript and for very useful comments and suggestions.

References

- [1] Alikhani H, Gopal D, Miandaragh MA, Rezapour Sh, Shahzad N, Some endpoint results for generalized weak contractive multifunction. Sci World J 2013:7. Article ID 948472, (2013).
- [2] Aydi H, Monica Felicia Bota, Erdal Karuinar and Slobodanka M, A fixed point theorem for set-valued quasi-contractions in b-metric spaces. Fixed Point Theory Appl 2012:88, (2012) doi:10.1186/1687-1812-2012-88, (2012).

- [3] Banach S, Sur les operations dans les ensembles abstraits et leur application aux equations integrales. *Fundam Math* 3:133–181, (1922).
- [4] Bakhtin, I.A. The contraction mapping principle in almost metric spaces. *Funct. Anal.* 30, 26–37, (1989).
- [5] Boriceanu M, Fixed point theory for multivalued generalized contraction on a set with two b-metrics. *Stud Univ Babes-Bolyai Math LIV(3)*:1–14, (2009).
- [6] Czerwik S, Contraction mappings in b-metric spaces. *Acta Math Inform Univ Ostraviensis* 1:5–11, (1993).
- [7] Czerwik S, Nonlinear set-valued contraction mappings in b-metric spaces. *Atti Semin Math Fis Univ Modena* 46(2):263–276, (1998).
- [8] Dhamodharan D, and Krishnakumar R, Common Fixed Points of (ψ, φ) -Weak Contractions in Regular Cone Metric Spaces, *International Journal of Mathematical Archive-8(8)*, 1-8, (2017).
- [9] Krishnakumar R and Dhamodharan D, Common Fixed Point of Four Mapping With Contractive Modulus on Cone Banach Space, *Malaya J. Mat.*, 5(2), 310–320, (2017).
- [10] Dhamodharan D, and Krishnakumar R, Cone C-class Function with Common Fixed Point Theorems for Cone b-metric Space, *Journal of Mathematics and Informatics*, Vol. 8, 83-94, (2017).
- [11] Khan M. S., Priyobarta M and Yumnam Rohen, Fixed points of generalised rational α^* - ψ -Geraghty contraction for multivalued mappings, *Journal of Advance Study*, 12(2), 156-169, (2019).
- [12] Laishram Shanjit and Yumnam Rohen, Best proximity point theorems in b-metric space satisfying rational contraction, *Journal of Nonlinear Analysis and Applications*, 2019(2), 12-22, (2019).
- [13] Özen Özer, A. Shatarah, An in depth guide to fixed point theorems, 2021, ISBN: 978-1-53619-565-1. NOVA Science Publisher, New York, U.S.A
- [14] Özen Özer , Saleh Omran, On the generalized C*- valued metric spaces related with Banach fixed point theory, *International Journal of Advanced and Applied Sciences*, 4(2), 35-37, (2017).
- [15] Huang L.G and Zhang X, Cone metric space and fixed point theorems of contractive mappings, *J. Math. Anal. Appl.* 332(2) 1468-1476, (2007).
- [16] Jiang, B., Xu, S., Huang, H and Cai, Z. Some fixed point theorems for generalized expansive mappings in cone metric space over Banach algebras. *Journal of Computational Analysis and Applications*, 21(6):1103-1114, (2016).
- [17] Liu H and Xu S, Cone metric spaces with Banach algebras and fixed point theorems of generalized Lipschitz mappings, *Fixed Point Theory Appl.* 2013, 320, (2013).
- [18] Nadler SB, Multi-valued contraction mappings. *Pac J Math* 30 (1969) 475–488.
- [19] Rudin W, *Functional analysis*, McGraw-Hill, New York, (1991).
- [20] Wang, Z., Li, H, Fixed point theorems and endpoint theorems for (α, ψ) - Meir-Keeler – Khan Multivalued mapping, *Fixed Point theory appl*, 2016, 12,(2016).