

Original Research Article

Pattern of Ocular Morbidity in Patients Attending Urban Specialist Eye Clinic in South East Nigeria.

Abstract

Background: The pattern of eye diseases varies across the world. Racial, ethnic, socio-cultural and socio-demographic characteristics are known to exert some influence on the pattern of these eye diseases. Knowledge of these patterns is important in planning for adequate eye care services.

Aim: To determine the pattern of ocular morbidity in an urban specialist eye clinic in Awka, South-East Nigeria.

Setting and Design: A two-year descriptive retrospective study of patients seen in an urban specialist eye clinic in South-East Nigeria

Materials and Methods: The needed information, which included socio-demographics and diagnosis, was extracted from patients' case files and entered into a proforma. The data were analyzed using SPSS version 20. Ninety five percent Confidence Interval was given for percentages and $p \leq .05$ was accepted as statistically significant.

Results: Five hundred and twenty seven patients comprising 277 (52.6%) females and 250 (47.4%) males with a mean age of 43.2 ± 21.7 years seen within the two year period were reviewed. Refractive error, glaucoma, allergic eye diseases, and cataract in that order were the commonest ocular morbidities in these patients. Spherical errors were commoner than cylindrical errors. Hypermetropia was the commonest refractive error while hypermetropic astigmatism was the commonest astigmatic error. All the refractive

errors were commoner in females than the males. Although 185 (35.1%) patients were presbyopic on clinical examination, only 41(7.7%) patients had presbyopia as their main diagnosis.

Conclusion: Most of the common reasons for ophthalmic consultations found were known causes of visual impairment and blindness. Adequate provision for the management of these conditions will help in reducing the burden of visual impairment and blindness.

Key Words: Ocular, Morbidity, Urban, Clinic, Nigeria, Cataract, Refractive error, Glaucoma.

Introduction

There are different patterns of eye diseases across the world. Racial, geographic, social, and cultural factors all have some influence on the pattern of these ocular diseases.^{[1], [2]} Ukponmwan in Benin- city, Nigeria reported the common causes of ocular morbidity in a tertiary hospital to include refractive errors, cataract, allergic conjunctivitis and glaucoma in that order.^[3] Oladigbolu et al^[4], in northern Nigeria, on the contrary reported that infective conjunctivitis, followed by allergic conjunctivitis, was the commonest reason for eye clinic consultation, though this was in a university sick bay. Ademe et al^[5] in Ethiopia, found that the commonest reason for eye clinic consultation was allergic conjunctivitis, followed by refractive error, cataract and glaucoma. Lipa et al^[6], also reported allergic conjunctivitis as the commonest reason for out- patient eye clinic consultations in a tertiary hospital in India. A study in a community centre in Nepal, similar to the finding in Benin-City, Nigeria, reported that refractive error, cataract and conjunctivitis were the most prevalent eye diseases seen in the centre.^[2] Refractive error and conjunctivitis were commoner in the younger age group while cataract and other posterior segment lesions were commoner in the older age group.^[2] A study among the elderly in Southern Germany recorded that cataract, dry eyes, glaucoma and age related macular degeneration were the commonest eye diseases found on examination.^[7] On the other hand, Lusambo et al^[8] in Kinshasa, Democratic Republic of Congo reported allergic conjunctivitis and refractive error as the commonest eye diseases in a population-based study in children aged 16 years and below.

Knowledge of the different patterns of eye diseases/reasons for eye clinic consultations are important in planning for eye care and allocation of scarce resources to the necessary and most needed eye care facilities/equipment especially in resource constrained regions of the world. This study aimed to determine the pattern of eye diseases at an urban specialist general clinic in Awka, South-East, Nigeria, with a view to generating information that may help in planning for adequate eye care provisions in the region and other regions that share similar characteristics.

Materials and Methods

This was a two-year descriptive retrospective study, carried out on patients who presented at a specialist eye clinic in Awka, Nigeria, between January 2020 and December 2021.

Case files of all patients who presented to the clinic within the two year period were retrieved from the medical records and the needed information extracted from the case files and recorded in a proforma designed for the study. Information obtained included socio-demographic data (age, sex, occupation), and ocular diagnosis on the first consultations. Where there were more than one diagnosis, the one that has the worst effect on vision and/or causes greatest discomfort to the patient was taken as the major diagnosis. Primary open angle glaucoma diagnosis was made when the following criteria were met on slit lamp binocular indirect ophthalmoscopy with 90D lens, and automated visual field testing : open anterior chamber angle (Shaffer's grade 2, 3 or 4); glaucomatous optic neuropathy [vertical cup/disc ratio (VCDR) ≥ 0.7 and/or VCDR asymmetry > 0.2], associated with glaucomatous visual field defect. Glaucomatous visual field defect presence was based on the Anderson's criteria.^[9] Ocular hypertension was diagnosed based on intraocular pressure > 21 mmHg in the absence of glaucomatous optic neuropathy and glaucomatous visual field defect^[10]. Shaffer's grade 0 and 1 were diagnosed as angle closure glaucoma. Cataract was defined as any visually significant lens opacity. The data were analyzed using statistical package for social science version 20 (SPSS-20). Ninety five percent Confidence Interval was given for percentages. Associations were tested with t-test and $p \leq .05$ was described as statistically significant. The tenets of Helsinki declaration was adhered to throughout the study.

Results

Five hundred and twenty seven (527) patients comprising 277 (52.6%) females and 250 (47.4%) males (female:male = 1:1.1) presented within the period under review. The age range of the patients was 1- 93 years (mean=43.2 \pm 21.7). Majority (60%) of the patients were aged between 20 and 60 years (table 1).

Table 1: Age Category of Patients

Age Category (Years)	Frequency	Percentage (%)
1-10	40	7.6
11-20	41	7.8
21-30	90	17.1
31-40	64	12.1
41-50	94	17.8
51-60	77	14.6
61-70	64	12.2
71-80	36	6.8
81-90	18	3.4
Above 90	3	0.6
Total	527	100

The commonest ocular morbidity in the study was refractive error which accounted for 27.3% [(144/527); 95% CI= 23.4-31.2] of all new consultations (table 2).

Different types of glaucoma, of which primary open angle glaucoma was the commonest, constituted the second commonest ocular morbidity. Glaucoma constituted 17.4% [(92/527); 95% CI= 14.1-20.7] of the ophthalmic diagnosis. Allergic eye disease, cataract, and presbyopia respectively were the third, fourth and fifth commonest diagnosis (table 2). Although only 41 (7.7%) patients had presbyopia as their main diagnosis (table 2), 185 (35.1%) patients were found to have presbyopia on clinical assessment.

Table 2: Main Diagnosis on First Eye Clinic Consultations

Diagnosis	Frequency	Percentage (%)
Refractive Error	144	27.3
Primary Open Angle Glaucoma	71	13.4
Angle Closure Glaucoma	10	1.9
Secondary Glaucoma	7	1.3
Neovascular Glaucoma	4	0.8
Allergic Eye Disease	65	12.3
Cataract	45	8.5
Presbyopia	41	7.8
Infective Conjunctivitis	14	2.6
Dry Eye Syndrome	13	2.5
Age Related Macular Degeneration	12	2.3
Uveitis	10	1.9
Pterygium	10	1.9
Diabetic Retinopathy	8	1.5
Posterior Vitreous Detachment	7	1.3
Corneal Ulcers	6	1.0
Non glaucomatous Optic Atrophy	6	1.0
Ocular Hypertension	5	0.9
Chorioretinal Scar	4	0.8
Corneoscleral Laceration	4	0.8

Preseptal/Orbital Cellulitis	4	0.8
Phthisis Bulbi	4	0.8
Migraine	4	0.8
Thyroid Eye Disease	4	0.8
Corneal dystrophy	3	0.6
Chalazion	3	0.6
Herpes Zoster Ophthalmicus	3	0.6
Conjunctival Cysts	2	0.4
Conjunctival Naevus	2	0.4
Conjunctival Squamous Cell Carcinoma	2	0.4
Corneal Foreign Body	2	0.4
Corneal Scar/Opacity	2	0.4
Lid Laceration	2	0.4
Accommodative Esotropia	2	0.4
Traumatic Hyphaema	2	0.4
Total	527	100

Spherical error was commoner than astigmatic error among these patients ($p < .000$), with hypermetropia being the commoner spherical error as well as the commonest refractive error (Table 3). Hypermetropic astigmatism was the commonest astigmatic error while mixed astigmatism was the least form of astigmatic error as well as the least common of all error (Table 3).

Table 3: Pattern of Refractive Error

Refractive Error	Frequency	Percentage (%)
Hypermetropia	43	29.9
Myopia	40	27.8
Hypermetropic astigmatism	29	20.0
Myopic astigmatism	24	16.7
Mixed astigmatism	8	5.6
Total	144	100

All the refractive errors were commoner in females ($p < .000$), being present in 32.5% (90/277) of females and constituting 62.5% (90/144) of all errors (table 4); female: male for refractive errors = 1.7:1.

Table 4: Sex of Patients * Refractive Errors Cross-tabulation

	REFRACTIVE ERROR					TOTAL
	Myopia	Hypermetropia	Myopic Astigmatism	Hypermetropic Astigmatism	Mixed Astigmatism	
Male	16	14	10	11	3	54
FEMALE	24	29	14	18	5	90

Total	40	43	24	29	8	144
--------------	-----------	-----------	-----------	-----------	----------	------------

Discussion

Refractive error, glaucoma, allergic eye disease and cataract in that order were the commonest ocular morbidities in this study. These commonest ocular morbidities, excluding allergic eye diseases, are among the commonest causes of visual impairment and blindness worldwide as well as in Nigeria.^{[11],[12]} This is similar to the findings of Ukponmwan^[3] in Benin-City, Nigeria where she reported refractive error as the most prevalent ocular morbidity in a tertiary eye centre. However, contrary to Ukponmwan's report of cataract being the second commonest eye disease, the present study found glaucoma to be the second commonest eye disease. Allergic eye disease was the third commonest in both Ukponmwan's study and the present study. The larger sample size of Ukponmwan's study (7220) compared to that of present study (527) may have accounted for some of the observed differences. Eze et al^[13] similarly found that refractive error was the commonest cause of ocular morbidity among public secondary school students in Enugu, Nigeria . They reported that more than half (57%) of ocular morbidities among these children were due to refractive error. The mean age of Eze et al's subjects were however much lower (14.2 years \pm 1.9) than that of the present study (43.2 \pm 21.7), and all were children and young adults aged 10 – 21 years while the present study included children as young as one year and adults as old as 93 years. Similarly, Razyal et al^[2] in Nepal and Mukwanseke et al^[14] in Kinshasa, Democratic Republic of Congo, reported refractive error as the commonest ocular morbidity. The Kinshasa study^[14] had similar sample size (500) and mean age (41.9 years) to those of the present study. Majority of the subjects in the Nepal study^[2] were also aged between 20- 60 years similar to the present study. These similarities in the sample population could account for the similarities in the findings despite the racial differences. Oladigbolu et al^[4], contrary to the present finding,

reported infective conjunctivitis as the commonest ocular morbidity in a university sick bay in Nigeria. Oladigbolu subjects, however, were younger (mean age=24.3) and majority were students. A study by Chinawa et al^[15] in Rivers state, Nigeria, contrary to the present finding, recorded glaucoma and then refractive error as the leading causes of ocular morbidity. Their study was however a community based study while the present study was a hospital based study. Also in contrast to the present study, Ademe et al^[5] in Ethiopia and Lipa et al^[6] in India both reported allergic conjunctivitis as the commonest diagnosis in eye clinic consultations. Both the Ethiopian and Indian studies however have lower mean ages compared to that of the present study, and this together with geographical differences could account for the observed differences.

Among patients with refractive error, hypermetropia was found to be the commonest error, followed by myopia, with mixed astigmatism being the least common error. This is similar to the finding of Bekibele et al^[16] among drivers in Ibadan, Nigeria. They found hypermetropia to be the commonest refractive error despite the study being community based and their subjects being all males. Ezepue^[17] in Enugu and Bagaiya et al^[18] in Kaduna similarly reported hypermetropia as the commonest refractive error. Ezepue and Bagaiya et al's studies, similar to the present study, were hospital based studies and included both adults and children. A study among secondary school teachers in Onitsha, Nigeria^[19] similarly recorded hypermetropia as the commonest refractive error although it was a community based study involving only adults. The present finding on the other hand is different from the findings of Ezelum et al^[20] in Nigeria and Anajekwu et al^[21] in Nsukka, Nigeria. Both Ezelum et al and Anajekwu et al's studies were community based and involved only adults unlike the present study. These differences could account for the observed variations in the pattern of refractive error.

There was a statistically significant difference between the sexes with all the refractive errors being commoner in females than the male. This is similar to what Lawan et al^[22] found in Kano, Nigeria. Lawan's subjects, though eye clinic patients like the present study, were all adults aged 35 years and above. Besufikad et al^[23] in Ethiopia, similarly found that refractive error is more prevalent in women in a

retrospective hospital based study. On the contrary, Malu et al^[24] in Jos, Nigeria and Abah et al^[25] in Zaria, Nigeria, did not find any statistically significant difference in the prevalence of refractive errors between male and female. This is surprising as both are hospital based studies and included adults and children just like the present study. Abah's subjects were however younger with a mean age of 24.6 ± 4.9 years. Anajekwu et al^[21] also did not find any statistical difference in the refractive errors between male and female. Anajekwu et al's subjects, however, were only adults and the study was a community based study unlike the present study. These could account for the observed differences.

Conclusion

Refractive error was the commonest reason for ophthalmic consultation in this study. Glaucoma, allergic eye disease and cataract were the second, third and fourth commonest reasons for ophthalmic consultations respectively. These common eye morbidities, with the exclusion of allergic eye disease, are known and common causes of visual impairment and blindness globally. There was also a high incidence of presbyopia that needed correction among these subjects. Adequate provisions in terms of facility, drugs and manpower, should be made available by relevant eye health planning bodies and hospitals to ensure proper management of these common eye morbidities. This will go a long way to help in reducing the burden of visual impairment and blindness, as well as the discomfort associated with these eye conditions.

References

1. Mahmoud AO, Olatunji FO, Buari SB. Survey of blindness and ocular morbidities in Kwara State, Nigeria. *Nig J Surg Sci.* 2005; 15: 26-30.
2. Rizyal A, Shakya S, Shresta RK, Shresta S. A study of ocular morbidity of patients attending a satellite clinic in Bhaktapur, Nepal. *Nepal Med Coll J.* 2010; 12(2): 87-89.
3. Ukponmwan CU. Pattern of ocular morbidity in Nigeria. *Asian Pac J Trop Dis.* 2013; 3(2):164-6.

4. Oladigbolu KK, Abah ER, Chinda D, Anyebe EE. Pattern of eye diseases in a university health service clinic in northern Nigeria. *Niger J Med.* 2012;21(3):334-7.
5. Ademe S, Edmealem A. Pattern of ocular diseases among patients attending ophthalmic outpatient department: A cross-sectional study. *Int J Clin Exp Ophthalmol.* 2020;4:049-053.
6. Lipa M, Banu TJ. A cross-sectional study to assess the morbidity pattern of ocular diseases in outpatient department of ophthalmology at a tertiary care hospital. *Int J Res Med Sci.* 2016;4(9):3797-3800.
7. Reitmeir P, Linkohr B, Heier M, Molnos S, Strobl R, Schulz H, et al. Common eye diseases in older adults of southern Germany: results from the KORA-Age study. *Age and Ageing.* 2017;46:481–6
8. Lusambo, NN, Ngoy,JK, Ahuka, AD, Bazitama, YM, Kapopo, AM, Muleya,JI, Mankiew, EM, Stahnke, T, Guthoff, R. Prevalence and patterns of childhood ocular morbidity in Kinshasa: A population-based study. *Global Epidemiology* 3. 2021; 10054.
9. Nayak BK, Dharwadkar S. Interpretation of autoperimetry. *J Clin Ophthalmol Res.* 2014;2:31-59.
10. Wolfs RCW, Borger PH, Ramrattan RS, Klaver CCW, Hulsman CAA, Hofman A. Changing Views on Open-Angle Glaucoma: Definitions and Prevalences—The Rotterdam Study. *Invest. Ophthalmol. Vis. Sci.* 2000;41(11):3309-21.
11. WHO media centre fact sheet. Blindness and vision impairment. 2021, Oct. Available from: <http://www.who.int/news-room/fact-sheet/detail/blindness-and-visual-impairment>. [Last accessed on 2022 Sep 1].
12. Abdull MM, Sivasubramaniam S, Murthy GV, Gilbert C, Abubakar T, Ezelum C, et al; Nigeria National Blindness and Visual Impairment Study Group. Causes of blindness and visual impairment in Nigeria: the Nigeria national blindness and visual impairment survey. *Invest Ophthalmol Vis Sci.* 2009 Sep;50(9):4114-20.
13. Eze GC, Anajekwu CC, Obeagu EI. Prevalence and Causes of Ocular Morbidities among School Children in Enugu-East Local Government Area, Enugu State, Nigeria. *Asian Journal of Research and Reports in Ophthalmology* 2022;5(2): 13-17

14. Mukwanseke E, Kilangalanga J, Lutete F, Hopkins A, Guthoff RF, Frech S. Ocular morbidity—A Critical analysis to improve outpatient services in an eye department in a Sub-Saharan megacity. *J Clin Med.* 2021;10:3791.
15. Chinawa NE, Odogu VK, Ezeh EI, Anyiam FE. Ocular morbidity pattern and presentation among residence of a semi-urban community in Rivers state, Nigeria. *AJMAH.* 2020;18(8):14-20.
16. Bekibele CO, Fawole OI, Bangboye AE, Adekunle LV, Ajayi R, Baiyeroju AM. Prevalence of refractive error and attitude to spectacle use among drivers of public institutions in Ibadan, Nigeria. *Ann Afr Med.* 2007;6:26-30.
17. Ezepue UF. Distribution of presbyopia in the eye clinic of the eastern Nigeria medical centre Enugu. *Orient J Med.* 2003;15:32-5.
18. Bagaiya T, Pam V. Refractive errors in Kaduna, Nigeria. *Niger J Surg Res.* 2003;5:106-9.
19. Anajekwu CC, Nwosu SN. Prevalence and causes of visual impairment among teachers in Onitsha, Nigeria. *Niger J Ophthalmol.* 2017;25:129-32.
20. Ezelum C, Razavi H, Sivasubramaniam S, Gilbert CE, Gudlavalleti VS, Entekume G, et al. Refractive error in Nigerian adults: Prevalence, type, and spectacle coverage. *Invest Ophthalmol Vis Sci.* 2011;52:5449-56.
21. Anajekwu CC, Kizor-Akaraiwe N. Uncorrected refractive error in a university community. *Niger J Clin Pract.* 2022;25:361-7.
22. Lawan A, Okpo E, Philips E. Refractive errors in presbyopic patients in Kano, Nigeria. *Ann Afr Med.* 2014;13:21-4.
23. Besufikad B, Hailemichael W, Tilahun L, Yiman W, Anteneh S. Refractive errors and associated factors among patients visiting BoruMeda Hospital's secondary eye Unit in Dessie Town, South Wollo Zone, Ethiopia. *BMC Ophthalmol.* 2022;22:312

24. Malu K, Ojabo C. Refractive error in patients attending a private hospital in Jos, Nigeria. Niger J Clin Pract. 2014;17:106-11.

25. Abah ER, Chinda D, Samaila E, Anyabe EE. Profile of refractive error and presbyopia in a university community: A clinical study. Ann Nigerian Med. 2010;4:55-8.

UNDER PEER REVIEW