

Original Research Article

Pre-harvest salicylic acid spray on shelf life and biochemical changes of litchi during storage

ABSTRACT

An experiment was conducted on litchi cv. China at Horticulture garden in the Department of Horticulture (Fruit and Fruit Technology), Bihar Agricultural University, Sabour, Bhagalpur, Bihar, to show the shelf life and biochemical changes of litchi during storage. The experiment was laid out in randomized block design with five treatments and four replications. Salicylic acid was sprayed twice i.e. 15 days and 30 days before anticipated harvesting time with different concentrations of salicylic acid (T₁- 50 ppm, T₂-100 ppm, T₃-150 ppm, T₄-200 ppm and T₅-control). Different concentrations have different effect on shelf life and biochemical parameters. Salicylic acid with T₃-150 ppm has increased TSS (20.02 °Brix), total sugar (13.10 %), ascorbic acid (38.21 mg/100 g pulp), anthocyanin (8.06 mg/100g pulp) and shelf life (4 days) on the 6th day of storage at ambient condition.

Keywords: Litchi, salicylic acid, biochemical parameters, shelf life

1. INTRODUCTION

As India has such a diverse climate, it is ideal for cultivation of different varieties of litchi. Litchi (*Litchi chinensis* Sonn.) belongs to sapindaceae family. It is native of Southern China. Its chromosome no. 2n=30 and type of fruit is one seeded nut and its edible part is fleshy aril. It is highly cross pollinated crop. Its name is derived from Chinese word “lee chee” which means “one who gives the pleasure of life”. It is grown in tropical and sub tropical regions. Due to its bright red peel colour and pleasant flavoured juicy aril this fruit is highly valued in national and international markets. Litchi fruit is very much liked as a table fruit as well as in dried and canned forms. Jam, jelly, squash and cordial are also prepared from this fruit. It is very nutritious and good source of vitamin C (64 mg/100 g pulp). It also contains vitamin E, B complex and trace amount of protein (0.7%), fat (0.3%) and minerals like phosphorous, calcium and iron. It is non climacteric type fruit; it does not ripe after harvesting from plants so it is harvested at ripe stage. Litchi is grown in different countries like China, West Indies, Myanmar, Japan, South Africa, Florida, Hawaii, etc. in the world; India is one among them. China is first and India is second largest producer of litchi fruits in the world. Litchi is very perishable in nature and it has very short shelf life. Litchi being non climacteric in nature it is harvested at ripe stage. Generally, the harvest maturity of the fruit is judged by the development of red colour on epicarp and flattened of the tubercles. After harvesting, the fruit turn brown within a couple of days due to loss of water from the pericarp. (Panwar *et al.* 2018) reported that salicylic acid decreases

respiration rate by the closure of stomata. Salicylic acid (SA) is a natural plant hormone and it acts as signaling molecule. It makes the plants resistance against biotic and abiotic stress (Khan *et al.* 2012). SA helps in plant growth, ion uptake and nutrient transport in plants. It is phenolic compound, and in plants it provides systemic resistance to fungal pathogens (Meena *et al.* 2014). Pre harvest application of various chemicals have been reported to enhance the shelf life of fruits by reducing physiological loss in weight and decay losses during storage (Gupta and Metha, 1988).

2. MATERIALS AND METHODS

The present investigation was carried out at the Horticulture Garden under the Department of Horticulture (Fruit & Fruit Technology), Bihar Agricultural College, Bihar Agricultural University, Sabour, Bhagalpur during the year of 2021-2022 with a view to study the pre-harvest application of salicylic acid on shelf life and biochemical changes in litchi during storage. Variety used for this experiment was “China”. The experiment was laid out with five treatments (T₁-50 ppm SA, T₂- 100 ppm SA, T₃- 150 ppm SA, T₄- 200 ppm SA and T₅- control) and four replications in randomized block design. Bihar Agricultural College Sabour is situated 25°50' N latitude and 87°19' E longitude at an altitude of 52.73 m above mean sea level. It has an average annual rainfall of 1348 mm precipitating mostly in between middle of June to middle of October.

Biochemical parameters namely titrable acidity (%) was calculated by the given titration method (AOAC, 2000), ascorbic acidity (mg/100 g pulp) was estimated by using 2-6-dichlorophenol indophenols dye method (Jones and Hughes, 1983), TSS (°Brix) was recorded with the help of digital hand refractometer, TSS: Acid ratio was calculated by dividing TSS with titrable acidity, total sugar (%) was estimate by Lane and Eyon (1923) copper titration method, anthocyanin (mg/100 g pulp) was calculated by peel pH- differential method (Wrolstad *et al.* 2005) using two different buffer systems: potassium chloride buffer (0.025 M, pH 1.0) and sodium acetate buffer (0.4 M, pH 4.5), antioxidant (μ mol Trolox Eq. /g pulp) was determined by cupric reducing antioxidant capacity (CUPRAC) method (Apak *et al.* 2008).

Statistical analysis

The experiment was laid out in RBD. Statistical analysis was performed in two way factorial RBD method. Least significant difference was calculated following significant F-test ($p \leq 0.05$). Effect of different treatments on various parameters and their interactions were assessed with ANOVA. Standard errors were computed by MS-Excel. This RBD design was adopted as suggested by Panse and Sukhatme (1967).

3. RESULT AND DISCUSSION

Total soluble solids (°Brix)

Maximum (20.99 °Brix) TSS was in T₃-150 ppm while minimum (10.20) in T₅- control. It was found that untreated (control) fruits exhibited a rapid increase in total soluble solids (TSS) from the day of harvest and a sharp decline, thereafter during storage at ambient conditions compared to the treated ones. The rapid and higher TSS in control fruits might be due to faster ripening associated with the hydrolysis of starch into simple sugars. The delayed increase of TSS in SA treated fruits might be due to inhibition of ethylene biosynthesis. A delayed increase in TSS of SA treated fruits was reported in kiwifruit (Kazemi *et al.* 2011), peach (Khadami and Ershadi, 2013) and persimmon (Khademi *et al.* 2012).

Total sugar (%)

Total sugar increased with increase in storage period. Among all the treatments fruits treated with T₃-150 ppm salicylic acid recorded highest (18.90 %) sugar as compared to other treatments, while lowest (9.30 %) in control. The increase in total sugars of fruits with SA may be due to loss of water from the fruits and conversion of polysaccharides and pectic substances in sugars. The higher rate of acceleration in SA treated fruits may be due to retardation in the rate of normal changes of polysaccharides to total sugars because of its low rate of respiration and oxidation in treated fruits reported by Roe and Brummer (1981) and Yuvraj *et al.* (1999) in mango fruits.

Titration acidity (%)

The highest (0.569 %) titration acidity was observed in treatment T₅-control whereas lowest (0.255 %) in T₃-150 ppm. A continual decrease in titration acidity was noticed during storage at ambient conditions. SA treated fruits shown a significant slower decline over the untreated fruits. The reduction in acidity during storage after attainment of maturity and ripening may be due to utilization of organic acids as a substrate (Islam *et al.* 2013). Similar results of greater titration acidity in mango fruits were also reported by Barman (2013).

Ascorbic acid (mg/100g pulp)

Highest (10.10 mg/100g pulp) ascorbic acid was in treatment T₃-150 ppm salicylic acid whereas lowest in T₅-control. Ascorbic acid was decreased with advancement of storage period. It was revealed that decrease in ascorbic acid was significantly higher throughout the storage period in the fruits treated with SA compared to untreated ones. It could be possible due to retardation of oxidation process and slow rate of conversion of L-ascorbic acid into dehydroascorbic acid (DHA) by ascorbic acid oxidase. DHA concentration was maintained at higher levels in salicylic acid treated fruits than in controls, leading to higher concentration in ascorbic acid throughout the storage period (Garcia-Pastor *et al.* 2020).

Anthocyanin (µmol Trolox Eq. /g pulp)

Highest (8.06 µmol Trolox Eq. /g pulp) anthocyanin was found in treatment T₄-200 ppm salicylic acid treated fruits whereas lowest (1.50 µmol Trolox Eq. /g pulp) was found in untreated fruits. SA maintained fruit colour by retarding the degradation of chlorophyll that declined skin colour

changes throughout the storage and retarding senescence rate in table grapes (Champa *et al.* 2015). Similar results were recorded by Ullah and Jawandha (2013), SA showed a slow rate of conversion from green to yellow and chlorophyll degradation in peach fruits.

PLW (Physiological loss in weight) (%)

The lowest (13.97 %) physiological loss in weight was observed in treatment T₄-200 ppm while highest (19.21 %) was in T₅-control. The higher PLW in untreated fruits might be due to active metabolism such as respiration and transpiration, which might have led to greater loss of water during storage in 'Amrapali' mango fruits (Singh and Tiwari, 1994). SA is an electron donor produces free radicals which prevent normal respiration thus leading to lower weight loss (Shafiee *et al.* 2010). The lower PLW per cent in SA treated fruits is due to reduced respiration, transpiration and ethylene production (Srivastava and Dwivedi, 2000).

Spoilage (%)

Fruits treated with different concentrations of SA significantly lowered decay incidence than control. Highest decay (48.02 %) was in T₅-control whereas lowest (17.20%) in T₃-150 ppm SA treated fruits. Salicylic acid helped in enhancement of expression of genes that control the production of phenolic compounds by activating the phenylalanine ammonia-lyase enzyme, increases resistance to infection (Eraslan *et al.* 2007). Dipping of pear fruit in the SA solution, effectively controlled fruit decay during 5 months of cold storage (Asghari *et al.* 2007). SA application had a positive effect in reducing berry decay (Samra, 2015).

Shelf life (Days)

Different concentrations of SA have different effect on shelf life of litchi. Maximum (4 days) shelf life of litchi was observed in T₃-150 ppm concentration over control. This enhancement effect on shelf life period might be due to the role of SA reducing physiological weight loss of fruits, the activity of oxidative enzymes and retarding fruit softening. Also, SA increased phenolic compounds and enhanced resistant against pest and diseases. These results are same as those of Perez-Vicente *et al.* (2002) who reported that, exogenous polyamines applications delayed color changes, reduced mechanical damage, chilling injury susceptibility and increase shelf life in both climacteric and non-climacteric fruits. The maximum shelf life in SA treated fruits might be due to its ability to serve as a physical barrier around the fruit which reduce transpiration and respiration activity along with delay in ethylene production. The positive effect of SA on storage life could probably be due to the modifying the atmosphere. The modified atmosphere created could, therefore, delay the ripening by delaying ethylene production and by reducing the level of internal oxygen and consequently prolonging the storage life of fruit (Gol and Rao, 2011).

4. CONCLUSION

From the above findings it may be concluded that pre-harvest spray of salicylic acid on litchi plant can enhance the shelf life and biochemical properties of fruits. Spray of salicylic acid T₃-150 ppm concentration was found superior in biochemical parameters and shelf life during storage at ambient condition. Biochemical parameters such as total soluble solids, total sugar, titrable acidity, ascorbic acid, anthocyanin and shelf life, physiological loss in weight (PLW) and spoilage.

REFERENCES

1. Asghari M, Shirzad H, Hajitagilo R. Postharvest treatment of salicylic acid effectively controls pear fruit diseases and disorders during cold storage. Novel approaches for the control of postharvest diseases and disorders. Proceedings of the international congress, bologna, Italy, Criofo, University of Bologna. 2007.
2. Barman K. Biologically safe approaches to control sap burn, chilling injury and postharvest diseases of mango. Ph. D. thesis, IARI, New Delhi, India. 2013.
3. Champa WA, Gill MI, Mahajan BV, Arora NK. Preharvest salicylic acid treatments to improve quality and postharvest life of stable (*Vitis vinifera* L.) cv. Flame seedless. *Journal of Food Science and Technol.* 2015; 52(6):3607-3616.
4. Eraslan F, Inal A, Gunes A, Alpaslan M. Impact of exogenous salicylic acid on the growth, antioxidant activity and physiology of carrot plants subjected to combined salinity and boron toxicity. *Science Hort.* 2007; 113:120-128.
5. Garcia-Pastor ME, Zapta PJ, Castillo S, Martinez-Romero D, Guillen F, Valero D. The effect of salicylic acid and its derivatives on increasing pomegranate fruit quality and bioactive compounds at harvest and during storage. *Frontiers in Plant Science.* 2020; 11:668.
6. Gupta OP, Metha N. Effect of pre harvest applications on the shelf life of ber (*Zizyphus mauritiana* Lamk) fruits cv. Gola . *Haryana .J. Hort. Sci.*1988; 17:183-189.
7. Hazarika TK, Marak T. Salicylic acid and oxalic acid in enhancing the quality and extending the shelf life of grape cv. Thompson seedless. *Food science and Tech International.* 2020; 1-13.
8. Islam Md K, Khan MH, Sarkar MAR, Absar N, Sarkar SK. Changes in acidity, TSS and sugar content at different storage periods of the postharvest mango (*Magnifera indica* L.) influenced by bavistin DF. *International Journal Food Science.*2013; 8:1-8.

9. Kazemi M, Aran M, Zamani S. Effect of calcium chloride and salicylic acid treatment and quality characteristics of kiwifruit (*Actinida deliciosa* cv. Hayward) during storage. *American Journal of Plant Physiology*. 2011; 6:183-189.
10. Khadami Z, Ershadi A. Postharvest application of salicylic acid improves storability of peach (*Prunus persica* cv. Elberta) fruits. *International Journal Agriculture Crop Science*. 2013; 5-6:651-655.
11. Khademi O, Zamini Z, Mostofi Y, Kalantari S, Ahmad A. Extending storability of persimmon fruit cv. Karaj by postharvest application of salicylic acid. *Journal Agriculture Science Tech*. 2012; 14:1067-1074.
12. Khan N, Nazar R, Iqbal N, Anjum NA. Phytohormones and Abiotic Stress Tolerance in Plants. *Springer*, Berlin, Heidelberg. 2012.
13. Meena D, Tiwari R, Singh OP. Effect of nutrient spray on growth, fruit yield and quality of aonla. *Annals Plant and Soil Research*. 2014; 16: 242-245.
14. Panse VG, Sukhatme PV. Statistical methods for agricultural workers, Indian council of Agricultural Research, New Delhi. 1967.
15. Panwar N, Rai PN, Kumar J, Shankar D, Singh MDP. Effect of different chemicals on litchi (*Litchi chinensis* Sonn.) cv. rose scented. *Journal of Pharmacognosy and Phytochemistry*. 2018; 7(4):1418-1422.
16. Roe B, Bruemmer JH. Changes in pectic substances and enzymes during ripening and storage of "Keitt mangoes". *Journal of Food Science*. 1981; 46(1):186-189.
17. Samara BN. Impact of postharvest salicylic acid and jasmonic acid treatments on quality of Crimson Seedless grapes during cold storage and shelf life. *International Journal of Advance Research*. 2015; 3: 483:490.
18. Shafiee M, TS Taghvae, Babler M. Application of SA to nutrient solution combined with postharvest treatments (hot water, SA and Ca dipping) improved postharvest fruit quality of strawberry. *Science Horticulture*. 2010; 124:40-45.
19. Singh J, Tiwari JP. Effect of ethephon on the post harvest quality of guava (*Psidium guajava* L.) cv. Sardar. *Progressive Horticulture*. 1994; 26:189-193.
20. Srivastava KM, Dwivedi UN. Delayed ripening of banana fruits by salicylic acid. *Plant Science*. 2000; 158:87-96.

21. Bassiony SS, Maha H, El-Aziz A, Fahmy HM. Effect of foliar application of putrescine and salicylic acid on yield fruit quality and storability of flame seedless grapes (*Vitis vinifera* L). *Journal of plant production*. 2018; 9(12): 1203-1214.
22. Ullah S, Jawandha SK. Effect of post harvest treatments of polyamines on colour of stored peach fruits. *The Asian Journal of Horticulture*. 2013; 8(2):784-787.
23. Yuvraj KM, Ughreja PP, Jambukia TK. Effect of post harvest treatments on ripening changes and storage life of mango fruits. In: National seminar on food processing India, G.A.U. Anand. 1999; 125-129.

Table 1. Effect of pre harvest spray of salicylic acid on TSS (°Brix) during storage at ambient condition

Treatment	Days of observation				Mean
	Day 0	Day 2	Day 4	Day 6	
T ₁ -SA 50 ppm	19.21	19.31	19.88	18.61	19.25
T ₂ -SA100 ppm	19.56	19.68	20.16	20.00	19.85
T ₃ -150 ppm	20.02	20.16	20.78	20.99	20.49
T ₄ -200 ppm	19.85	19.98	20.69	20.64	20.29
T ₅ -Control	18.75	18.86	16.11	10.20	15.98
Mean	19.49	19.60	19.52	18.09	19.17
CD(p≤0.05)	T				0.50
	D				0.45
	T*D				0.99
CV (%)					7.35

Table 2. Effect of pre harvest spray of salicylic acid on Total sugar (%) during storage at ambient condition

Treatment	Days of observation				Mean
	Day 0	Day 2	Day 4	Day 6	
T ₁ -SA 50 ppm	11.66	12.92	13.59	17.98	14.03
T ₂ -SA100 ppm	12.75	13.36	14.72	18.06	14.72
T ₃ - SA150 ppm	13.10	14.46	16.01	18.90	15.62
T ₄ - SA 200 ppm	12.86	13.85	15.26	18.10	15.02
T ₅ -Control	10.84	11.91	10.22	9.30	10.57
Mean	12.24	13.30	13.96	16.47	13.99
CD(p≤0.05)	T				0.26
	D				0.24

	T*D				0.53
CV (%)					5.34

Table 3. Effect of pre harvest spray of salicylic acid on Titrable acidity (%) during storage at ambient condition

Treatment	Days of observation				Mean
	Day 0	Day 2	Day 4	Day 6	
T ₁ -SA 50 ppm	0.605	0.533	0.449	0.317	0.476
T ₂ -SA100 ppm	0.580	0.512	0.435	0.310	0.459
T ₃ -150 ppm	0.440	0.428	0.324	0.255	0.362
T ₄ -200 ppm	0.520	0.493	0.356	0.312	0.420
T ₅ -Control	0.760	0.682	0.568	0.569	0.645
Mean	0.581	0.530	0.427	0.352	0.472
CD (p≤0.05)	T				0.011
	D				0.010
	T*D				0.022
CV (%)					6.70

Table 4. Effect of pre harvest spray of salicylic acid on Ascorbic Acid (mg/100 g) during storage at ambient condition

Treatment	Days of observation				Mean
	Day 0	Day 2	Day 4	Day 6	
T ₁ -SA 50 ppm	34.86	27.26	22.51	8.44	23.27
T ₂ -SA100 ppm	35.44	28.55	22.91	8.73	23.91
T ₃ -150 ppm	38.21	31.90	24.52	10.10	26.18
T ₄ -200 ppm	36.15	30.16	23.30	8.81	24.61
T ₅ -Control	33.01	26.00	19.00	0.00	19.50
Mean	35.54	28.77	22.45	7.22	23.49
CD (p≤0.05)	T				0.50
	D				0.45
	TxD				1.00
CV (%)					6.07

Table 5. Effect of pre harvest spray of salicylic acid on Anthocyanin of peel (mg/100 g pulp) during storage at ambient condition

Treatment	Days of observation				Mean
	Day 0	Day 2	Day 4	Day 6	
T ₁ -SA 50 ppm	19.54	18.74	14.61	6.76	14.91
T ₂ -SA100 ppm	19.96	18.93	14.70	7.66	15.31
T ₃ - SA150 ppm	20.88	19.66	15.11	8.06	15.92
T ₄ - SA 200 ppm	21.54	20.33	16.07	7.75	16.42
T ₅ -Control	18.72	16.58	6.07	1.50	10.72
Mean	20.12	18.85	13.31	6.35	14.66
CD (p≤0.05)	T				0.30
	D				0.26
	TxD				0.59
CV (%)					5.70

Table 6. Effect of pre harvest spray of salicylic acid on PLW (%) during storage at ambient condition

Treatment	Days of observation				Mean
	Day 0	Day 2	Day 4	Day 6	
T ₁ -SA 50 ppm	-	5.35 (21.41)	10.70 (42.81)	15.52 (62.08)	10.53
T ₂ -SA100 ppm	-	4.99 (19.95)	10.35 (41.40)	14.07 (56.26)	9.80
T ₃ - SA150 ppm	-	4.42 (17.67)	9.80 (39.18)	14.29 (57.16)	9.50
T ₄ - SA 200 ppm	-	4.10 (16.40)	9.64 (38.57)	13.97 (55.86)	9.24
T ₅ -Control	-	7.92 (31.67)	13.85 (55.40)	19.21 (76.84)	13.66
Mean	-	5.36	10.87	15.41	10.54
CD(p≤0.05)	T				0.26
	D				0.24
	TxD				0.53
CV (%)					7.06

Table 7. Effect of pre harvest spray of salicylic acid on Spoilage (%) during storage at ambient condition

Treatment	Days of observation	
-----------	---------------------	--

	Day 0	Day 2	Day 4	Day 6	Mean
T ₁ -SA 50 ppm	-	-	18.43 (73.72)	29.16 (116.64)	23.80
T ₂ -SA100 ppm	-	-	17.20 (68.81)	24.72 (98.90)	20.96
T ₃ - SA150 ppm	-	-	14.47 (57.90)	17.20 (68.81)	15.84
T ₄ - SA 200 ppm	-	-	15.17 (60.68)	21.76 (87.04)	18.46
T ₅ -Control	-	-	45.72 (182.87)	48.02 (192.07)	46.87
Mean	-	-	22.20	28.17	25.19
CD(p ≤0.05)	T				0.44
	D				0.28
	TxD				0.62
CV(%)					3.49

Table 8. Effect of pre harvest spray of salicylic acid on shelf life during storage at ambient condition

Treatment	Shelf life enhanced	Shelf life extended over control
T ₁ -SA 50 ppm	4	2
T ₂ -SA100 ppm	4	2
T ₃ - SA150 ppm	6	4
T ₄ - SA 200 ppm	4	2
T ₅ -Control	2	-