

Original Research Article

Effect of Conjunctive use of Fe, Zn and Bio-fertilizers on Growth, Nodulation and Chlorophyll Content of Urdbean (*Vigna mungo* L.)

ABSTRACT

An experiment was conducted on urdbean during *kharif* 2021 at Instructional Farm, College of Agriculture, Ummedganj, Kota (Rajasthan). The experiment comprised 10 treatments viz. (Control, 75% RDF, 100% RDF, 75% RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation, 75% RDF+ *PSB* @ 600 g ha⁻¹ seed inoculation, 75% RDF + *Rhizobium* @ 600g ha⁻¹ + *PSB* @ 600 g ha⁻¹, 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage, 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹, 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹, 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600

Key words: Bio-fertilizer, chlorophyll, Ferrous, nodules, urdbean and zinc.

INTRODUCTION

Pulses are the integral part of Indian dietary system due to richest source of protein and other nutrients. Indian population is predominantly vegetarian and protein requirement for the growth and development of the human being is mostly

g ha⁻¹) was carried out in randomized block design with three replications. Significantly higher plant height (25.28 and 34.43 cm) at 60 DAS and at harvest, nodules plant⁻¹ (29.22) at 45 DAS, nodule fresh weight (86.31 mg plant⁻¹) at 45 DAS, chlorophyll content (3.19 mg g⁻¹) at 45 DAS and dry matter accumulation (2.71, 5.94 and 9.99 g plant⁻¹) at 30, 60 DAS and at harvest were recorded with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ in urdbean over rest of treatments. Among these treatments, the application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ produced maximum seed yield (1269.30 kg ha⁻¹) of urdbean.

met with pulses. The availability of pulses per capita per day has proportionately declined from 71 g to 52 g against the minimum requirement of 70 g day⁻¹/capita [1]. Pulses are improving soil fertility by atmospheric nitrogen fixation through *Rhizobium culture*. In India pulses are

Comment [PS2]: Kindly mention the significant level (P<0.05)

Comment [PS1]: What is RDF; mention in the Abstract. Certainly, I know, it is Recommended dose of fertilizer but for better clarity to reader it will be better to mention it. Simultaneously also mention the dose and source such as urea, SSP, DAP or something.

grown area of 28.34 million ha with total production of 23.15 million tonnes and productivity of 817 kg ha⁻¹[2]. Among the pulses, urdbean [*Vigna mungo* (L.) Hepper], is an important crop in India. Urdbean is the third most important pulse crop after chickpea and pigeonpea in India. It is highly nutritious containing 24-26% protein, 1.3% fat and 60% carbohydrates on dry weight basis and it is rich source of calcium, iron and vitamins[3].

The major constraints for low yield of urdbean are lack of micronutrient application and non-adoption of proper agronomic practices. Micronutrients are equally important in plant nutrition as major nutrients. The incidence of micronutrient deficiencies in crops has increased markedly in recent years due to intensive cropping, loss of top soil by erosion, losses of micronutrients through leaching, liming of acid soils, decreased proportions of farmyard manure or other organic sources compared to chemical fertilizers and use of marginal lands for crop production [4]. The optimum supply of micronutrients under balanced condition is very important for achieving higher productivity of urdbean. Foliar application of micronutrients has been proved to be an important asset in fertilizer application with a specific aim of increasing nutrient availability at the time of need, especially at the later stage of plant growth. Foliar application of nutrients using water soluble fertilizers is one of the possible ways to

enhance the productivity of pulses. The foliar fertilization of Zn and Fe at flowering stage improves growth and yield of urdbean[5].

Bio-fertilizer plays a vital role in maintaining long term soil fertility and sustainability [6]. Incorporation of FYM alone or bio-fertilizers improves available nutrient status of the soil with enhanced soil biological activity which in turn provides a congenial physical condition and improved availability of nutrient in the rhizosphere, resulting in an improvement in the crop growth and providing a better source-sink relationship [7]. Among bio-fertilizers, *Rhizobium* inoculation is cheapest, easiest and safest method of supplying nitrogen to legumes through well-known symbiotic nitrogen fixation process. Inoculation of appropriate strain enhances nodule formation resulting better nitrogen fixation. *Rhizobium* species in association with plant roots in urdbean improved soil fertility by fixing atmospheric nitrogen and produces plant growth substances in the soil. *Rhizobium* inoculation can increase the grain yield of pulse crops [8].

MATERIALS AND METHODS

A field experiment was conducted on urdbean during *kharif* 2021 at Instructional Farm, College of Agriculture, Ummedganj, Kota (Rajasthan). Geographically, is situated at 25.11^o North latitude and 75.50^o East

Comment [PS3]: Productivity of the Urdbean should be provided for better clarity

Comment [PS4]: Should be written Farm yard manure (FYM) first time and then subsequently can be used in the whole manuscript.

Comment [PS5]: The cited reference does not provide any relevant data to written statement that the biofertilizers improve the available nutrient status of the soil with enhanced biological activity. Herewith, I suggest authors, to add one more reference along with [7] to proper supporting the statement.

For example: the author can add the following reference
Garima, Bhardwaj, D. R., Thakur, C. L., Kaushal, R., Sharma, P., Kumar, D., & Kumari, Y. (2021). Bamboo-based agroforestry system effects on soil fertility: Ginger performance in the bamboo subcanopy in the Himalayas (India). *Agronomy Journal*, 113(3), 2832-2845.
As, it properly support the written statement on both parameters, yield and soil properties on ginger crop but that does not matters.

Comment [PS6]: One more thing, the authors not only considering the individual treatment but also the conjunctive application of the inorganic fertilizers and biofertilizers. Thus, the statement should be added like
Previously researchers have indicated that the conjunctive application of the inorganic, organic and biofertilizers enhanced the growth of the agricultural crops (Meena et al., 2022).
Meena, N. K., Latore, A., Singh, M., Sharma, P., & Verma, K. (2022). Conjunctive Application of Organic and Inorganic Fertilizers to Improve Soil Fertility and Wheat (*Triticum aestivum* L.) Productivity in Semi-arid Region. *Annals of Arid Zone*, 60(3 & 4), 85-92. Retrieved from <https://eprints.icar.org.in/index.php/AAZ/article/view/131562>

Comment [PS7]: The statement regarding objective and aim of the research is missing. So kindly add it.

longitude at an altitude of 258 meters above mean sea level (MSL). In Rajasthan, this region falls under the Agro-Climatic Zone-V (Humid South Eastern Plain Zone). This zone possesses typical subtropical conditions with maximum temperature range in summer is 42.2-43.0°C and minimum 12 - 27°C. In this zone annual average rainfall is received 840 mm. The soil of experimental site was clay loam in texture, slightly saline in reaction, medium in available nitrogen (264 kg ha⁻¹) and phosphorus (21.7 kg ha⁻¹) while high in potassium (388 kg ha⁻¹) and sufficient in DTPA extractable micronutrients with pH neutral (7.61) and EC (0.52 dS m⁻¹). Source of nutrients were applied urea for nitrogen, DAP for phosphorus and mutate of potash for potassium. The full dose of fertilizer 100 % RDF (20:40:30 NPK kg ha⁻¹) was applied as basal dose. Before sowing, seed of urdbean was inoculated with liquid *rhizobium* culture @ 10 ml kg⁻¹ seed and liquid *PSB* culture @ 10 ml kg⁻¹ seed as per treatment. Foliar fertilization of zinc sulphate (ZnSO₄) and ferrous sulphate (FeSO₄) were applied at pre flowering and pod formation stage. Spraying was done with knapsack sprayer and the leaves were wetted thoroughly with fine mist. The absorption of solution by urdbean leaves, a sticker was added in the spray solution.

The experiment laid out in randomized block design with three replications. The experiment comprised 10 treatments viz. (Control, 75 % RDF, 100 % RDF, 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation, 75 % RDF + *PSB* @ 600 g ha⁻¹ seed inoculation, 75 % RDF + *Rhizobium* @ 600g ha⁻¹ + *PSB* @ 600 g ha⁻¹, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹, 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹, 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹) was carried out in randomized block design with three replications. For recording pre and post-harvest observations, five plants were randomly selected for each plot and tagged with labels for various observations on growth parameters and yield. Data were recorded plot wise as per standard procedures and statistically analysed by adopting appropriate method of standard analysis of variance [9].

Chlorophyll content (mg g⁻¹)

Total chlorophyll content of leaves at 45 DAS was determined by the method advocated by [10]. Take 20 mg of leaf

Formatted: Highlight

Formatted: Highlight

Comment [PS8]: Also write about the EC.

Comment [PS9]: It is a suggestion that it will be better if the author should provide the information in the form of

sample and crush and suspended in test tube containing 5ml of Dimethyl Sulphoxide (DMSO). Test tube were

incubated at 60°C for one hour in oven after that take reading on spectrophotometer at 645 and 663 nm.

$$\text{Total chlorophyll content (mg g}^{-1}\text{)} = \frac{(20.2 \times A_{645}) + (8.03 \times A_{663})}{1000 \times \text{weight of leaf sample taken (g)}}$$

RESULTS AND DISCUSSION

Effect of Fe, Zn and bio-fertilizer on growth parameters

The growth of urdbean crop was measured in terms of plant population, plant height, nodules plant⁻¹, nodule fresh weight, chlorophyll content and dry matter production presented in Table 1.0. A perusal of data indicated that the plant populations at 30, 60 DAS and at harvest and plant height at 30 DAS did not significantly influence, among all the treatments. It is evident from the data, plant height increased with the advancement in the age of the plant and reached the maximum at harvest. Significantly higher plant height at 60 DAS (25.28 cm) and at harvest (34.43 cm) was recorded with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ culture inoculation in urdbean as compared to control (12.33 and 16.18 cm), 75 % RDF (14.67 and 19.41 cm), 100 % RDF (16.42 and 21.96 cm), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation (17.48 and 23.32 cm), 75 % RDF + PSB @ 600 g ha⁻¹ seed inoculation (17.26 and

22.73 cm), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ (18.23 and 24.67 cm), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage (17.46 and 23.42 cm), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹ (20.59 and 26.06 cm), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + PSB @ 600 g ha⁻¹ (19.39 and 26.35 cm) in urdbean. The number of nodules plant⁻¹ at 45 DAS (29.22) was registered significantly higher under 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ culture inoculation over control (14.06), 75 % RDF (18.42), 100 % RDF (20.12), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation (22.30), 75 % RDF + PSB @ 600 g ha⁻¹ seed inoculation (21.0), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ (22.58), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage (22.68), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage +

Comment [PS10]: Write only Growth Parameters, similarly follow for other heading

Rhizobium @ 600 g ha⁻¹(24.99), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹(24.03) in urdbean. Significantly higher fresh nodule weight of urdbean(86.31 mg plant⁻¹) at 45 DAS was observed with application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ culture inoculation control (41.17mg plant⁻¹), 75 % RDF (49.71mg plant⁻¹), 100 % RDF (55.28mg plant⁻¹), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation (59.10mg plant⁻¹), 75 % RDF + *PSB* @ 600 g ha⁻¹ seed inoculation (58.20mg plant⁻¹), 75 % RDF + *Rhizobium* @ 600g ha⁻¹ + *PSB* @ 600 g ha⁻¹(61.40mg plant⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage (60.03mg plant⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹(68.66mg plant⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹(67.96mg plant⁻¹).

The chlorophyll content at 45 DAS (3.19 mg g⁻¹) was recorded significantly higher under 75% RDF + Fe 0.1 % FeSO₄ + Zn 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹

culture inoculation over control (1.71 mg g⁻¹), 75 % RDF (1.93 mg g⁻¹), 100 % RDF (2.04 mg g⁻¹), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation (2.17 mg g⁻¹), 75 % RDF + *PSB* @ 600 g ha⁻¹ seed inoculation (2.13 mg g⁻¹), 75 % RDF + *Rhizobium* @ 600g ha⁻¹ + *PSB* @ 600 g ha⁻¹(2.25 mg g⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage (2.25 mg g⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹(2.61 mg g⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹(2.38mg g⁻¹). Among the treatments, application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + *PSB* @ 600 g ha⁻¹ culture inoculation exerted significantly higher dry matter accumulation (2.71, 5.94 and 9.99g plant⁻¹) at 30, 60 DAS and at harvest over control (1.14, 2.18 and 4.25 g plant⁻¹), 75 % RDF (1.38, 2.53 and 5.15 g plant⁻¹), 100 % RDF (1.68, 3.63 and 6.29 g plant⁻¹), 75 % RDF + *Rhizobium* @ 600 g ha⁻¹ seed inoculation (1.71, 3.57 and 6.43 g plant⁻¹), 75 % RDF + *PSB* @ 600 g ha⁻¹ seed inoculation (1.71, 3.67 and 6.38g plant⁻¹), 75 % RDF + *Rhizobium* @ 600g ha⁻¹ + *PSB* @ 600 g ha⁻¹(1.75, 3.76 and 6.55 g plant⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod

formation stage (1.74, 3.70 and 6.43g plant⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha⁻¹ (2.17, 4.59 and 7.88 g plant⁻¹), 75 % RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ at pre flowering and pod formation stage + *PSB* @ 600 g ha⁻¹ (2.09, 4.48 and 7.79 g plant⁻¹) at 30, 60 DAS and at harvest in urdbean.

In the present study application of ferrous and zinc and bio-fertilizers affected the overall growth of urdbean plants measured in terms of plant height, fresh weight of nodule, chlorophyll content and dry matter accumulation. The significant increase in plant height was found due to application of ferrous, zinc and bio-fertilizers. *Rhizobium* inoculation, which may maintain favourable balance between the applied nutrients in the plant for its optimum growth while elongation and chlorophyll biosynthesis. [Tripathi et al. [11]] found similar result related to plant height with [12]. The foliar application of nutrient helps in spreading of root system and gives more site for rhizobia infection, and increase their proliferation in rhizosphere, helps in forming more effective root nodules. These results were in accordance with the findings of [13]. The increase in these growth attributes due to foliar application of Zn and Fe were also reported by [14]. The increase in

chlorophyll synthesis due to application of Fe, Zn and bio-fertilizers enhances the activity of participation of Fe in the formation of chlorophyll and Zn enzymatic role in starch formation and in protein synthesis. The higher nitrogen, phosphorus and zinc content was increased the availability throughout the growth period and reproductive period of black gram leading to increase the number of cell and size resultant profound vegetative growth in terms of plant height, dry matter accumulation, nodule count and their fresh weight was observed [15]. The significant increase of dry matter accumulation was due to the fact that nitrogen helps in maintaining higher auxin level which might have resulted in better plant height, leaf area and presumably chlorophyll content of the leaves [16]. Increase in chlorophyll content due to the stimulation effect of Fe, Zn and Bio-fertilizers on plant processes helped in cell division and root elongation. These results were in accordance with the findings of [17]. Zinc helps in chlorophyll formation and plays an important role in nitrogen metabolism. The application of zinc in a deficient soil improved the overall growth and development of plant was observed by [18].

Effect of Fe, Zn and bio-fertilizer on seed yield

Comment [PS14]: Kindly add the following statement to strengthen the discussion, "Moreover, the micronutrient (Fe and Zn) provided through foliar application leads to better uptake of the nutrient, thus enhanced growth and yield (Dey et al., 2021). Dey, S., Prasad, S., Tiwari, P., & Sharma, P. (2017). Effect of urea, KCl, Zinc placement & spray on growth of cowpea. Journal of Pharmacognosy and Phytochemistry, 6(6S), 971-973.

Comment [PS11]: Need to re-write the statement.

Comment [PS12]: The supporting citation should be provided.
For instance:
"Sharma et al., 2021 also reported the similar that "PSB culture increases the number of root nodules/plant in mungbean." Sharma, I., Singh, R. K., Tiwari, P., & Sharma, P. (2017). Effect of different Spacing and Phosphorus levels on growth and yield parameters of Mungbean under guava based Agri-horti System. Journal of Pharmacognosy and Phytochemistry, 6(6S), 993-996.

Comment [PS13]:
"Moreover, the micronutrient (Fe and Zn) provided through foliar application leads to better uptake of the nutrient than soil application, thus enhanced growth and yield (Dey et al., 2021). Dey, S., Prasad, S., Tiwari, P., & Sharma, P. (2017). Effect of urea, KCl, Zinc placement & spray on growth of cowpea. Journal of Pharmacognosy and Phytochemistry, 6(6S), 971-973.

Seed yield is a function of various growth and yield attributing parameters. The data pertaining to seed yield (kg ha^{-1}) as influenced by application of Fe, Zn and Bio-fertilizers was presented in Table 2.0. Significantly higher seed yield ($1269.30 \text{ kg ha}^{-1}$) was recorded with application of 75% RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4 foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha^{-1} + PSB @ 600 g ha^{-1} culture inoculation as compared to control (605 kg ha^{-1}), 75 % RDF (733 kg ha^{-1}), 100 % RDF (817 kg ha^{-1}), 75 % RDF + *Rhizobium* @ 600 g ha^{-1} seed inoculation (867 kg ha^{-1}), 75 % RDF + PSB @ 600 g ha^{-1} seed inoculation (854 kg ha^{-1}), 75 % RDF + *Rhizobium* @ 600 g ha^{-1} + PSB @ 600 g ha^{-1} (881 kg ha^{-1}), 75 % RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4 at pre flowering and pod formation stage (902 kg ha^{-1}), 75 % RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4 at pre flowering and pod formation stage + *Rhizobium* @ 600 g ha^{-1} (1004 kg ha^{-1}), 75 % RDF + 0.1 % FeSO_4 + 0.5 % ZnSO_4 at pre flowering and pod formation stage + PSB @ 600 g ha^{-1} (993 kg ha^{-1}) seed yield of urdbean,

The significant improvement in seed yield under the influence of application of Fe, Zn and bio-fertilizers was largely a function of improved growth and the consequent increase in different

yield attributes [19]. It might also be due to availability of nutrients to the urdbean crop which increased number of pods plant^{-1} . This might have significantly increased the number of pods plant^{-1} . Similar results were also reported by [20] for number of pods of chickpea when nutrients applied at the initial stages, might have been effectively absorbed and translocated to the pods resulting in more number of pods plant^{-1} . Similar result was also found by [21]. The result showed that test weight was not significantly influenced by the different treatments due to adequate supply of zinc and iron contributed to accelerate the enzymatic activity and auxin metabolism in plants, as auxins are involved in cell division and root formation resulted in more number of pods plant^{-1} and number of seeds pod^{-1} . These results are in close conformity with those of [18], who reported significant increase in the number of pods plant^{-1} and number of seeds pod^{-1} in urdbean due to application of zinc and iron. The profound influence of nutrient application on biological yield seems to be on account of its influence on vegetative and reproductive growth [22]. The increase in growth attributes due to application of 75% RDF might have improved the yield parameters of black gram [23]. Application of *Rhizobium* and PSB significantly increased pods plant^{-1} , seeds pod^{-1} as well

Comment [PS15]: Supporting citation should be provided.

as seed, straw and biological yield over absolute control due to the fact that *Rhizobium* inoculation increased the root nodulation through better root development and more nutrient availability ultimately there was beneficial effect on seed yield. The test weight and harvest index remain non-significant due to application of different biofertilizers. The findings of this investigation confirm the observations of earlier workers, [24] and [25]. The significant increase in yield due to zinc and iron fertilization could be attributed to the increased plant growth and biomass production, possibly as a

REFERENCES

1. Anonymous. The Directorate of Pulses Development. Pulses in India *Retrospect and Prospects*. Bhopal, M. P. 2017.
2. Anonymous. Agricultural Statistics at a Glance. Government of India. Ministry of Agricultural and Farmers Welfare. Department of Agriculture, Cooperation & Farmers Welfare. *Directorate of Economics and Statistics, New delhi, India*. 2020.
3. Jadhav SC, Sawant SP P, Sanap SP, Puranik UY, Prabhudesai SS and Devmore JP. Effect of micronutrients on yield, nutrient uptake and quality of coriander (*Coriandrum sativum*) in lateritic soil Konkar region.

result of the uptake of nutrients. These findings are supported by [26] and [18] in various crops.

CONCLUSION

Amongst all the treatments, application of 75% RDF + 0.1 % FeSO₄ + 0.5 % ZnSO₄ foliar spray at pre flowering and pod formation stage + *Rhizobium* 600 g ha⁻¹ + PSB @ 600 g ha⁻¹ culture inoculation gave maximum productivity. Therefore, it is suggested that most suitable dose for urdbean to obtain higher yield.

Comment [PS16]: The conclusion should be elaborated.

- International Journal of Chemical Studies*. 2017; **5**(4): 214-216
4. Laddha K C, Sharma R K, Sharma S K and Jain P M. Integrated nitrogen management in maize and its residual effect on black gram under dry land conditions. *Indian Journal of Dry land Agricultural Research and Development*. 2006; **21**(2): 177-184.
5. Saviour M N and Stalin P. Influence of zinc and boron in residual blackgram productivity. *Indian Journal of Science and Technology*. 2013; **6**(8): 5105-5108.
6. Khandelwal R, Choudhary SK, Khangarot SS, Jat MK and Singh P. Effect of inorganic and biofertilizers on productivity and nutrients uptake in cowpea [*Vigna Unguiculata* (L.) walp].

- Legume Research*.2012; **35**(3): 235-238.
7. KudiV K and Singh J K.Effect of biofertilizers and fertility levels on blackgram (*Vignamungo* L.) under custardapple (*Annona squamosa* L.) based agri-horti system in Vindhyan region of U.P.*International Journal of Agriculture Sciences*.2016;**8**(52): 2534-2537.
 8. AliM and Chandra S. Rhizobium inoculation of pulse crop. *Indian Farming*.1985; **35**(5): 22-25.
 9. Fisher RA. Statistical methods for research workers, Oliver and Boyd, Edinburg, London. 1950.
 10. ArnonD I. Copper enzymes in isolated chloroplasts, polyphenol oxidase in *beta vulgaris*. *Plant Physiology*.1949; **24**:1-15.
 11. Tripathi S, Kumar S, Kumar M, KumarA, Dhyani B P and KumarY.Effect of rhizobiuminoculation methods on growth, nodulation and yield of black gram (*Vigna mungo* L.). *International Journals of Current Microbiology and Applied Sciences*. 2021; **10**(01):1588-1598
 12. KumarA and Elamathi S. Effect of nitrogen levels and rhizobium application methods on yield attributes, yield and economics of black gram (*Vigna mungo* L.). *International Journal of Agriculture Sciences*. 2007;**3**(1): 179-180.
 13. Meena D, Bhushan C, Shukla A, Chaudhary S and Sirajudin S. Effect of foliar application of nutrients on nodulation yield attributes, yield and quality parameters of urdbean (*Vigna mungo* L.). *The Bioscan*. 2017; **12**(1): 411-414,
 14. Sitaram T, Sharma S K and Reager M L. Nutrient uptake of green gram as influenced by vermicompost and zinc in arid western Rajasthan. *Advance Research Journal of Crop Improvement*.2013;**4**(1): 65-69.
 15. Gajendra S, ChoudharyP, MeenaBL, RawatRS and JatBL.Integrated nutrient management in blackgramunder rainfed condition. *International Journal of Recent Scientific Research*. 2016; **7**(10):13875-13894.
 16. Devaraju B, Singh D and Senthivel T. Effect of foliar application of different sources of nutrients on growth and yield of blackgram under Irrigated Conditions. *International Journal of Current Microbiology and Applied Sciences*. 2018; **7**(1): 3105-3109.
 17. Balai K, Jajoria M, Verma R, Deewan P and Bairwa S K. Nutrient content, uptake, quality of chickpea and fertility status of soil as influenced by fertilization of phosphorus and zinc.

- Journal of Pharmacognosy and Phytochemistry*. 2017; **6**(1): 392-398.
18. Mahilane C and Singh V. Effect of zinc and molybdenum on growth, yield attributes, yield and protein in grain on summer blackgram (*Vigna mungo* L.). *International Journal of Current Microbiology and Applied Science*. 2018; **7**(1): 1156-1162.
 19. Pal S S. Acid tolerant strains of phosphatesolubilizing bacteria and their interactions in soybean-wheat crop sequence. *Journal of the Indian Society of Soil Science*. 2010; **45**: 742-746.
 20. Venkatesh M S and Basu P S. Effect of foliar application of urea on growth, yield and quality of chickpea under rainfed conditions. *Journal of Food Legumes*. 2011; **24**(2): 110-112.
 21. Thakur V, Patil R P, Patil J R, Suma T C and Umesh M R. Influence of foliar nutrition on growth and yield of blackgram under rainfed condition. *Journal of Pharmacognosy and Phytochemistry*. 2017; **6**(6): 33-37.
 22. Jangir C K, Singh D and Kumar S. Yield and economic response of biofertilizer and fertility levels on black gram (*Vigna mungo* L.). *Progressive Research – An International Journal*. 2017; **11**(8): 5252-5254.
 23. Rathore D S, Purohit H S and Yadav B L. Integrated phosphorus management on yield and nutrient uptake of urdbean under rainfed conditions of Southern Rajasthan. *Journal of Food Legumes*. 2010; **23**: 128-131.
 24. Sheikh T A, Akbar P I, Bhat A R and Khan I M. Response to biological and inorganic nutritional applications in Black gram (*Vigna mungo* L.). *Journal of Environmental Biology*. 2012; **35**: 851-854.
 25. Kumawat P K, Tiwari R C, Golada S L, Godara A and Garhwal R S. Effect of phosphorus sources, levels and biofertilizers on yield attributes, yield and economics of black gram (*Phaseolus mungo* L.). *Legume Research*. 2013; **36**(1): 70-73.
 26. Tak S, Sharma S K and Reager M L. Growth attributes and nutrient uptake of green gram as influenced by vermicompost and zinc in arid western Rajasthan. *Advance Research Journal of Crop Improvement*. 2014; **2**(1): 65-69.

Table 1.0: Effect of Fe, Zn and Bio-fertilizers application on growth parameters and seed yield of urdbean.

Treatments	Plant height (cm)			Nodules plant ⁻¹	Nodule fresh weight (mg plant ⁻¹)	Chlorophyll content (mg g ⁻¹)	Dry matter accumulation (g plant ⁻¹)			Seed yield (kg ha ⁻¹)
	30 DAS	60 DAS	At harvest				30 DAS	60 DAS	At harvest	
T ₁ : Control	8.97	12.33	16.80	14.06	41.17	1.71	1.14	2.18	4.25	604.78
T ₂ : 75 % RDF	9.33	14.67	19.41	18.42	49.71	1.93	1.38	2.53	5.15	733.36
T ₃ : 100 % RDF	9.96	16.42	21.96	20.12	55.28	2.04	1.68	3.63	6.29	817.20
T ₄ : T ₂ + <i>Rhizobium</i> @ 600 g ha ⁻¹ seed inoculation	10.12	17.48	23.32	22.30	59.10	2.17	1.71	3.57	6.43	867.24
T ₅ : T ₂ + PSB @ 600 g ha ⁻¹ seed inoculation	9.99	17.26	22.73	21.00	58.20	2.13	1.71	3.67	6.38	853.63
T ₆ : T ₂ + <i>Rhizobium</i> @ 600g ha ⁻¹ + PSB @ 600 g ha ⁻¹	10.18	18.23	24.67	22.58	61.40	2.25	1.75	3.76	6.55	881.23
T ₇ : T ₂ + 0.1 % FeSO ₄ + 0.5 % ZnSO ₄ at pre flowering and pod formation stage	10.84	17.46	23.42	22.68	60.03	2.25	1.74	3.70	6.43	901.78
T ₈ : T ₂ + T ₇ + <i>Rhizobium</i> @ 600 g ha ⁻¹ seed inoculation	11.16	20.59	26.06	24.99	68.66	2.61	2.17	4.59	7.88	1003.66
T ₉ : T ₂ + T ₇ + PSB @ 600 g ha ⁻¹ seed inoculation	10.93	19.39	26.35	24.03	67.96	2.38	2.09	4.48	7.79	993.05
T ₁₀ : T ₂ + T ₇ + <i>Rhizobium</i> 600 g ha ⁻¹ + PSB @ 600 g ha ⁻¹ seed inoculation	11.86	25.28	34.43	29.22	86.31	3.19	2.71	5.94	9.99	1269.30
SEm±	0.59	1.00	1.37	0.97	4.02	0.11	0.10	0.20	0.39	43.86
CD (p=0.05)	NS	2.98	4.06	2.89	11.95	0.33	0.30	0.60	1.17	130.30

Formatted: Highlight

UNDER PEER REVIEW

