

Application of uricase isolated from *Bacillus subtilis* SP6 in uric acid assay diagnostic kit

ABSTRACT

Uricase enzyme is a major component of the diagnostic kit for the estimation of uric acid for diagnosis, monitoring and treatment of gout and joint inflammations. Uric Acid/Uricase assay kit is a simple assay for measuring uric acid concentrations in biological samples such as serum, plasma, and urine without any need for pretreatment. The level of uricase enzyme activity was detected in the crude extract of some animal liver tissues, plant leaves and a microbial source (*Bacillus subtilis* SP6 bacteria) and expressed as specific activity (unit/mg protein). It was found difficult to produce high yield of uricase in short time from animal or plant tissues and stability of uricase from both sources is still unclear; therefore, it was isolated from *Bacillus subtilis* SP6. The extraction procedure of uricase from *Bacillus subtilis* SP6 involved, isolation and extraction of bacterial cells, determining the uricase activity in both intracellular and extracellular fractions, pooling for both fractions and ammonium sulfate precipitation which seemed to be convenient since 74.7 % of uricase activity was recovered. The isolated uricase was applied in the preparation of uric acid diagnostic kit that found sensitive and comparable with commercially available ones.

Keywords: Uricase; mammalian tissues; plant tissues; *Bacillus subtilis* SP6; diagnostic kit

INTRODUCTION

“For humans and higher primates, uric acid is the final oxidation end product of purine nucleotide metabolism. The enzyme xanthine oxidase produces uric acid from xanthine and hypoxanthine, which are derived from purines. Although most animals can metabolize uric acid to the easily excreted product allantoin, humans lack the necessary enzyme, uricase, due to two nonsense mutations in the uricase gene. Uric acid is released in hypoxic conditions and is usually excreted in the urine via glomerular filtration. Approximately 70% of daily uric acid disposal occurs via the kidneys” [1-3]. “High levels of uric acid have been linked to impaired renal function, polycythemia, leukemia and consumption of foods high in nucleoproteins. Hyperuricemia induces or accelerates the development of gout, kidney stones, hypertension, metabolic syndrome and renal or cardiovascular diseases. Gout is an inflammatory condition that results from uric acid deposits within the body joints” [4-6]. “Urate oxidase or uricase (urate: oxygen oxidoreductase, EC 1.7.3.3) is an enzyme that catalyze the oxidation of uric acid to allantoin and plays an important role in purine metabolism. It catalyzes the oxidative opening of the purine ring of urate to yield allantoin, carbon dioxide and hydrogen peroxide. This enzyme is widely present in most vertebrates but is absent in humans. It was first found in bovine kidney” [7, 8]. “Uricase has been produced from various microbial sources such as bacteria, fungi and eukaryotic cells” [9-11]. Uricase has been previously isolated from various sources like microorganisms [9, 12], plants like chickpea (*Cicer arietinum* L.), broad bean (*Vicia faba* major L.), wheat (*Triticum aestivum* L.) and animals like porcine and fish [13-15]. An uricase from *Aspergillus flavus* (Uricozyme) was in the market for the treatment of tumor lysis syndrome, hyperuricemia and renal failure [16]. Uricase has also been purified from several microbial sources like *Gliocladium viride* [17], *Streptomyces* [18] and *Bacillus subtilis* SP6, a strain that was evaluated for its ability to produce uricase [19]. The objective of present study was screening of uricase activity in the locally available and safe sources as mammalian livers, plant tissues and certain bacterial source. Isolation of uricase enzyme from the source displayed the highest specific activity and using it in the preparation of uric acid diagnostic kit which is mainly used in diagnosis, monitoring and treatment of gout .

MATERIALS AND METHODS

Collection of samples:

Fresh liver samples of buffalo (*Bubalus bubalis*), camel (*Camelus dromedaries*) and sheep (*Ovis aries*) were obtained from a local slaughter-house and stored at -40 °C. The samples were collected from at least six different individuals for each animal. Different plant tissues (Pea, Chard, Clover, Guava, Mango, Wheat and Thermos leaves) were collected from the open fields and stored at -40 °C. Soil samples contaminated with chick faecal materials were collected from certain poultry farm for isolation of *Bacillus subtilis* SP6 bacteria.

Chemicals

Uric acid, glucose, Beef extract, yeast extract, phenol red, BSA (bovine serum albumin), agar powder, peptone, manganese sulfate, 4-aminoantipyrine, peroxidase from horseradish, diaminobenzidine, 2,4,6-tribromo-3-hydroxybenzoic acid (TBHBA) and chemicals for electrophoresis were purchased from Sigma-Aldrich Chemical Co. The other chemicals were of analytical grade.

Assay of uricase activity

“The uricase activity reaction mixture contained 0.6 ml of 2 mM uric acid dissolved in 0.1 M sodium borate buffer (pH 8.5), 0.1 ml of 1.5% phenol, 0.15 ml of 30 mM 4-aminoantipyrine, 0.05 ml of 15 U/ml peroxidase from horseradish, and 0.1 ml of enzyme solution. The mixture was incubated at 25 °C for 20 min. The reaction was terminated by the addition of 1 ml of ethanol, and the absorbance at 540 nm was read against the blank. One unit of uricase was defined as the amount of enzyme that produces 1.0 μmol of H₂O₂ per minute under the standard assay conditions. To determine the uricase activity, samples of the culture were withdrawn, and the cells were

centrifuged at 3000g for 10 min and filtered through 0.2 μ m. The filtered supernatant (crude enzyme solution) was used for the analysis of enzyme activity” [20].

Uricase activity staining on polyacrylamide gels

“Activity staining of uricase was determined after electrophoresis by submerging the gel in 0.1 M Tris-HCl, pH 9.0, 0.1 mM urate, 2.8 mM diaminobenzidine and 3.3 U of horseradish peroxidase and staining was continued till the activity band appears on the gels” [21].

Extraction of proteins from animal and plant tissues:

All of the procedures were performed at 4 °C unless stated otherwise. 10 grams of animal liver were homogenized in 50 mM Tris-HCl buffer, pH 8.0 using Omni mixer homogenizer. Cell debris and insoluble materials were removed by centrifugation at 8000 x g for 20 min and the supernatant was saved and designated as crude extract. Pea, Chard, Clover, Guava, Mango, Wheat and Thermos leaves were collected and proteins were extracted according to Lanna *et al.*, [22]. “One gram fresh weight was ground in a mortar and pestle containing liquid nitrogen. The resulting powder was macerated for 30 sec in 3 ml extraction buffer [50 mM Tris-HCl buffer, pH 8.0], then centrifuged at 20,000 g for 25 min at 4 °C. The supernatant was divided and kept in ice at -20 °C for the following determination” [22].

Isolation of bacteria

“*Bacillus subtilis* SP6 formerly isolated in our lab from poultry waste soil samples was used for the production of uricase enzyme” [19]. “Soil samples contaminated with chick faecal materials were collected from certain poultry farm. The culture was maintained on uric acid agar slants and stored at 4 °C. For uricase production, the composition of pre-culture medium was 1% peptone, 2% glucose, 0.1% K₂HPO₄, 0.05% MgSO₄.7H₂O and 0.5% NaCl. The pH of the medium was adjusted to 8.0. The production medium was prepared by the addition of 0.05% uric acid to the pre-culture medium” [19].

Cell growth and uricase production

1% (v/v) of pre-cultured organism was inoculated into a 3000 ml Erlenmeyer flask containing 500 ml of sterilized production medium. The flask was incubated in an orbital shaker incubator for 48 h at 175 rpm maintained at 37 °C. Samples were taken periodically to determine the cell biomass and uricase production.

Extraction and partial purification of uricase

“The cells free culture broth was sonicated for 21 sec at max speed then centrifuging at 5000 rpm for 20 min. The uricase enzyme in the cell filtrate was further concentrated by 80% ammonium sulfate precipitation, centrifuged at 10000 rpm for 30 min followed by dialysis. Enzyme activity was determined for each separate fraction” [23].

Electrophoretic analysis

Native gel electrophoresis was carried out with 7% PAGE [24] and proteins were stained with 0.25% Coomassie Brilliant Blue R-250.

Protein determination

“Protein content was determined by the dye binding assay method using bovine serum albumin (BSA) as a standard protein” [25].

Construction of uric acid diagnostic kit

The partially purified *Bacillus subtilis* SP6 uricase enzyme was used in the construction of uric acid diagnostic kit. The uric acid kit is composed of two reagents: (R1: 100 mM phosphate buffer pH 7.0 and 1.25 mM TBHBA (2, 4, 6-Tribromo- 3-hydroxybenzoic acid) and (R2: 100 mM phosphate

buffer pH 7.0, 1.5 mM 4-Aminoantipyrine, 50 mM $K_4 [Fe (CN)_6]$, ≥ 10 kU/L Peroxidase (POD) and ≥ 150 U/L bacterial uricase). The standard: 6 mg/dL (357 μ mol/L) uric acid [26].

RESULT AND DISCUSSION

Comparison of uricase activity

“In gout, the increased serum levels of uric acid lead to formation of mono-sodium urate crystals around the joints. Further causes of elevated blood concentrations of uric acid are renal diseases with decreased excretion of waste products, starvation, drug abuse and increased alcohol consume as well as use of certain medicaments. High uric acid levels also constitute an indirect risk factor for coronary heart disease. Uricase catalyzes the oxidation of uric acid to allantoin and plays an important role in purine metabolism. It catalyzes the oxidative opening of the purine ring of urate to yield allantoin, carbon dioxide, and hydrogen peroxide” [27, 28, 1, 2]. The level of uricase activity was detected in the crude extract of different samples and expressed as specific activity (units / mg protein) and the data of uricase specific activity are summarized in (Table 1). The specific activity of the uricase was found to be (0.214 units / mg protein) for buffalo liver, (0.015 units / mg protein) for camel liver, (0.185 units / mg protein) for sheep liver, (0.19 units / mg protein) for Pea leaves, (0.085 units / mg protein) for Chard leaves,, (0.24 units / mg protein) for Clover leaves, (0.87 units / mg protein) for Guava leaves, (0.24 units / mg protein) for Mango leaves, (0.1 units / mg protein) for Wheat leaves, (0.194 units / mg protein) for Thermos leaves and (0.92 units / mg protein) for the *Bacillus subtilis* SP6 bacteria. The specific activity of uricase from Guava leaves and the bacterial source is higher than other sources. Both of the protein and uricase isoenzyme patterns of different samples were compared by analysis on 7 % native PAGE. For comparison, a definite amount of protein (100 μ g) was applied for protein and uricase isoenzyme patterns (Fig. 1). The uricase isoenzyme pattern indicated the presence of one isoenzyme in the bacterial crude extract. Different protein patterns were monitored on the native PAGE from different tissues. Uricase has been produced from various microbial sources such as bacteria, fungi and eukaryotic cells [9-11]. Uricase from leaves of chickpea, broad bean and wheat has been purified to electrophoretic homogeneity [14].

Extraction of *Bacillus subtilis* SP6 bacteria uricase

In this study, it was found difficult to produce high amounts of uricase enzyme in short time either from animal or plant tissues. Therefore, we isolated the desired amount of uricase from the intracellular and the extracellular fractions of *Bacillus subtilis* SP6. The two fractions were pooled together, brought to 80% saturation by gradually adding solid $(NH_4)_2SO_4$, stirred for 30 min at 4 °C and centrifuged at 10000 x g for 20 min. The pellet of this step was dissolved in 0.05 M sodium phosphate buffer pH 7.0 and dialyzed extensively against the same buffer. Most of the uricase activity was precipitated in the ammonium sulfate fraction that 74.7 % of the activity was recovered. The uricase specific activity was found to be 1226.4 m units / mg protein (Table 2). Various uricase specific activities were reported; 0.007 U/mg for broad bean, 0.11 U/mg for chickpea and 0.009 mU/mg for wheat [14], 32 U/mg for *Bacillus subtilis* [8], 0.131 U/ml for *Halobacillus sp* [29], 0.05 U/mg for *Candida sp.* [30] and the intracellular uricase ranged from 0.09 to 0.5 U/ml and the extracellular ranged from 0.08 to 0.43 U/ml for *Streptomyces exfoliates* UR10 [23].

Comparison of the constructed uric acid kit with commercially available kits

The isolated *Bacillus subtilis* SP6 uricase is used in the construction of uric acid diagnostic kit. The constructed kit (prepared uric acid diagnostic kit with the isolated uricase enzyme) has been compared with commercially available kits utilizing different individual samples (Table 3 & Fig. 2). Uric acid diagnostic kit is very sensitive in the measurement of uric acid concentration and mainly used in diagnosis, monitoring and treatment of gout and joint inflammations. The prepared kit was found to be comparable to the commercial kits. The variance between the constructed and commercial kits was found more or less within the experimental error.

CONCLUSION

In conclusions, the uricase enzyme has been isolated from *Bacillus subtilis* SP6 bacteria as a newer source of uricase which is cost effective as well as having more specificity that makes it suitable for various medical applications. The isolated bacterial uricase enzyme was used in the preparation of uric acid diagnostic kit which found comparable to commercially available kits.

ACKNOWLEDGEMENTS

This work was funded by the National Research Centre, Egypt (Agreement No. E121101).

Availability of data and material

All data and materials are available.

Competing interests

No potential conflict of interest was reported by the authors.

Authors' contributions

All authors contributed to the experimental design, hands on work, discussions, and commented on the manuscript.

REFERENCES

1. Heinig M, Johnson RJ. Role of uric acid in hypertension, renal disease, and metabolic syndrome. *Cleve Clin J Med*. 2006;73(12): 1059-1064.
2. Shani M, Vinker S, Dinour D, Leiba M, Twig G, Holtzman EJ, Leiba A. High normal uric acid levels are associated with an increased risk of diabetes in lean, normoglycemic healthy women. *J Clin Endocrinol Metab*. 2016;101(10): 3772-3778.
3. Watanabe S, Kang DH, Feng L, Nakagawa T, Kanellis J, Lan H, Mazzali M, Johnson RJ. Uric acid, hominoid evolution, and the pathogenesis of salt-sensitivity. *Hypertension*. 2002;40: 355-360.
4. Votyakova TV, Reynolds IJ. DeltaPsi(m)-Dependent and -independent production of reactive oxygen species by rat brain mitochondria. *Neurochem*. 2001;79:266 .
5. Ichida, K. What lies behind serum urate concentration? Insights from genetic and genomic studies. *Genome Med*. 2009;1(12): 118.
6. Moriya C, Satoh H. Teneligliptin decreases uric acid levels by reducing xanthine dehydrogenase expression in white adipose tissue of male wistar rats. *J Diabetes Res*. 2016;doi:10.1155/2016/3201534.
7. Schiavon O, Calicati P, Ferruti P, Veronese FM. Therapeutic proteins: a comparison of chemical and biological properties of uricase conjugated to linear or branched poly (ethylene glycol) and poly (N-acryloylmorpholine). *Il Farmaco*. 2000;55: 264-69.
8. Jagathy K, Pushparaj A, Ronald J. Uricase production from *Bacillus subtilis* isolated from Poultry waste. *Int. J. Adv. Res. Biol. Sci*. 2016;3(6): 255-262.

9. Farley PC, Santosa S. Regulation of expression of the *Rhizopus oryzae* uricase and urease enzymes. *Can J Microbiol.* 2002;48: 1104-1108.
10. Ishikawa J, Yamashita A, Mikami Y. The complete genomic sequence of *Nocardia farcinica* IFM 10152. *Proc. Natl. Acad. Sci. U.S.A.* 2004;101: 14925-14930.
11. Zhou X, Ma X, Sun G, Li X, Guo K. Isolation of a thermostable uricase producing bacterium and study on its enzyme production conditions. *Proc. Biochem.* 2005;40: 3749-3753.
12. Abdollah GA, Moradpour Z, Baniasad M, Ghasemi Y. Isolation, molecular identification and characterization of the culture conditions for extracellular uricase production by a new strain of *Pseudomonas* sp. *J Pure Appl Microbio.* 2015;9:2813-2821.
13. Conley TG, Priest DG. Purification of uricase from mammalian tissue, *Prep Biochem Biotechnol.* 1979;9 (2): 197-203.
14. Montalbini P, Redondo J, Caballero JL, Cardenas J, Pineda M. Uricase from leaves: its purification and characterization from three different higher plants. *Planta Heidelberg.* 1997;202: 277-283.
15. Kinsella E, German B, Shetty J. Uricase from fish liver: isolation and some Properties. *Comp Biochem Physiol.* 1985;82 (4): 621-624.
16. Pui C.H, Relling MV, Lascombes F, Harrison PL, Struxiano A, Mondesir JM, Ribeiro RC, Sandlund JT, Rivera GK, Evans WE, Mahmoud HH. Urate oxidase in prevention and treatment of hyperuricemia associated with lymphoid Malignancies. *Leukemia.* 1997;11:1813-1816.
17. El-Naggar NE. Isolation, screening and identification of actinobacteria with uricase activity: statistical optimization of fermentation conditions for improved production of uricase by *Streptomyces rochei* NEAE-25. *Int J Pharmacol.* 2015;11: 644-658.
18. Nanda P, Babu JP, Fernandes J, Hazarika P, Dhabre RR. Studies on production, optimization and purification of uricase from *Gliocladium viride*. *Res Biotechnol.* 2012;3 (4): 35-46.
19. Pustake S, Bhagwat P, Dandge P. Statistical media optimization for the production of clinical uricase from *Bacillus subtilis* strain SP6. *Heliyon.* 2019;5: e01756.
20. Ravichandran R, Hemaasri S, Cameotra SS, Jayaprakash NS. Purification and characterization of an extracellular uricase from a new isolate of *Sphingobacterium thalpophilum* (VITPCB5). *Protein Exp. Purific.* 2015;114: 136-142.
21. Pineda M, Fernandez E, Cardenas J. Urate oxidase of *Chlamydomonas reinhardtii*. *Physiol Plant.* 1984;62: 453-457.
22. Lanna AC., Oliveira MGA, Barros EG, Moreira MA. Kinatic parameters of leaf lipoxygenase pool from normal soybean genotypes and from a line devoid of seed lipoxygenase. *Rev.Bras.Fisiol.Vegetal.* 1996;8: 87-92.
23. Aly M, Tork S, Al-Garni S, Allam R. Production and characterization of uricase from *Streptomyces exfoliates* UR10 isolated from farm wastes. *Turk J Biol.* 2013;37: 520-529.
24. Smith I. Acrylamide gel disc electrophoresis. In "Electrophoretic techniques" (Edited by Smith, I.) 1969;pp. 365-515, Academic press, New York.

25. Bradford M.M.. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 1976;72: 248-254.
26. Bakker AJ, Mücke M. Gammopathy interference in clinical chemistry assays: mechanisms, detection and prevention. *Clin Chem Lab Med.* 2007;45(9):1240-1243.
27. Thomas L. *Clinical Laboratory Diagnostics.* 1st ed. Frankfurt: TH-Books Verlagsgesellschaft; 1998;pp. 208-214.
28. Newman DJ, Price CP. Renal function and nitrogen metabolites. In: Burtis CA, Ashwood ER, editors. *Tietz Textbook of Clinical Chemistry.* 3rd ed. Philadelphia: W.B Saunders Company; 1999;pp. 1204-1270.
29. Honarbakhsh F, Abolmaali S, Amoozegar MA. Uricase activity of halophilic bacteria from Iranian salt lakes. *Iran J Sci Technol Trans Sci.* 2021;45:1597-1606.
30. Liu J, Li G, Liu H, Zhou X . Purification and properties of uricase from *Candida* sp. and its application in uric acid analysis in serum. *Appl Biochem Biotechnol.* 1994;47(1): 57-63.

Table (1): uricase specific activity of different animal, plant and microbial tissues:

Sample	Uricase specific activity*
Buffalo liver	0.214
Camel liver	0.015
Sheep liver	0.185
Pea leaves	0.19
Chard leaves	0.085
Clover leaves	0.24
Guava leaves	0.87
Mango leaves	0.24
Wheat leaves	0.1
Thermos leaves	0.194
<i>Bacillus subtilis</i> SP6 bacteria	0.92

* The specific activity is expressed as units / mg protein.

* Each value represents the mean of at least 4 tissues.

Table (2): Purification scheme of uricase from *Bacillus subtilis* SP6 bacteria:

Purification step	Total protein (mg)	Total Activity (m unit)	Specific Activity (mU/mg protein)	Yield (%)	Fold Purification
<i>Bacillus subtilis</i> SP6 intracellular fraction	6.7	6300	940.3	100.0	1.0
<i>Bacillus subtilis</i> SP6 extracellular fraction	3.5	2400	685.7	100.0	1.0
The pooled fraction	10.2	8700	852.9	100.0	1.0
80% (NH ₄) ₂ SO ₄ fraction	5.3	6500	1226.4	74.7	1.4

The specific activity is expressed as m units / mg protein.

Table (3): Comparison of the constructed uric acid kit with commercially available kits:

Serum samples	Uric acid concentration		
	commercial kit 1	commercial kit 2	Constructed kit
1	5.4	5.34	5.8
2	4.68	5.94	5.6
3	4.96	5.37	5.66
4	9.57	6.2	7.88
5	5.48	5.67	6.8
6	4.37	5.85	6.26
7	4.73	5.82	4.84
8	4.61	6.24	6.96
9	5.05	5.67	6.73
10	3.82	4.11	4.43
11	6.11	5.88	6.67
12	5.22	5.92	5.56




Fig. (1): (a) Electrophoretic analysis of protein and uricase isoenzyme patterns on 7 % native PAGE: (1) buffalo, (2) camel and (3) sheep. (b) Electrophoretic analysis of protein and uricase isoenzyme patterns on 7 % native PAGE: (1) *Bacillus subtilis* SP6 extract, (2) Guava leaves extract, (3) Mango leaves extract (4) Thermos leaves extract.




Fig. (2): Comparison of the constructed uric acid kit with commercially available kits.