

Original Research Article

Response of Different Levels of Nitrogen on Growth and Yield of Cauliflower (*Brassica oleracea* var. botrytis) Varieties

ABSTRACT

Nitrogen is the major limiting nutrient in commercial cauliflower production. The effect of six nitrogen doses (0, 70, 80, 90, 115, 130 kg N/ha) were evaluated in two cauliflower varieties (Snow Mystique and KhumalJyapu) at the Institute of Agriculture and Animal Science, Lamjung Campus from September 2018 to January 2019. This experiment was carried out in two factorial Completely Randomized Block Design with twelve treatments and five replications. FYM (25 t/ha), full dose of phosphorus (80 kg/ha) and potassium (60 kg/ha), and half dose of nitrogen were applied as a basal dose and the remaining half dose of nitrogen was applied in two equal split doses at 30 DAT and 60 DAT. The application of 130 kg N/ha gave superior results in plant height (26.32 cm), plant spread (4103 cm²), leaf area (264 cm²), stem diameter (1.96 cm), number of leaves (20.70), root length (26.26 cm), girth of curd (49.46cm) and adjusted yield (21.78 ton/ha). Similarly, the performance of the Snow Mystique variety was significantly superior as compared to KhumalJypau in terms of growth and yield. Therefore, the combination of optimum level of nitrogen (130 kg N/ha) and hybrid variety (Snow Mystique) could be recommended for the commercial production of

Keywords: Cauliflower, Growth, Nitrogen, Yield

1. INTRODUCTION

Cauliflower (*Brassica oleracea* var. botrytis L. 2n= 18) is a popular vegetable crop commercially grown from terai to high hills of Nepal with area, production, and productivity of 35402 ha, 528,738 tons, 14.94 ton/ha respectively [1]. Cauliflower is regarded as an economically important crop of the cole group having high market value and potentiality of offseason production in the different parts of the country. Besides its economic value, cauliflower is a good source of dietary fiber, vitamin C, and minerals [2]. While analyzing cauliflower's production trend, there has not been a satisfactory increase in the productivity of cauliflower in the last five years [1]. Among different factors, poor nutrient management and the use of low yielding open pollinated varieties are major limiting factors associated with the lower production of cauliflower

Comment [PS1]: Instead of writing it in value, write in text like 35.4 thousand ha likewise for yield.

in the country. Being a heavy feeder crop, the role of macronutrients (NPK) and micronutrients (boron, zinc, and molybdenum) are inevitable for the luxuriant growth and higher yield of cauliflower. The nitrogen requirement in cauliflower is much higher than other nutrients [3]. Nitrogen is directly related to the vegetative growth and curd development of cauliflower, as it involves in several physiological processes and enzymatic activities. Nitrogen is integral to chlorophyll, protoplasm, proteins, nucleic acids, and enzymes [4]. Thus, N should be applied in the optimum dose to avoid toxicity and deficiency and to reduce production costs.

Nitrogen deficiency checks the growth and reduces the yield significantly [5]. Nitrogen deficiency in plants results in poor plant performance with stunted growth, leaves discoloration, interveinal yellowing, chlorosis, acceleration of the reproductive stage, and senescence [6]. On the other hand, the excessive application of nitrogen is not only uneconomical but also stimulate vegetative growth, induces physiological disorder, negatively affect the nutritional value by limiting the synthesis of sugar and enhancing nitrate accumulation, high nitrogen leaching from soil and environment pollution [7,8,9,10]. High nitrogen applications have been shown to reduce the vitamin C content [11], and increasing nitrogen content from 80 to 120 kg/ha was reported to decrease vitamin C content by seven percent in cauliflower and broccoli [12]. Similarly, the varietal response to nitrogen fertilization may be different, different varieties have specific requirements for nitrogen. Hence, the balanced dose of nitrogen application and selection of suitable varieties is crucial for the commercial production of cauliflower. Therefore, the present study was carried out to determine the optimum level of nitrogen and suitable variety for the commercial production of cauliflower in Lamjung, Nepal.

2. MATERIALS AND METHODS

2.1 Location and Site of the Experiment

This experiment was conducted at the Institute of Agriculture and Animal Science (IAAS) Lamjung Campus from September 2018 to January 2019. It is located at 28.13° N latitude and 84.42° E longitude. During the research period, the monthly average temperature and relative humidity were assessed using the official website of the department of hydrology and meteorology. The maximum temperature (21°C) and relative humidity (75%) were observed in September whereas the minimum temperature (5°C) and relative humidity (45%) were observed in January [13].

2.2 Design of Experiment

The experiment was designed in two factorial completely randomized block design with five replications. The first factor consisted of six doses of nitrogen (control, 70 kg, 80 kg, 90 kg, 115 kg, and 130 kg) and the second factor consisted of two cauliflower varieties (Snow Mystique and KhumalJyapu).

Table 1. Treatment details

S.N.	Treatments	Treatments name	Combination
1.	T1	0 kg N + Snow Mystique	N1V1
2.	T2	70 kg N + Snow Mystique	N2V1
3.	T3	80 kg N + Snow Mystique	N3V1
4.	T4	90 kg N + Snow Mystique	N4V1
5.	T5	115 kg N + Snow Mystique	N5V1
6.	T6	130 kg N + Snow Mystique	N6V1
7.	T7	0 kg N + KhumalJyapu	N1V2
8.	T8	70 kg N + KhumalJyapu	N2V2
9.	T9	80 kg N + KhumalJyapu	N3V2
10.	T10	90 kg N + KhumalJyapu	N4V2
11.	T11	115 kg N + KhumalJyapu	N5V2
12.	T12	130 kg N + KhumalJyapu	N6V2

2.3 Preparation of potting media for transplanting

On September 25, 2018, the soil was collected from the field and registered at the Agricultural Technology Center in Jwagal, Lalitpur, Nepal for testing (Table 2). On October 1, 2018, an estimated quantity of FYM was mixed properly with the soil. After one month, pots (top diameter: 22.5 cm, base diameter: 16.5 cm, height: 18 cm) were filled with well-prepared media (FYM and soil) and watered to the saturation point, and left for three days to allow for field capacity.

Comment [PS2]: Note an appropriate word.

Formatted: Highlight

Table 2. Soil analysis report

Test	Result	Remarks
pH	6.5	Slightly Acidic
N (%)	0.10	Medium
P ₂ O ₅ (kg/ha)	8.24	Low
K ₂ O (kg/ha)	112.40	Medium
Organic matter (%)	2.01	Medium
Sand (%)	39.60	
Silt (%)	38.00	
Clay (%)	22.40	
Soil texture	Loamy	

2.4 Seedling Raising

A raised bed of 15 cm height and 1m width was made and seeds of two varieties were sown separately on 8th October 2018. Line sowing with 10 cm spacing was done at the depth of 4-5 cm. The seeds were then finely covered with a thin layer of a mixture of FYM and soil and nursery beds were mulched with straw. Light irrigation was done after seed sowing. The straw mulch was removed after five days and the overhead plastic tunnel was made to protect seedlings from extreme cold and hot. Light irrigation was done at three days intervals.

2.5 Transplanting of Seedlings

Uniform size vigorous seedlings (4-5 leaf stage and 27 days old) were transplanted in the pot containing well-prepared media (soil and FYM) and recommended doses of fertilizers. Light irrigation was done immediately after transplanting.

Comment [PS3]: Why on pot rather than planting it on the field condition.

2.6 Manure and Fertilizers Application

Nitrogen, phosphorous, and potash were applied through Urea, Single Super Phosphate (SSP,) and Muriate of Potash (MOP) respectively. FYM (25t/ha), phosphorous, and potassium (80:60 kg/ha) were applied as a basal dose. The fertilizer requirement per pot was calculated on the basis of 60x45 cm plant spacing. Half a dose of different levels of urea (0, 2.05, 2.34, 2.64,

3.37, 3.81 g pot⁻¹) was applied as a basal dose and the remaining dose of urea was applied in two equal split doses at 30 and 60 days after transplanting (DAT) by band placement. A foliar spray of boron (0.25%) was done at 30, 45, and 60 DAT and a foliar spray of molybdenum (0.2%) was done at 35 DAT.

Table 3. NPK application at basal and split dose during the experiment

Fertilizers	Basal dose (g)	Split dose (g)
Urea	0, 2.05, 2.34, 2.64, 3.37, 3.81 g pot ⁻¹	At 30 DAT (0, 1.025, 1.17, 1.32, 1.68, 1.90 g pot ⁻¹) At 60 DAT (0, 1.025, 1.17, 1.32, 1.68, 1.90 g pot ⁻¹)
Single Super Phosphate	13.50 g pot ⁻¹	
Muriate of Potash	2.70 g pot ⁻¹	

2.7 Intercultural Operations

The pots were kept weed free throughout the growth period to avoid crop weed competition. Manual weeding was done at 30 and 60 DAT and earthing up was done during the second weeding. During the weeding operation split dose of nitrogen was applied.

2.8 Observations

2.8.1 Growth parameters

2.8.1.1 Plant height (cm)

The plant height was measured with the help of a measuring scale. The plant height was measured from the ground level to the growing tip of the plant.

2.8.1.2 Plant spread (cm²)

The spreading of the plants was measured by measuring the distance between perpendicular directions at harvesting time.

2.8.1.3 Leaf area (cm²)

The length of leaves was measured between the attachment of the petiole and the apex of the leaf and breadth was measured at the point of maximum expansion with the help of a measuring scale. The area of leaves per plant was calculated by the Montgomery equation [14].

$$\text{Leaf area} = \frac{\pi}{4} * \text{Length} * \text{Width}$$

2.8.1.4 Diameter of the stem (cm)

The girth of the stem was measured with the help of Vernier's caliper at the time of harvesting.

2.8.1.5 Number of leaves per plant

The total number of fully opened leaves was counted at the time of harvesting.

2.8.1.6 Root length (cm)

The root length was measured with the help of a measuring scale. The plant was uprooted and immersed in water for one hour. The soil in the root was washed away, and the length of the root was measured from the base of the plant (ground level) to the tip of the root.

2.8.2 Yield parameters

2.8.2.1 Days to curd initiation

The number of days taken to curd initiation from the date of transplanting of seedlings was observed.

2.8.2.2 Diameter of curd(cm)

After harvesting the curd, the diameter of the curd was measured horizontally with the help of scale.

2.8.2.3 Fresh weight of leaves (g)

After harvesting, the crop fresh weight of the leaves was measured with the help of an electrical weighing balance.

2.8.2.4 Fresh weight of curd (g)

After harvesting, the fresh weight of the curd was measured with the help of an electrical weighing balance.

2.8.2.5 Adjusted yield per hectare

The weight of curd was measured by the electrical weighing balance and yield per hectare was computed by using the following formula:

$$\text{Yield per hectare} = \frac{\text{curd weight per pot (gm)} \times \text{total plant population per hectare}}{1000 \times 1000}$$

2.9 Statistical Analysis

The data were entered into Microsoft Excel 2016 and analyzed using the Genestat (15th edition). The data were subjected to Analysis of Variance (ANOVA) and mean separation was carried out using Duncan's Multiple Range Test (DMRT) at 0.05 level of significance ($P < 0.05$).

3. RESULTS

3.1 Effect of Nitrogen and Varieties on Growth Parameters of Cauliflower

Analysis of variance showed that effect of different doses of nitrogen and varieties was significant for the different growth parameters of cauliflower (Table 4). The highest mean height of the cauliflower plant was obtained in treatment applied with 130 kg N/ha (26.32 cm) and the lowest plant height (16.10 cm) was observed in the control. Maximum plant spread was observed in the treatment with 130 kg N/ha (4103 cm²) which was statistically at par with the treatment treated with 115 kg N/ha and 90 kg N/ha while minimum plant spread was observed in the control (627 cm²). The leaf area of the cauliflower plant increased significantly with the application of successive doses of nitrogen. Treatment that received 130 kg N/ha had the maximum leaf area (264 cm²) while the minimum leaf area was observed in the control treatment (35.1 cm²). The maximum stem diameter was observed in the 130 kg N/ha (1.96 cm) and the minimum stem diameter was observed in the control (0.92 cm). The highest number of leaves per plant was recorded in the 130 kg N/ha which was statistically similar to 115 kg N/ha. The lowest number of leaves per plant was observed in the control (12.70). Similarly, the root length was also maximum in the 130 kg N/ha (26.26 cm) and minimum in the control (19.50 cm). Among the varieties, the results were significant for all growth parameters except leaf area. The highest plant height (22.16 cm), plant spread (4011 cm²), diameter of the stem (1.70 cm), number of leaves per plant (20.10), and root length (26.22 cm) were observed in the Snow Mystique.

Table 4. Effect of different levels of nitrogen and varieties on growth parameters of cauliflower

Treatments	Plant height (cm)	Plant spread (cm ²)	Leaf area (cm ²)	Diameter of stem (cm)	Number of leaves per plant	Root length (cm)
Nitrogen level (kg/ha)						
0	16.10 ^e	627 ^c	35.1 ^d	0.92 ^e	12.70 ^d	19.50 ^e

70	20.92 ^d	3554 ^b	192.2 ^c	1.49 ^d	17.50 ^c	22.37 ^d
80	21.84 ^{cd}	3762 ^{ab}	194.0 ^c	1.62 ^c	18.70 ^{bc}	22.88 ^{cd}
90	22.75 ^{bc}	3915 ^a	198.4 ^{bc}	1.65 ^{bc}	20.00 ^{ab}	24.17 ^{bc}
115	23.80 ^b	4007 ^a	217.2 ^b	1.74 ^b	20.50 ^a	25.37 ^{ab}
130	26.32 ^a	4103 ^a	264.0 ^a	1.96 ^a	20.70 ^a	26.26 ^a
LSD	1.63 ^{**}	335.3 ^{**}	19.41 ^{**}	0.09 ^{**}	1.65 ^{**}	1.64 ^{**}
Varieties						
Snow	22.16	4011 ^a	185.0	1.70 ^a	20.10 ^a	26.22 ^a
Mystique						
KhumaJyapu	21.75	2645 ^b	182.0	1.40 ^b	16.60 ^b	20.57 ^b
LSD	0.77 ^{ns}	193.6 ^{**}	11.21 ^{ns}	0.05 ^{**}	0.956 ^{**}	0.97 ^{**}
CV	6.8	11.2	11.8	6.6	10.0	8.0
Mean	21.96	3328	183.5	1.6	18.35	23.40

Means separation in column followed by the same letters are not significantly different at $p=0.05$, **= highly significant, NS= Non-significant, LSD = Least Significant Difference, CV = Coefficient of Variance

3.2 Effect of Different Levels of Nitrogen and Varieties on Yield Parameters of Cauliflower

The yield parameters were significant for the different doses of nitrogen and varieties as shown by the analysis of variance (Table 5). The minimum number of days (44 days) taken for curd initiation was observed in treatment at 80 kg N /ha which was similar to 70 kg N/ha while the maximum days for curd initiation was 53.30 days in treatment applied with 130 kg/ha. The highest girth (49.46 cm) of curd was observed in 130 kg N/ha and the lowest girth was observed in the control (15.27 cm). The maximum fresh weight of leaves was observed in the treatment 130 kg N /ha (592.20 g) and the minimum weight of fresh leaves was observed in the control (33.1 g). The fresh weight of curd and adjusted yield were also increased with the successive dose of nitrogen. The highest adjusted yield was observed in 130 kg N/ha (21.78 ton/ha) and the lowest adjusted yield was observed in the control (1.05 ton/ha). Among varieties, the results were significant for the yield parameters. The minimum number of days (45.77 days) for curd initiation was observed in the KhumaJyapu and the maximum girth of curd (43.59 cm), fresh weight of leaves (583.0 g), fresh weight of curd (569.7 g) and adjusted yield (21.10 ton/ha) was observed in Snow Mystique.

Table 5. Effect of different levels of nitrogen on yield parameters of cauliflower

Treatments	Curd initiation days	Girth of curd (cm)	Fresh weight of leaves (g)	Fresh weight of curd (g)	Adjusted yield (t/ha)
Nitrogen level (kg/ha)					
0	48.80 ^b	15.27 ^e	33.10 ^d	28.50 ^e	1.05 ^e
70	45.10 ^{cd}	45.71 ^d	423.10 ^c	420.40 ^d	15.57 ^d
80	44.00 ^d	46.53 ^{cd}	467.70 ^{bc}	481.70 ^c	17.84 ^c
90	47.50 ^{bc}	47.66 ^{bc}	514.90 ^b	518.20 ^{bc}	19.19 ^{bc}
115	51.70 ^a	48.42 ^{ab}	524.20 ^b	557.90 ^{ab}	20.66 ^{ab}
130	53.30 ^a	49.46 ^a	592.20 ^a	588.20 ^a	21.78 ^a
LSD	2.751 ^{**}	1.286 ^{**}	65.81 ^{**}	53.82 ^{**}	1.993 ^{**}
Varieties					
Snow	51.03 ^a	43.59 ^a	583.0 ^a	569.7 ^a	21.10 ^a
Mystique					
KhumaJyapu	45.77 ^b	42.04 ^b	268.8 ^b	295.3 ^b	10.94 ^b
LSD	1.588 ^{**}	1.160 [*]	38.00 ^{**}	31.07 ^{**}	1.151 ^{**}
CV	6.3	5.2	17.2	13.8	13.8
Mean	48.40	42.81	425.9	432.5	16.02

Means separation in column followed by the same letters are not significantly different at $p=0.05$, ^{**}= highly significant, LSD = Least Significant Difference, CV = Coefficient of Variance

4. DISCUSSION

Nitrogen application is attributed to the vegetative growth of plants and plant height increases with the increase in the levels of nitrogen. The maximum growth under a higher dose of nitrogen might be due to increasing the photosynthetic rates and the assimilation rates, which lead to an increase in the stalk length of cauliflower. Being one of the important nutrients with its roles in plant metabolism and development, its application boosts plant growth [15]. These findings are

Comment [PS4]: Reference should be provided such as Garima et al., 2021.
Garima, Bhardwaj, D. R., Thakur, C. L., Kaushal, R., Sharma, P., Kumar, D., & Kumari, Y. (2021). Bamboo-based agroforestry system effects on soil fertility: Ginger performance in the bamboo subcanopy in the Himalayas (India). *Agronomy Journal*, 113(3), 2832-2845.

in agreement with those reported by [16] in broccoli, [5, 17] in cauliflower, [18] in Chinese cabbage and [19] in broccoli. Plant spread is directly related to the available nitrogen in the root zone of plants. When the supply of nitrogen to the root increases, the synthesis of cytokinin increases, and ultimately more cytokinin is transferred to leaves for their growth and expansion [20, 21]. These findings are in agreement with [18] in Chinese cabbage. Nitrogen in the vegetative growth of the plant enhanced leaf number in broccoli brought about by high nitrogen fertilization [22]. The maximum growth under a higher supply of nitrogen might be due to increasing the photosynthetic and assimilation rates, which lead to increased metabolic functions in the stem diameter of cauliflower. Similar results were observed by [5,23, 24] in cauliflower, [25] in broccoli. Nitrogen is an integral part of chlorophyll and a major constituent of all proteins. With the increase in nitrogen, the enzymatic reactions within a plant system operate with optimum efficiency which leads to an increase in vegetative growth, leaf area, leaf number, and chlorophyll content [26].

Nitrogen promotes vegetative growth; low or high doses of nitrogen fertilizer can delay curd initiation while an optimum dose of nitrogen fertilizer can promote early curd initiation of cauliflower. It is a well-known fact that nitrogen enhances vegetative growth and increases dry matter accumulation[27].Nitrogen increases the production of green leaves which help in the synthesis of carbohydrates, protein,and nitrogenous compounds essential for building up new tissues. Nitrogen also stimulates chlorophyll formation which in turn accelerates the rate of photosynthesis and eventually increases the growth and dry matter formation. It also promotes the growth and development of plants while increasing the uptake and utilization of potassium and phosphorus [28]. However, with the increase in nitrogen, curd initiation,and maturity are delayed due to an increase in the vegetative period [29]. The increased supply of nitrogen might have accelerated the synthesis of chlorophyll and amino acids and the effective utilization of carbohydrates and other organic compounds which might have resulted in the enlargement of curd. Nitrogen has a significant impact on the production of leaf area, the duration of leaf area, and the net assimilation rate, all of which are directly related to increased yield [30].Carbohydrate metabolism, enzyme activation, and translocation of sugars and starch alsoincrease with the supply of optimum level of nitrogen [31]. With the increase in yield attributing traits of plants (plant height, leaf number, leaf area, and plant spreading), the photosynthesis, net assimilation,and dry matter accumulation of plants increase. This result is supported by [3,32,33] in broccoliand [29] in cauliflower.

Comment [PS5]: Author also need to mention the following details:
"Simultaneously, the foliar sprayed nutrients (urea) increased the nutrient availability in the rhizo-ecosystem, resulting in improved plant growth and development (Dey et al., 2017). But the nutrient spray seems to be labor intensive and costly.

Dey, S., Prasad, S., Tiwari, P., & Sharma, P. (2017). Effect of urea, KCl, Zinc placement & spray on growth of cowpea. *Journal of Pharmacognosy and Phytochemistry*, 6(6S), 971-973.

The hybrid varieties have greater potential to uptake and utilize nitrogen more efficiently than the open pollinated varieties[30]. Similarly, hybrid varieties contain higher amounts of chlorophyll in their leaves, promoting the accumulation of photosynthates[34]. Higher photosynthetic activity correlated with the chlorophyll content of leaves [35]. The superiority of hybrid varieties might be due to improved photosynthetic activity and efficient transportation of photosynthates to developing sink tissues which ultimately increased plant biomass and yield [36]. The better growth and performance throughout the plant's development may therefore be influenced by the better accumulation of carbohydrate and pigment components in hybrid varieties[37]. Due to superior genetic makeup, hybrid varieties exhibited accelerated maturity, more complete apical dominance, higher harvest index, superior heading quality [38], and more productive and resistant to biotic and abiotic stresses [39]. Hybrid broccoli varieties outperformed open-pollinated varieties by 40-90%, owing to the hybrids' superior quality characteristics [40].

5. CONCLUSION

Nitrogen is a crucial nutrient for the growth and yield of cauliflower. This study indicated that increasing nitrogen levels significantly improved the yield and yield-attributing traits of cauliflower. Curd yield and growth parameters of cauliflower were highest at 130 kg N/ha. Similarly, the varietal response to nitrogen was superior in the Snow Mystique variety. Thus, the application of nitrogen and selection of varieties should be judiciously managed to obtain the maximum yield of cauliflower.

REFERENCES

1. MOALD. Statistical Information on Nepalese Agriculture (2020/21). Ministry of Agriculture & Livestock Development, Planning and Development Cooperation Coordination Division, Government of Nepal. Singhadurbar, Kathmandu, Nepal; 2022.
2. Bana ML, Kaushik RA, Bana RS, Dhakar MK. Integrated weed management in cauliflower cv. 'Pusa Snowball K-1'. *Annals of Agricultural Research* 2012;33(3):163-9.
3. Singh MK, Chand T, Kumar M, Singh KV, Lodhi SK, Singh VP, Sirohi VS. Response of Different Doses of NPK and Boron on Growth and Yield of Broccoli (*Brassica oleracea* L. var. *italica*). *International Journal of Bio-resource and Stress Management*. 2015;6(1):108-112.

Comment [PS6]: The manuscript is written well, however, the authors need to make some corrections as suggested in the manuscript. Moreover, in future studies, I suggested author to more focus on the scheduling of the nutrients rather than just on the different levels of crops which are already standardized.

4. Yeshiwas Y. Effect of different rate of nitrogen fertilizer on the growth and yield of cabbage (*Brassica oleraceae*) at Debre Markos, North West Ethiopia. African Journal of Plant Science. 2017;11(7):276-281.
5. Sharma V. Effect of nutrient management on growth and yield of cauliflower (*Brassica oleracea* var *botrytis*) inside low-cost polyhouse. Himachal Journal of Agricultural Research. 2016;42(1):88-92.
6. Barker AV, Pilbeam DJ, editors. Handbook of plant nutrition. CRC Press; 2015 May 14.
7. Sorensen J. Nitrogen effects on vegetable crop production and chemical composition. International Workshop on Ecological Aspects of Vegetable Fertilization in Integrated Crop Production. 1998; 506:41-50.
8. Sainju UM, Ghimire R, Pradhan GP. Nitrogen fertilization Impact on crop, soil, and environment. Nitrogen Fixation. 2019; 9:1-9.
9. Anas M, Liao F, Verma KK, Sarwar MA, Mahmood A, Chen ZL, Li Q, Zeng XP, Liu Y, Li YR. Fate of nitrogen in agriculture and environment: agronomic, eco-physiological and molecular approaches to improve nitrogen use efficiency. Biological Research. 2020;53(1):1-20.
10. Johnson A, Laney K. National Academies of Sciences, Engineering, and Medicine, Environmental Health Matters Initiative. Reducing the Health Impacts of the Nitrogen Problem; 2021 January 28 - February 25; Washington (DC): National Academies Press;2021.
11. Lee SK, Kader AA. Preharvest and postharvest factors influencing vitamin C content of horticultural crops. Postharvest biology and technology. 2000;20(3):207-20.
12. Lisiewska Z, Kmiecik W. Effects of level of nitrogen fertilizer, processing conditions and period of storage of frozen broccoli and cauliflower on vitamin C retention. Food Chemistry. 1996;57(2):267-70.
13. Department of Hydrology and Meteorology. Meteorological information. Ministry of Energy, Water Resources and Irrigation. Kathmandu, Nepal; 2019. Available: <https://www.dhm.gov.np/>
14. Shi P, Liu M, Yu X, Gielis J, Ratkowsky DA. The proportional relationship between leaf area and the product of leaf length and width of four types of special leaf shapes. Forests. 2019;10(2):178.
15. Neethu TM, Tripathi SM, Narwade AV, Sreeganesh S. Effect of N and P levels on growth and yield parameters of broccoli (*Brassica oleracea* L. var. *italica*) under South Gujarat soil conditions. International Journal of Tropical Agriculture. 2015;33(Part II):913-917.
16. Kumar P, Sharma SK. Performance of different sprouting broccoli [*Brassica oleracea* var. *italica*] cultivars under different combinations of nitrogen, phosphorus and potassium. Indian Journal of Agricultural Sciences. 2001;71(8):558-560.
17. Kanaujia SP, Singh PK. Integrated nutrient management for quality production of cauliflower in acid alfisol of Nagaland. Karnataka Journal of Agricultural Sciences. 2015;28(2):244-247.
18. Prasad PH, Bhunia P, Naik A, Thapa U. Response of nitrogen and phosphorus levels on the growth and yield of chinese cabbage (*Brassica campestris* L. var. *pekinensis*) in the gangetic plains of West Bengal. Journal of Crop and Weed. 2009;5(2):75-77.
19. Dheerendra K, Tripathi SM, Dwivedi AK, Vivek P. Effect of nitrogen and phosphorus on crop growth, head yield and quality of broccoli (*Brassica oleracea* L. var. *italica*). HortFlora Research Spectrum. 2012;1(2):172-174.
20. Van der Werf A, Nagel OW. Carbon allocation to shoots and roots in relation to nitrogen supply is mediated by cytokinins and sucrose: opinion. Plant and Soil. 1996; 185:21-32.

21. Agarkar UR, Dadmal KD, Nikas NS, Piwlatkar GK. Effect of nitrogen levels and spacing on growth and yield of broccoli (*Brassica oleracea* var. *italica* L.). *Green farming*. 2009;1(5):477-9.
22. Tripathi S, Patel J M, Singh N, Naik J, Naik V R. Effect of different NPK levels on growth and yield attributes of broccoli (*Brassica oleracea* L) under South Gujarat condition. *International Journal of Chemical Studies*. 2020;8(3):1335-1339.
23. Meena Y, Sharma R, Kushwah S, Gallani R. Effect of varieties and nutrient levels on growth and yield of cauliflower (*Brassica oleracea* var. *Botrytis* L.). *The Bioscan*. 2017; 12(3):1731-1734.
24. Tekasangla S, Kanaujia P, Singh PK. Integrated nutrient management for quality production of cauliflower in acid alfisol of Nagaland. *Karnataka Journal of Agricultural Science*. 2015;28(2):244-247.
25. Bika R, Bhandari N, Khanal A. Response of different doses of nitrogen on broccoli (*Brassica oleracea* var. *italica*) in Lamjung district. *International Journal of Applied Sciences and Biotechnology*. 2018;6(3):270-273.
26. Shah DA, Narayan R, Ahmad N, Narayan S, Wani KP. Influence of boron and zinc on growth yield and quality of knolkhol cv. Early White Veinna. *Indian Journal of Horticulture*. 2010;67(4):323-328.
27. Yadav HK, Rawat TS, Dogra P, Yadav V. Effect of foliar application of nitrogen and zinc on quality of cauliflower (*Brassica oleracea* var. *Botrytis* L.) cv. Snowball-16. *Progressive Horticulture*. 2015;47(2):338-340.
28. Alhrout HH. Response of growth and yield components of sweet pepper to tow different kinds of fertilizers under greenhouse conditions in Jordan. *Journal of Agricultural Science*. 2017;9(10):265-276.
29. Bashyal LN. Response of cauliflower to nitrogen fixing biofertilizer and graded levels of nitrogen. *Journal of Agriculture and Environment*. 2011; 12:41-50.
30. Leghari SJ, Wahocho NA, Laghari GM, Laghari AH, Bhabhan GM, Talpur KH, Bhutto TA, Wahocho SA, Lashari AA. Role of nitrogen for plant growth and development: A review. *Advances in Environmental Biology*. 2016;10(9):209-219.
31. Singh AK, Akhilesh S. Influence of nitrogen and potassium on growth and head yield of broccoli (*Brassica oleracea* L. var. *italica*) under low hills subtropical condition of HP. *Vegetable Science*. 2000;27(1):99-100.
32. Moniruzzaman M, Rahman SM, Kibria MG, Rahman MA, Hossain MM. Effect of boron and nitrogen on yield and hollowstem of broccoli. *Journal of Soil and Nature*. 2007;1(3):24-29.
33. Dhakal M, Shakya SM, Bhattarai S. Yield and quality of broccoli (*Brassica oleracea* L. var. *italic* Plenck.) cv. Calabrese affected by nitrogen and farm yard manure in Chitwan, Nepal. *Journal of Plant Health*. 2016;1(1):102.
34. Tang WB, Zhang GL, Xiao YH, Deng HB, Ke FA, Liu GH, Chen LY. Physiological and biochemical characteristics in flag leaves of the *C liangyou* series of hybrid rice combinations at late growth stages. *Rice Science*. 2010;17(4):319-25.
35. Ni Z, Kim ED, Ha M, Lackey E, Liu J, Zhang Y, Sun Q, Chen ZJ. Altered circadian rhythms regulate growth vigour in hybrids and allopolyploids. *Nature*. 2009;457(7227):327-31.
36. Sreeharsha RV, Mudalkar S, Sengupta D, Unnikrishnan DK, Reddy AR. Mitigation of drought-induced oxidative damage by enhanced carbon assimilation and an efficient antioxidative metabolism under high CO₂ environment in pigeon pea (*Cajanus cajan* L.). *Photosynthesis Research*. 2019; 139:425-39.

37. Elahi E, Wali A, Nasrullah GA, Ahmed S, Huma Z, Ahmed N. Response of cauliflower (*Brassica oleracea* L. botrytis) cultivars to Phosphorus levels. *Pure and Applied Biology (PAB)*. 2021;4(2):187-94.
38. Stansell Z, Björkman T. From landrace to modern hybrid broccoli: the genomic and morphological domestication syndrome within a diverse *B. oleracea* collection. *Horticulture research*. 2020; 7:159-176.
39. Ficiyan AM, Loos J, Tschardtke T. Similar Yield Benefits of Hybrid, Conventional, and Organic Tomato and Sweet Pepper Varieties Under Well-Watered and Drought-Stressed Conditions. *Frontiers in Sustainable Food Systems*. 2021; 5:628537.
40. Wehner TC. Heterosis in vegetable crops. *Genetics and Exploitation of Heterosis in Crops*. 1999:387-97.

UNDER PEER REVIEW