

## Original Research Article

### Surveillance of Mycotoxigenic Fungi and Production of Aflatoxin by *A. flavus* in Contaminated Maize Seeds- Bihar.

#### **Abstract**

Aflatoxin- producing fungi like *Aspergillus flavus* contaminate maize crops in the agricultural field at harvest, post- harvest and during storage making them one of the most widespread and dangerous mycotoxins. It has been directly correlated to adverse the health effects, such as liver cancer in many animal species as well as plant system.

Maize samples collected from seven districts of Bihar viz, Begusari, East champaran, Khagaria, Bhagalpur, Saharsa, Madhepura and Samastipur. Collected sample examined for associated mycoflora and aflatoxin producing prospective of *Aspergillus flavus*. It was determined that *A. flavus* was of presiding occurrence accompanied by *Penicillium spp.*, *Fusarium*, *A. nigar*, *Rhizopus*, *Trichoderma*, *Mucor* and *A. ochraceus*. 119 strains of the total *A. flavus* isolates were toxigenic, producing aflatoxin B<sub>1</sub>, B<sub>2</sub> and G<sub>1</sub>. Highest Aflatoxin is all probability due to complementary environmental conditions, undeveloped agricultural practices, poor storage circumstances of grains and because of yearly flood problems in this region.

**Key words:** Maize seeds, Aflatoxin, Mycoflora, Seven districts of Bihar.

#### **INTRODUCTION**

Mycotoxin (Greek word Mykes = Fungus and Toxin = Toxin) is toxic secondary metabolite produced by organism of Kingdom Fungi (Richard JL, 2007; Cimen *et. al.*, 2022) and is capable of causing disease and death in humans and animals (Bennett, J. W and Klick, M. 2003). They are stable, invisible and toxic chemical compounds, found to be common in the farm environment surviving in several places as well as on many different types of feed stuffs.

The growing moulds and production of mycotoxin are most encouraged by environmental factors like temperature (cool / hot) and moisture (wet / dry). However, mycotoxins may also be produced by moulds when other stress conditions occur to the host plant or the mould.

Aflatoxins are a type of mycotoxin produced by *Aspergillus spp.* such as *A. flavus* and *A. parasiticus* (Martins *et. al.*, 2001). The umbrella term aflatoxin refers to four different compounds which are B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub> and G<sub>2</sub> (Yin YN *et. al.*, 2008) where Aflatoxin B<sub>1</sub>, the most toxic, is a potent carcinogen and has been directly correlated to adverse health effects, such as Liver cancer in many animal species (Martins *et. al.*, 2001). Aflatoxin is largely associated with

commodities produced in the tropics and subtropics region in cotton, wheat, millet, spices, rice, sorghum, peanuts, sunflower, pistachios and maize crops (Martins *et. al.*, 2001; Yin YN *et. al.*, 2008).

Maize is widely cultivated throughout the world and a greater weight of maize is produced each year than any other grain (International grains council, 2013). In 2021, total world production was 1.2 billion tonnes. Maize is the most widely grown grain crop throughout the America. Out of 38 administrative districts of Bihar only seven districts, viz. Begusarai, East champaran, Khagaria, Bhagalpur, Saharsa, Madhepura and Samastipur constitute about half of the total maize acreage. These districts are historically flood-prone areas during rainy season and fall north of the Ganges, having several seasonal river tributaries.

These seven districts of Bihar have recorded in April 2021, the highest maize productivity in the world. Maize cultivation provides livelihood to approx. 1.3 million farmers of Bihar state. Maize has become a staple food in many parts of the world with the total production surpassing that of wheat or rice. In addition to being consumed directly by humans (often in the form of masal). Maize is used for corn ethanol, animal feed and other products, such as corn starch and corn syrup (Foley, J. 2019). It is also used in making ethanol and other biofuels.

Therefore, an attempt has been made to surveillance of mycotoxigenic fungi and production of aflatoxin by *A. flavus* in contaminated maize seeds of major growing districts of Bihar.

### **Occurrence of mycotoxin in maize crop**

Aflatoxin – producing fungi like *A. flavus* contaminate maize crops in the agricultural field at harvest and during storage making them one of the most widespread and dangerous mycotoxins. In Kenya (2004), about 125 people died and nearly approx. 200 others required medical treatment after eating aflatoxin contaminated maize (Lewis L. *et. al.*, 2005). The death was associated with home-grown maize that had not been treated with fungicides or properly dried before storage. At that time, due to food shortages, farmers may have been harvesting maize earlier than normal to prevent thefts from their agricultural field, so that the grain had not fully matured and was more susceptible to infection with *A. flavus*.

## **MATERIALS AND METHODS**

### **Sample collection**

Maize seed samples were collected from farmers field in seven districts viz. Begusarai, East Champaran, Khagaria, Bhagalpur, Saharsa, Madhepura and Samastipur of Bihar. Maize seed samples kept in sterile polythene bags for further experiments. Moisture content of each sample was recorded with the help of OSAW moisture meter and PH reading were taken by using digital PH meter (IJBAF, 2013).

### **Isolation and identification of Mycoflora**

100 kernels of all seven- sampling site (districts) was surface sterilized in 2% NaOCl and plated on moist blotting paper in sterile petri dishes (ISTA, 1966). Plates were incubated at room temperature for 7 days followed by macro and microscopic identification (Mc Clenny, N. 2005; Adame- Garcia, J. *et. al.*, 2015). Fungal colonies were maintained on PDA media for further use and identification.

### **Aflatoxin analysis of toxigenic isolates**

*Aspergillus flavus* isolates were allowed to grow on SMKY liquid media (Diener and Davis, 1966) for one week at  $30 \pm 2^\circ\text{C}$  and aflatoxin extracted (Thomas *et. al.*, 1975). It was extracted with chloroform, dried on water bath and concentrated extracts were kept in screw tight bottle for qualitative and quantitative analysis.

### **Qualitative and Quantitative analysis**

Qualitative analysis of aflatoxin was done using TLC (Thin layer chromatography), toluene-isoamyl alcohol- methanol (90: 32: 2 v/v) solvent system (Reddy *et. al.*, 1970) and for chemical conformation trifluoroacetic acid (Stack and Pohland, 1975) or 25% sulphuric acid spray was used.

Quantitative estimation of aflatoxin was done by spectrophotometrically (Nabney and Nesbitt, 1965).

## **RESULTS**

Table -1 and fig. A, indicates the association of mycoflora in seven districts of Bihar in flood – prone area during rainy season (Agroclimaic zone in Fig- B) and observed during the study, moisture content and PH of the sample. *A. flavus*, *Penicillium* spp., *Fusarium*, *A. nigar* and *Rhizopus* were present in all the seven districts in large amount whereas *Trichoderma*, *Mucor* and *A. ochraceus* present in rare amount of all districts but absent in Khagaria and Saharsa districts, respectively (Fig- C). PH ranged between 5.8 to 6.7 and moisture content were recorded 10.8, 10.5, 10.0, 9.6, 10.1, 10.4 and 10.7 for Begusarai, East Champaran, Khagaria, Bhagalpur, Saharsa, Madhepura and Samastipur, respectively.

Table-2 represent that *A. flavus* sample of 7 districts having 385 isolates of *A. flavus*. 119 strains were found to be toxigenic in which 86 positives to AFT- B<sub>1</sub>, 24 positives to AFT-B<sub>1</sub>B<sub>2</sub> and only 9 positives for AFT-B<sub>1</sub>B<sub>2</sub>G<sub>1</sub>. The amount of AFT-B<sub>1</sub> was, however very low at 0.2- 1.3, 0.0- 1.0 and 0.0- 1.1 µg/ml, respectively. Whenever, the maximum aflatoxin was recorded from

Begusarai samples at 0.2- 1.8 µg/ml followed by East champaran, Khagaria, Saharsa, Bhagalpur, Samastipur and Madhepura district, respectively.

Maize samples collected from seven districts in flood- prone area during rainy season were highly contaminated with *A. flavus* and other genera of fungi like, *Penicillium*, *Fusarium*, *A. nigar*, *Rhizopus*, *Trichoderma*, *Mucor* and *A. ochraceus*. Mycotoxin producing fungi like *Aspergillus*, *Fusarium* spp. were of predominant occurrence (Fig. D and E). The moisture contents were also observed in all the samples which influence the aflatoxin production.

## **DISCUSSION**

Recently, about 125 people died and nearly approx. 200 others required medical treatment after eating aflatoxin contaminated maize (Lewis L. *et. al.*, 2005). In Kenya (2004), the death was associated with homegrown maize that had not been treated with fungicides or properly dried before storage. At that time, due to food shortages, farmers may have been harvesting maize earlier than normal to prevent thefts from their agricultural field, so that the grain had not fully matured and was more susceptible to infection with *A. flavus* and aflatoxin producing potentiality of toxigenic strains of *A. flavus* were higher, due to moisture content in rainy season in flooded area as well as poor storage conditions that provide an opportunity for fungal growth like Aflatoxin to easily invade the maize seeds.

Bihar has experienced serious aflatoxicosis outbreaks associated with maize which has claimed lives as well as maize yield losses. Our findings from the various sites (districts) revealed that, potentially mycotoxigenic fungal isolates were found on maize samples. Maize samples from each district were more infested by a specific fungal genus. Maize grains from Begusarai were heavily contaminated by *Aspergillus* spp. while those collected from Khagaria and Saharsa were not contaminated with *Trichoderma* and *Mucor*.

## **CONCLUSION**

Maize samples from the seven districts tested were infested by different mycotoxigenic fungi. The existence of mould on the maize samples shows the possibility of occurrence of more than one mycotoxin but dominant by Aflatoxin B<sub>1</sub>. This may be maize has the potential to enhance the income of 1.3 million maize grower in Bihar State, thereby significantly reducing poverty in us of the poorest states in India.

So, it was important to determine the distribution and incidence of fungi that exist in maize from different districts.

## **REFERENCES**

1. Richard, JL (2007). "Some major mycotoxins and their mycotoxicosis- an overview". *Int. J. Food microbiol.* 119 (1-2): 3-10.
2. Cimen, Duygu, Bereli, Nilay, Denizli, Adil (2022). "Patulin Imprinted Nanoparticles Decorated surface Plasmon Resonance chips for Patulin Detection". *Photonic Sensors.* 12 (2): 117-129.
3. Bennett, J.W; Klich, M (2003). Mycotoxins. *Clinical microbiology Reviews.* 16 (3). 497-516.
4. Lewis, L. Onsongo, M. Njapau, H. *et. al.*, (2005). "Aflatoxin contamination of commercial maize products during an outbreak of acute aflatoxicosis in eastern and central Kenya. *Environ. Health Perspect.* 113 (12): 1763-67.
5. Liu. Yue, Yamdeu, Joseph Hubert Golani, Gong, Yun Yun, orfila, Caroline (2020). "A review of postharvest approaches to reduce fungal and mycotoxin contamination of foods". *Comprehensive Reviews in Food Science and Food safety.* 19 (4): 1521-1560.
6. Martins, ML. Martins, HM, Bernardo, F. (2001). "Aflatoxins in spices marketed in Portugal". *Food Addit. Contam.* 18 (4): 315-19.
7. Mc Clenny N. (2005). Laboratory detection and identification of *Aspergillus* traditional approach. *Medical Mycology* 43 (Supplement- I): S125- S128.
8. Miller JD. (2008). Mycotoxins in small grains and maize: old problems, new challenges. *Food Additives and Contaminants Part A* 25: 219-230.
9. Adame- Garcia, J. Rodriguez- Guerra, R. Iglesias- Andreu, LG. Ramos- Prado, JM. Luna- Rodriguez, M. (2015). Molecular identification and Pathogenic variation of *Fusarium* species isolated from vanilla planifolia in Papantla Mexico. *Botanical Sciences* 93 (3): 669-678.
10. Thomas, F. Eppley, RM, Trucksess, MW. (1975). Rapid screening method for aflatoxins and zearalenone in corn. *Journal of the Association of official Analytical chemists.* 58 (1): 114-116.
11. Yin, YN. Yan, LY. Jiang, JH. Ma, ZH (2008). "Biological control of aflatoxin contamination of crops". *J. Zhejiang univ. sci. B.* 9 (10): 787-92.
12. Foley Jonathon (2019). "Its Time to Rethink Americas corn sys. *Scientific American.* Retrieved. Feb 15, 2019.
13. International grains council (International organization 2013). International grains council market Report 28 Nov 2013.
14. IJBAF (2013). Mycoflora and aflatoxin contamination of some foodstuffs. \**ORANUSI SU AND OLAREWAJU SA.* 1 (1): 9-18.
15. Diener, U.L. and Davis, N. D. (1966). Aflatoxin production by isolates of *Aspergillus flavus*. *Phytopathology*, 56, 1390-1393.
16. Nabney, J. and Nesbitt, B. F. (1965). A spectrophotometric method for determining the aflatoxins Analysis. 90: 155-160.
17. Reddy, T.V. Viswanathan, L. and Venkita subramania, J.A. (1970). Thin layer chromatography of aflatoxins. *Analytical Biochemistry*, 38: 568-571.
18. Stack, M.E. and Pohland, A.E. (1975). Collaborative study of a method for chemical conformation of identity of aflatoxin. *J. Assoc. off. Anal. Chem.* 58:110-113.


UNDER PEER REVIEW

**Table -1: Mycoflora associated with Maize Seeds collected from seven Districts of Bihar**


S l . N o .	Fungus	Maize Seed Samples from Seven District																								
		Begusari (N =100)			East Champaran (N =100)			Khagaria (N =100)			Bhagalpur (N =100)			Saharsa (N =100)			Madhepura (N =100)			Samstipur (N =100)						
		No. of seed infested	Moisture content	PH	No. of seed infested	Moisture content	PH	No. of seed infested	Moisture content	PH	No. of seed infested	Moisture content	PH	No. of seed infested	Moisture content	PH	No. of seed infested	Moisture content	PH	No. of seed infested	Moisture content	PH				
1	<i>A. Flavus</i>	75			66				63				57				46				41			37		
2	<i>Penicillium Spp.</i>	68			60				57				49				42				38			33		
3	<i>Fusarium Spp.</i>	62			58				43				34				30				28			25		
4	<i>A. nigar</i>	74	10.8	6.7	65	10.5	6.3	62	10.0	6.0	56	9.6	5.8	45	10.1	5.9	40	10.4	6.2	36	10.7	6.5	36	10.7	6.5	
5	<i>Rhizopus Spp.</i>	35			32				24				21				19				16			13		
6	<i>Trichoderma</i>	9			5				0				3				6				9			4		
7	<i>Mucor</i>	4			9				1				7				0				1			5		
8	<i>A. ochraceus</i>	5			8				0				4				7				2			1		

**Table- 2: *Aspergillus flavus* isolates from Maize Seed samples**

Sl. No.	Districts of Bihar	No. of <i>A. flavus</i> strains isolates	No. of toxigenic isolates of <i>A. flavus</i>	Positive isolates			Range of aflatoxin B <sub>1</sub> concentration µg/ml (ppm)
				B <sub>1</sub>	B <sub>1</sub> B <sub>2</sub>	B <sub>1</sub> B <sub>2</sub> G <sub>1</sub>	
1	Begusari	75	25	15	7	3	0.2- 1.8
2	East Champaran	66	19	13	5	1	0.1- 1.6
3	Khagaria	63	18	12	4	2	0.2- 1.4
4	Bhagalpur	57	13	11	1	1	0.2- 1.3
5	Saharsa	46	17	14	2	1	0.0- 1.2
6	Madhepura	41	12	10	2	0	0.0- 1.0
7	Samstipur	37	15	11	3	1	0.0- 1.1
	<b>Total</b>	<b>385</b>	<b>119</b>	<b>86</b>	<b>24</b>	<b>9</b>	-



**Fig- A:** Line showing No. of Maize Seed infested in Seven Districts of Bihar




**Fig B:** Pink dot indicates Agroclimatic Zone of Seven Districts of Bihar



Mycoflora Grown on PDA media  
on various Districts - Bihar.

**Fig- C:** Association of Mycoflora in PDA media of Seven Districts of Bihar



**Fig- D:** Pie chart showing No. of toxigenic isolates of *A. flavus* (%) in Seven Distri Bihar



**Fig- E:** Association of Mycoflora in maize comb during flood situation