

## Original Research Article

### **EVALUATION OF DETRUSOR CONTRACTILITY PARAMETERS HELP TO ANALYZE CONSEQUENCES OF STEADY ABDOMINAL PRESSURE CHANGES DURING VOIDING IN WOMEN.**

**Comment [CO1]:** The topic does not reflect the objectives of the study

#### **Abstract**

#### **Introduction**

Urodynamic study interpretation leading to urodynamic diagnosis is mainly based on pressure recordings and the value of detrusor pressure at maximum flow ( $p_{det.Qmax}$ ). Detrusor pressure is calculated by subtracting the abdominal pressure ( $p_{abd}$ ) from the vesical pressure ( $p_{ves}$ ). Hence, there is a critical role for  $p_{abd}$  in this process. The goal of our study was to evaluate the contribution of detrusor contractility parameters (DCP) to confirm urodynamic misdiagnosis (UmD) due to steady abdominal pressure changes during voiding in women.

#### **Material – Methods**

Urodynamic tracings of 271 non-neurologic women referred for investigation of various LUTS were retrospectively analyzed. UmD could be bladder outlet obstruction (BOO) and normal (N) for decreased  $p_{abd}$ , normal (N) and detrusor underactivity (DU) for increased  $p_{abd}$ . Detrusor contractility parameters were VBN parameter  $k$  and PIP1.

#### **Results**

Among the whole population 125 women had a significant change ( $\geq 5$  cmH<sub>2</sub>O) of  $p_{abd}$  during voiding (73 decrease, 52 increase).

In the “decrease” sub-group, only 3 N became DU with decreased DCP values; in the “increase” sub-group 1 N and 1 DU patients gained BOO diagnosis with increased DCP values.

In total, analysis of changes in abdominal pressure leads to 5/271 (1.8%) changes in urodynamic diagnosis; no correlation between previous surgery of incontinence or main complaint.

### **Conclusion**

A high percentage of the non-neurologic female population has steady changes of abdominal pressure during voiding, that condition leads to few changes in urodynamic diagnosis; evaluation of detrusor contractility parameters values help to verify the new conclusions.

**Key words:** abdominal pressure; detrusor contractility; urodynamic diagnosis; voiding; non-neurologic women.

## Introduction

After completion of a voiding cystometry, the study interpretation leading to a urodynamic diagnosis (UD) is mainly based on pressure recordings and the value of detrusor pressure at maximum flow ( $p_{det.Qmax}$ ). Detrusor pressure is calculated by subtracting the abdominal pressure  $p_{abd}$  (assumed equal to rectal) from the vesical pressure ( $p_{ves}$ ). Hence, there is a critical role for abdominal pressure in this process.

According to the report of Good Urodynamic Practice guidelines [1], abdominal pressure is recorded using a punctured intrarectal balloon catheter filled with 2 mL of saline. During voiding, the changes in abdominal pressure can be threefold, decreased, unchanged, or increased (generally from straining efforts). Two of these three processes, increase or decrease in abdominal pressure, can have an effect on the final urodynamic diagnosis, especially for the diagnoses of detrusor underactivity (DU) or bladder outlet obstruction (BOO) which rely primarily on  $p_{det.Qmax}$  [2-3-4].

The goal of this study was to analyze the consequences of steady changes in abdominal pressure during voiding in women on urodynamic diagnosis, and the contribution of analysis of detrusor contractility parameters (DCP) to confirm urodynamic misdiagnosis (UmD).

## Material and Methods

Urodynamic tracings of 271 non-neurologic women age range [20-88 years old] who were referred for investigation of various lower urinary tract symptoms (LUTS) to our specialized unit, run by the same team over time were

retrospectively analyzed. Exclusion criteria, in addition of neurological condition, were advanced cognitive impairment (MMSE  $\leq$  20), diabetes mellitus, grade  $\geq$  2 pelvic organ prolapse, complete urinary retention and/or severe mobility impairment. Each patient file comprised demographic data, medical history, 3-day bladder diary, and current medications. Main complaint was categorized as stress urinary incontinence (SUI), urge urinary incontinence (UUI) mixed urinary incontinence (MUI), and "OTHER" (dysuria-frequency complaint, meaning LUTS but no urinary incontinence). Each analyzed file included a filling cystometry followed by a voiding study with an intubated flow (IF). Cystometry was performed with the patient in a sitting position with a 7-F triple-lumen urethral catheter perfused with saline at room temperature using a medium filling rate of 50 mL/min. Abdominal pressure was recorded using a punctured intrarectal balloon catheter filled with 2 mL of saline according to the report of Good Urodynamic Practice guidelines [1]. Post-void residuals (PVR) were measured using bladder-scan after the IF.

After completion of the urodynamic session, the tracings were interpreted to reach a urodynamic diagnosis which conformed to the ICS/IUGA recommendations [5]. UD included: bladder outlet obstruction (BOO), detrusor hyperactivity with impaired contractility (DHIC), detrusor overactivity (DO), detrusor underactivity (DU). Some investigations were found "normal" (N) and others related to urethral dysfunction (intrinsic sphincter deficiency (ISD) and voiding triggered by urethral relaxation (URA)). Some combined diagnoses were observed between DO, DU or DHIC with ISD.

Diagnoses of BOO and DU are based on the following criteria:

1- for BOO:  $p_{\text{det.Qmax}} \geq 25 \text{ cm H}_2\text{O}$  and  $Q_{\text{max}} \leq 12 \text{ mL}\cdot\text{s}^{-1}$  proposed by Defreitas et al. [4].

2- for DU the cutoff criteria, usable for all women, proposed by Gammie et al. [2] were used:  $p_{\text{det.Qmax}} < 20 \text{ cm H}_2\text{O}$ ,  $Q_{\text{max}} < 15 \text{ mL}\cdot\text{s}^{-1}$  and BVE (bladder voiding efficiency)  $< 90\%$ .

To add more consistency, an evaluation of detrusor contractility was obtained from the VBN detrusor contractility parameter [6]  $k$  and the projected isovolumetric pressure PIP1 [3] ( $\text{PIP1} = p_{\text{det.Qmax}} + Q_{\text{max}}$ ). Parameters necessary for  $k$  computation include initial bladder volume  $V_{\text{ini}}$  (voided volume + post void residual) with voided volume  $\geq 100 \text{ mL}$ , intubated maximum flow ( $Q_{\text{max}}$ ) and detrusor pressure at maximum flow ( $p_{\text{det.Qmax}}$ ) [7]. It had been demonstrated that  $k$  and PIP1 gave consistent evaluations of detrusor contractility for females [8].

After evaluation of  $p_{\text{abd}}$  at maximum flow, a correction of  $p_{\text{det.Qmax}}$  erasing the artificially increase of  $p_{\text{det.Qmax}}$  was applied; when  $p_{\text{abd}}$  decreased during voiding; a similar scheme for correction of  $p_{\text{det.Qmax}}$  was used.

Then, after evaluation of the real value of  $p_{\text{det.Qmax}}$ , we investigated possible changes in urodynamic diagnosis and, to check the validity of the proposed changes, we computed values of detrusor contractility parameters.

This study was conducted in accordance with the Declaration of Helsinki. According to the local practice of Ethics Committee, there is no formal Institutional Review Board approval required for retrospective studies.

### **Statistical analysis**

Data are presented as mean  $\pm$  SD and range. Analysis of variance (ANOVA), and the Chi-square test were used as appropriate. All statistical results were considered significant at  $p < 0.05$ . Statistical analyses were performed using SAS, version 5.0 (SAS Institute, Inc., Cary, NC).

## Results (Tables 1-2-3)

### 1- Steady decrease of abdominal pressure during voiding (Figure1) (Tables 1-2)

Among the whole population, 145 women (53.5%) had a steady decrease of abdominal pressure from baseline to the time of  $Q_{max}$  during the intubated flow among which 73 (26.9%) had a significant decrease ( $\geq 5$  cmH<sub>2</sub>O).

After new evaluation of  $p_{det;Q_{max}}$  taking into account the decrease of  $p_{abd;Q_{max}}$ :

- all BOO remained BOO
- among 51 N, 3 had criteria following Gammie's criteria [2] ( $p_{det}$ ,  $Q_{max}$  and BVE) and then gained DU diagnosis.

For these patients with initial UD "normal" the complaint and the decrease of values of contractility characteristics ( $k$  and PIP1) were as follow:

- complaint UUI, age 75y, from  $k = .326$ , PIP1 = 33 to  $k = .070$ , PIP1 = 21
- complaint MUI, age 72y, from  $k = .257$ , PIP1 = 30 to  $k = .153$ , PIP1 = 25
- complaint OTHER, age 68y, from  $k = .253$ , PIP1 = 40 to  $k = .030$ , PIP1 = 22

Comment [CO2]: State the outcome

The decrease of  $p_{abd}$  for these three patients was respectively -12, -5 and -18 cm H<sub>2</sub>O.

Change of UD was 3/271 (1.11%) for the whole population and 3/145 (2.07%) for the patients with steady decreasing abdominal pressure during voiding.

## 2- No change of detrusor pressure during voiding (Tables 1-2)

Forty five (35.7%) patients had no change of detrusor pressure during voiding and thus no change in urodynamic diagnosis.

## 3- Steady increase of abdominal pressure during voiding (Figure 1) (Tables 1-2)

Among the whole population, 81 women (53.5%) had an steady increase of abdominal pressure from baseline to the time of  $Q_{max}$  during the intubated flow and 52 women (19.2%) had a significant increase ( $\geq 5$  cmH<sub>2</sub>O) in  $p_{abd}$  from baseline to the time of  $Q_{max}$  resulting in artificially decrease in  $p_{det.Qmax}$ .

After correction of abdominal pressure and withdrawal of patients who strained, among 5 patients with N diagnosis, 1 gained BOO diagnosis and among 3 patients with DU diagnosis, 1 gained BOO diagnosis.

For these 2 patients the complaint and the changes of values of contractility characteristics ( $k$  and PIP1) were as follow:

- patient with initial UD "normal": complaint MUI, age 69 y, from  $k=.410$ , PIP1 = 38 to  $k=.565$ , PIP1=45

- patient with initial UD "detrusor underactivity" :complaint SUI, age 37y, from  $k=.103$ , PIP1 =21 to  $k=.472$ , PIP1=41

Their correction of  $p_{det}$  was respectively +7 and +20 cm H<sub>2</sub>O.


**Comment [CO3]:** The calculation is not necessary

In total, among 81 patients with steady increase of abdominal pressure during voiding 2 (0.6%) had change in urodynamic diagnosis and mainly gain of BOO diagnosis.

#### 4- Influence of previous surgery of incontinence


Thirty eight women had surgery for urinary incontinence: 31 TVT or TOT, 1 Burch while 6 had TVT ablation.

Among these women 9 had significant decrease and 9 significant increase of  $p_{det}$  during voiding. None had change in urodynamic diagnosis after correction of  $p_{det.Qmax}$ .




**Figure 1:** Steady changes of  $p_{abd}$  from baseline during voiding and correction which must be made for  $p_{det}$ .

In total, analysis of changes in abdominal pressure led to 5/271 (1.8%) changes in urodynamic diagnosis. There were no correlation between previous surgery of incontinence and changes in urodynamic diagnosis due to changes in abdominal pressure during the voiding phase. Changes in values of detrusor contractility parameters were consistent with usual values of the new urodynamic diagnoses [8] (Figures 2-3).



**Figure 2:** VBN detrusor contractility  $k$  vs. age



**Figure 3 :** Values of detrusorcontractilityparameters ( $k$  without unit and PIP1 in cm H<sub>2</sub>O) vs urodynamic diagnoses : BOO bladder outlet obstruction; DHIC detrusor hyperactivity with impaired contractility, DO detrusor overactivity, DU detrusor underactivity; N investigations found "normal"; investigations related to urethral dysfunction: intrinsic sphincter deficiency (ISD) and voiding triggered by urethral relaxation (URA).

### Interpretation of the results

Changes of abdominal pressure during voiding are very frequent in women whether it be decrease or increase. Small fluctuations are frequently due to live signals and are rubbed out with acute examination of the traces. This study is interested in the permanent and regular variations during voiding. Muscular relaxation causes decrease of abdominal pressure with consequence a rise of detrusor pressure while rectal contraction causes increase in abdominal

pressure with consequence a decrease in detrusor pressure. It is the first study which assesses the exact variations in abdominal pressure during voiding between baseline and the time of maximum flow and, then evaluates the consequences on urodynamic diagnoses. In their study, Valdevenito et al. [9-10] only evaluate the changes due to a decrease of abdominal pressure.

**Comment [CO4]:** This statement should be modified or deleted. Proper literature search will yield similar studies.

Some decreases and increases are of low amplitude consequently without effect. Due to accuracy of static pressure measurement in urodynamic system [11], a gap of +5 to -5 cm H<sub>2</sub>O between the beginning of voiding and the time of Q<sub>max</sub> is considered as irrelevant. Note that for URA urodynamic diagnosis, after the initial urethral relaxation which triggers voiding, the three options (decrease, statu quo and increase) for evolution of abdominal pressure during voiding can be observed.

In our non neurologic female population, 145 (53.5%) have a decrease of abdominal pressure without association with one complaint and 81 (29.9%) an increase. However, 50.3% of the population with decrease and 64.2% of the population with increase doesn't need detrusor pressure correction; more care should be taken in the population with increase because there are possible straining episodes.

Whereas the different urodynamic diagnoses, two depend of the evaluation of detrusor pressure at maximum flow: bladder outlet obstruction and detrusor underactivity. So, between the urodynamic diagnosis proposed in the initial analysis of the urodynamic testing bladder outlet obstruction and normal urodynamic can be revised with a correction of abdominal pressure in case of abdominal pressure decrease during voiding while normal urodynamic and

**Comment [CO5]:** Unclear statement.

detrusor underactivity can be revised in case of abdominal pressure increase during voiding. PIP1 and  $k$  values are of the order of magnitude expected for age and urodynamic diagnosis whether it be a steady decrease or an steady increase of abdominal pressure during the voiding phase<sup>7</sup>.

### **Conclusion**

A high percentage of the non-neurologic female population has steady changes of abdominal pressure during voiding the consequence of which may be a urodynamic misdiagnosis. If the correction of these changes leads to few changes, an evaluation of detrusor contractility parameters values helps to verify the new findings.

### **References**

- 1- Rosier PFWM, Schaefer W, Lose G, Goldman HB, Guralnick M, Eustice S et al. International Continence Society Good Urodynamic Practices and Terms 2016: Urodynamics, uroflowmetry, cystometry, and pressure-flow study. *Neurourol Urodyn.* 2017; 36(5):1243-1260. doi: 10.1002/nau.23124. Epub 2016 Dec 5.
- 2- Gammie A, Kaper M, Dorrepaal C, Kos T, Abrams P. Signs and Symptoms of Detrusor Underactivity: An Analysis of Clinical Presentation and Urodynamic Tests From a Large Group of Patients Undergoing Pressure Flow Studies. *Eur Urol.* 2016; 69(2): 361-369. doi:10.1016/j.eururo.2015.08.014

3- Tan TL, Bergmann MA, Griffiths D, Resnick NM. Stop test or pressure-flow study? Measuring detrusor contractility in older females. *NeurourolUrodyn.* 2004; 23: 184-189.[doi.org/10.1002/nau.20020](https://doi.org/10.1002/nau.20020)

4-Defreitas GA, Zimmern PE, Lemack GE, Shariat SF. Refining diagnosis of anatomic female bladder outlet obstruction: comparison of pressure-flow study parameters in clinically obstructed women with those of normal controls. *Urology* 2004; 64: 675-679.

5-Haylen BT, de Ridder D, Freeman RM, Swift SE, Berghmans B, Lee J et al. An international urogynecological association (IUGA)/international continence society (ICS) joint report on the terminology for female pelvic floor dysfunction. *NeurourolUrodyn* 2010; 29: 4-20.

6-Valentini FA, Besson GR, NelsonPP, Zimmern PE. A mathematical micturition model to restore simple flow recordings in healthy and symptomatic individuals and enhance uroflow interpretation. *Neurourol. Urodyn.* 2000; 19(2) 153-176.

7-Valentini FA, Nelson PP, Zimmern PE, Robain G. Detrusor contractility in women: influence of ageing and clinical conditions. *ProgrUrol* 2016; 26: 425-431 DOI 10.1016/j.purol.2016.03.004

8- ValentiniFA,MartiBG, RobainG, ZimmernPE, NelsonPP.Comparison of indices allowing an evaluation of detrusor contractility in women.*ProgUrol*2020 ;30 : 396-401 [doi.org/10.1016/j.purol.2019.11.004](https://doi.org/10.1016/j.purol.2019.11.004)

9- Valdevenito JP, Lopez-Fando L, Abad J, Alvarez D, Valdivieso A. Changes of abdominal pressure during voiding in women. Can abdominal pressure decrease modify the urodynamic diagnosis ? Abstract #112, ICS 2019 meeting Gothenburg.

10- Valdevenito JP, Mercado-Campero A, Lopez-Fando L, Calvo CI, Manriquez V, Medina L. Dropped abdominal pressure at void in women. IUJ 2022;33:3275-3281. doi.org/10.1007/s00192-022-05202-9

11-Hogan S, Gammie A, Abrams P. Urodynamic features and artefacts. NeurourolUrodyn. 2012 Sep;31(7):1104-17.

UNDER PEER REVIEW

**Table 1:** Changes of abdominal pressure during voiding. Nbr number of patients

Nbr (number of women with decrease in $p_{abd}$ ) (%)	Decrease in $p_{abd}$ Unit : cm H <sub>2</sub> O	Nbr (number of women with increase in $p_{abd}$ ) (%)	Increase in $p_{abd}$ Unit : cm H <sub>2</sub> O
145 (53.5%)		81 (29.9%)	
72 (49.6%)	-1 → -4 cm H <sub>2</sub> O	29 (35.8%)	1 → 4 cm H <sub>2</sub> O
36 (24.8%)	-5 → -9 cm H <sub>2</sub> O	17 (21.0%)	5 → 9 cm H <sub>2</sub> O
22 (15.2%)	-10 → -14 cm H <sub>2</sub> O	9 (11.1%)	10 → 14 cm H <sub>2</sub> O
8 (5.5%)	-15 → -19 cm H <sub>2</sub> O	9 (11.1%)	15 → 19 cm H <sub>2</sub> O
7 (4.8%)	≥ -20 cm H <sub>2</sub> O	17 (20.9%)	≥ 20 cm H <sub>2</sub> O

**Table 2:** Number of women with steady change of abdominal pressure during voiding (increase of abdominal pressure = IAP or decrease of abdominal pressure = DAP; No = no change) vs. main complaint (SUI stress urinary

incontinence; MUI mixed urinary incontinence; UUI urge urinary incontinence;  
OTHER: dysuria, frequency complaint, without incontinence)

DAP/IAP	SUI	MUI	UUI	OTHER	p
IAP (Nbr=81)	10	25	31	15	.6552
No (Nbr=45)	13	11	11	10	.1167
DAP(Nbr=145)	32	44	39	30	.6971

**Table 3:**Number of women with changes with steady change of abdominal pressure (increase of abdominal pressure = IAP or decrease of abdominal pressure = DAP; No = no change) vs. urodynamic diagnosis (BOO bladder

outlet obstruction;DHICdetrusor hyperactivity with impaired contractility, DOdetrusor overactivity, DUdetrusor underactivity; N investigations found “normal”; investigations related to urethral dysfunction: intrinsic sphincter deficiency (ISD) and voiding triggered by urethral relaxation (URA)).

DAP/IAP	BOO	DHIC	DO	DU	ISD	N	URA	p
IAP (Nbr=81)	8	5	20	9	16	16	7	.0344
No (Nbr=45)	8	3	8	5	6	12	3	.8433
DAP(Nbr=145)	18	2	30	13	26	51	5	.7220