

Influence of sources of sulphur and zinc on growth and yield of Sesame (*Sesamum indicum* L.)

Comment [A1]:

Abstract:

A field experiment was conducted during *kharij* 2022 at Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Prayagraj, (UP). The soil had a sandy loam texture, with a pH of 7.5, organic carbon (0.625%) and EC (0.564 m.m cm⁻¹), Nitrogen (278.48 kg ha⁻¹), Phosphorus (38.2 kg ha⁻¹) and Potassium (240.7 kg ha⁻¹). The experiment was laid out in Randomized Block Design with nine treatments and control which are replicated thrice on the basis of one year of experimentation. The treatments consisted of 3 levels of source of sulphur (Gypsum 300 kg/ha, Single super phosphate 20 kg/ha, Elementary Sulphur 40 kg/ha) and zinc (5, 10, 15 kg/ha) as a Basal application and a control. The application of Gypsum 300 kg/ha + Zinc 15 kg/ha recorded significantly higher Plant height (110.73 cm), Plant dry weight (15.71 g/plant), number of Capsules/plant (51.40), Seeds/capsules (55.47), Test weight (2.70 g), Seed yield (1.07 t/ha) were recorded with the treatment of Gypsum 300 kg/ha + Zinc 15 kg/ha.

Keywords: Sesame, Gypsum, single super phosphate, sulphur, zinc.

Introduction:

Sesame ranks first for having an oil content of 46-64% and 6355 kcal/kg dietary energy in seeds (Sanjay Kumar Kumar & Goel, 1994). Seeds of sesame is also a rich source of protein (20-28%), sugar (14-16%) and minerals (5-7%). This oil has 85% unsaturated fatty acid viz., is highly stable and has a washing effect on cholesterol & prevents coronary heart disease. Sesame as a valued oil seed appears to have numerous industrial applications. The forecasted results showed for area, production and productivity of the Sesame crop for the year 2020-21 to be 51.66 thousand hectare hectares, 17.64 thousand tonnes and 323.43 kg/hectare respectively. However, improved varieties and agro production technologies capable of increasing the productivity levels of sesame are now developed for different agro ecological situations in the country. A well-managed crop of sesame can yield 1200 - 1500 kg/ha under irrigated and 800 - 1000 kg/ha under rainfed conditions. The crop is grown in almost all parts of the country. More than 85% production of sesame comes from West Bengal, Madhya Pradesh, Rajasthan, Uttar Pradesh, Gujarat, Andhra Pradesh and Telangana (Priyanka 2020).

Comment [A2]: <https://www.thepharmajournal.com/archives/2022/vol11issue4/PartE/11-3-384-994.pdf/>

Comment [A3]: <https://www.ijcmas.com/abstractview.php?ID=18108>

Comment [A4]: <http://ndpublisher.in/ndpbookpage.php?book=319>

Comment [A5]: https://www.business-standard.com/economy/news/why-is-growing-rice-becoming-a-challenge-with-each-passing-year-123041100196_1.html

Comment [A6]: <https://www.ijcmas.com/9-7-2020/N.%20Priyanka%20Evangelin,%20et%20al.pdf>

Sufficient application of Sulphur fertilizer has been documented to improve sesame seed yield and yield related traits [Raja 2007, Vaiyapuri 2004, Sarkar 2005, Shah 2013], as well as oil and protein content in sesame.

The sesame oilcake is a very good cattle feed since it contains protein of high biological value and appreciable quantities of phosphorus and potassium. The cake is also used as manure (Malik et al., 2003).

Gypsum is a soluble source of ~~the~~ essential plant nutrients, (calcium 29.2% and sulfur 18.6%), and can improve overall plant growth. Gypsum amendments can also improve the physical properties of some soils (especially heavy clay soils). Such amendments promote soil aggregation and thus can help prevent ~~the~~ dispersion of soil particles, reduce surface crust formation promote seedling emergence, and increase water infiltration rates and movement through the soil profile. It can also reduce erosion losses of soils and nutrients and reduce concentrations of soluble phosphorus in surface water runoff. Chemical properties improved by ~~the~~ application of gypsum include the mitigation of subsoil acidity and ~~aluminum~~ aluminium toxicity. This enhances deep rooting and the ability of plants to take up adequate supplies of water and nutrients during drought periods.

SSP is an excellent source of three plant ~~nutrient-nutrients~~ (P₂O₅ 16%, Calcium 20%, Sulphur 12%). The P component reacts in soil similarly to other soluble fertilizers. The presence of both P and sulfur (S) in SSP can be an agronomic advantage where both of these nutrients are deficient. In agronomic studies where SSP is demonstrated to be superior to other P fertilizers, it is usually due to the S and/or Ca that it contains. When locally available, SSP has found ~~wide-spread~~ widespread use for fertilizing pastures where both P and S are needed. As a source of P alone, SSP often costs more than other more concentrated fertilizers, therefore it has declined in popularity.

Elemental sulphur, which usually occurs in the form of eight-member rings, is a relatively nontoxic and chemically inert substance and is insoluble in water and most other liquids. It is solid at ambient temperature and exists in two crystalline and amorphous ~~form~~ forms

Zinc is an indispensable micronutrient for crop growth, an important component of carbonic anhydrase and a stimulator of aldolase, which ~~are-is~~ involved in carbon metabolism (Tsonev 2012) Zn is also an integral component of several biomolecules such as lipids, and proteins and co-factor of auxins, and, therefore, it plays an important role in plant nucleic acid metabolism. Zn application has been ~~proved-proven~~ beneficial in improving crop yield and quality [Chattha 2017], while its deficiency reduces yield and deteriorates crop quality. Moreover, higher Zn contents have toxic effects on plants and lead to the suppression of cell division and elongation, affecting biomass production.

Comment [A7]: <https://fabe.osu.edu/sites/fabe/files/imce/files/Soybean/Gypsum%20Bulletin.pdf>

Comment [A8]: http://soilhealth.ucdavis.edu/download_file/view/156/230

Comment [A9]: <https://agribegri.com/products/buy-agriculture-grade-gypsum-online--buy-sulphur-fertilizer.php#:text=Gypsum%20amendments%20can%20also%20improve.phosphorus%20in%20surface%20water%20runoff>

<https://ecotika.in/product/gypsumfertilizer/>

<https://fabe.osu.edu/sites/fabe/files/imce/files/Soybean/Gypsum%20Bulletin.pdf>

[http://ipni.net/publication/nss.nsf/0/5540C741907C7657852579AF007689EC/\\$FILE/NSS-21%20SSP.pdf](http://ipni.net/publication/nss.nsf/0/5540C741907C7657852579AF007689EC/$FILE/NSS-21%20SSP.pdf)

<https://grow-organic-spirulina.com/glossary/single-superphosphate-ssp/>

<https://www.cropnutrition.com/resource-library/single-superphosphate>

<https://www.sciencedirect.com/topics/engineering/elemental-sulfur>

<https://www.mdpi.com/2077-0472/10/9/396>

<https://www.mdpi.com/2077-0472/10/9/396/>

Material and Methods:

The experiment was conducted to know the **Influence of sources of sulphur and zinc on the growth and yield of Sesame (*Sesamum indicum* L.)** was carried out at Crop Research Farm of Sam Higginbottom University, Prayagraj, Uttar Pradesh during in 2022, which is located at 25°39' 42''N latitude, 81°67'56'' E longitude and 98 m altitude above the mean sea level. The soil had a sandy loam texture, with a pH of 7.5, organic carbon (0.625%) and EC (0.564 m.m cm⁻¹), Nitrogen (278.48 kg ha⁻¹), Phosphorus (38.2 kg ha⁻¹) and Potassium (240.7 kg ha⁻¹). The experiment was laid out in a RBD consisting of Ten treatments with 3 replications, with the treatment combinations (T₁) Gypsum 300 kg/ha+Zinc 5 kg/ha, (T₂) Gypsum 300 kg/ha+Zinc 10 kg/ha, (T₃) Gypsum 300 kg/ha+Zinc 15 kg/ha, (T₄) Single super phosphate 20 kg/ha+Zinc 5 kg/ha, (T₅) Single super phosphate 20 kg/ha+Zinc 10 kg/ha, (T₆) Single super phosphate 20 kg/ha+Zinc 15 kg/ha, (T₇) Sulphur 40 kg/ha+Zinc 5 kg/ha, (T₈) Sulphur 40 kg/ha+Zinc 10 kg/ha, (T₉) Sulphur 40 kg/ha+Zinc 15 kg/ha, (T₁₀) Control.

Comment [A10]: <http://eprints.stmone.org/1630/1/Kumar35172023IJPSS101722.pdf>

Results and Discussion:

At Harvest, there was a significant difference among the treatments. However, the highest plant height (110.73 cm) was recorded with the application of Gypsum 300 kg/ha+Zinc 15 kg/ha, whereas minimum plant height (97.57 cm) control, and Gypsum 300 kg/ha+Zinc 10 kg/ha (109.60 cm) was statistically at par with T₃.

At Harvest, there was a significant difference among the treatments. However, the highest dry weight (15.71 gm) was recorded with the application of Gypsum 300 kg/ha+Zinc 15 kg/ha, whereas minimum dry weight (13.23 gm) control, and Gypsum 300 kg/ha+Zinc 10 kg/ha (15.48 gm) was statistically at par with T₃.

The perusal of the data of the Number of Capsules/plants was recorded at harvest, is presented in Table 1. The data reveals that there was a significant effect among different treatments on the Number of Capsules/plants. Significantly ~~Maimum~~ **Maximum** Number of Capsules/plants (51.40) was recorded with the treatment of application Gypsum 300 kg/ha+Zinc 15 kg/ha over all the treatments and the minimum was recorded in (37.40) control. However, treatment Gypsum 300 kg/ha+Zinc 10 kg/ha (50.67) was found to be statistically at par with Gypsum 300 kg/ha+Zinc 15 kg/ha.

Significantly Maximum Number of seeds per ~~capsules~~ **capsule** (55.47) was recorded with the treatment of application Gypsum 300 kg/ha+Zinc 15 kg/ha over all the treatments and the minimum was recorded (45.20) in control. However, treatment Gypsum 300 kg/ha+Zinc 10 kg/ha

(54.40) was found to be statistically at par with Gypsum300 kg/ha+Zinc15 kg/ha.

The perusal of the data of Test weight was recorded at harvest, is presented in Table 1.

The data reveals that there was no significant effect among different treatments on Test weight (g).

Non significantly significant Maximum test weight (2.70 g) was recorded with the treatment of application Gypsum300 kg/ha+Zinc15 kg/ha over all the treatments and a minimum was recorded (2.27 g) in control.

Significantly Maximum seed yield (1.07 t/ha) was recorded with the treatment of application Gypsum300 kg/ha+Zinc15 kg/ha over all the treatments and minimum was recorded (0.88 t/ha) in control. However, treatment Gypsum300 kg/ha+Zinc10 kg/ha (1.06 t/ha) was found to be statistically at par with Gypsum300 kg/ha+Zinc15 kg/ha.

Discussions:

Sulphur through gypsum might have promoted the uptake and translocation of food assimilates from source to sink effectively, resulting in higher yield attributes viz., no. of capsules/plant, the weight of capsules/plant leading to higher seed yield. These results are in agreement with the findings of Duhoon et al. (2005) and Ramakrishna-Ramakrishna (2013) in sesame. The increase in yield might be attributed to the easy availability of sulphate (SO₄) sulphur present in gypsum compared to sulphide form in elemental sulphur, which essentially requires its oxidation to be converted into SOS prior to its absorption by the plants. Among sources of sulphur gypsum proved significantly superior to other sources for seed yield have been reported by Chaurasia-Era (2009) and Ramakrishna (2013). Zinc plays as an activator of several enzymes in plants and it is directly involved in the biosynthesis of growth substances such as auxin thereby producing more plant cells and enhanced dry matter (Christopher Christopher et al., 2019). It has been reported that the Sulphur application not only improves the Sulphur availability itself but also improves the availability of other nutrients too, which are essential for the growth and development of plant plants. It has been also reported that Sulphur helps in reducing soil pH, which helps in the greater availability and mobility of nutrients, especially P, Fe, Mn, and Zn. Reason-The reason for the enhancement of growth parameters might be due to adequate ready supply and increased uptake of sulphur which have resulted into a larger photosynthesizing surface and accelerated the process of formation and translocation of photosynthates and hence overall development of the plant. Higher growth with gypsum source was also reported by Warkad (2018).

Conclusion:

It was concluded that for obtaining higher yield components with better quality of Sesame by application of Gypsum300 kg/ha+Zinc15 kg/ha was recorded a significantly higher number of

Comment [A11]: https://papers.ssrn.com/sol3/Delivery.cfm/SSRN_ID3690067_code4064153.pdf?abstractid=3690067

Capsules/plant (51.40), Seeds/capsules (55.47), seed yield (1.07 t/ha), as compared to other treatments. Since, the finding based on the research done in one season.

UNDER PEER REVIEW

Table 1: Effect of sulphur and zinc on growth and yield parameters of Sesame.

Treatments	Plant Height (cm)	Dry weight (g/plant)	No. of capsules/plant	No. of seeds/capsule	Test weight (g)	Grain Yield (t ha ⁻¹)
T1	107.50	14.82	46.33	51.60	2.57	1.01
T2	109.60	15.48	50.67	54.40	2.67	1.06
T3	110.73	15.71	51.40	55.47	2.70	1.07
T4	98.80	13.40	39.40	46.60	2.40	0.91
T5	99.60	13.55	41.20	47.53	2.43	0.93
T6	103.80	14.20	43.07	49.47	2.50	0.98
T7	101.70	13.99	42.53	47.93	2.47	0.96
T8	104.73	14.49	44.87	51.33	2.53	0.99
T9	108.20	15.06	48.27	53.27	2.60	1.04
T10	97.57	13.23	37.40	45.20	2.27	0.88
Sem(±)	0.76	0.09	0.61	0.51	0.08	0.01
CD(p=0.05)	2.27	0.27	1.82	1.52	-	0.03

References:

- Amudha, A., Vaiyapuri, M. V., SriramaCehandra Ssekharan S., & M. Ravichandran, (2005). Effect of sulphur and organics on yield and sulphur use efficiency in sesame. *Research on Crops*, 6: 468–469.
- Asif, M., Anjum, S.A., Wahid, M.A., Bilal, M.F. (2013). Effect of nitrogen and zinc sulphate on growth and yield of maize (*Zea mays L.*). *Journal of Agricultural Research*. 51 (14): 455464.
- Bairagi IV, Mahulikar DN, Hiwarale JS 2007. Effect of zinc and phosphorus on yield, oil and protein content of soybean *J Soils and Crops*;17(2):292-293.
- Brady NC, Well RR (2002). *The nature and properties of soils*. 13th Ed. Pearson Education (Singapore) Pvt. Ltd. India branch.
- Chaurasia, S. K. Jain, N. & Jain, N. (2009) Effects of integrated use of fertilizer, organic manure and micronutrients on the productivity of sesame (~~Sesamum~~ Sesamum indicum L.) *Annale Agriculture Research New Series* 30 (384), 91-94.
- Chattha, M.U.; Hassan, M.U.; Khan, I.; Chattha, M.B.; Mahmood, A.; Chattha, M.U.; Nawaz, M.; Subhani, M.N.; Kharal, M.; Khan, S 2017. Biofortification of Wheat Cultivars to Combat Zinc Deficiency. *Front. Plant Sci*, 8, 281
- Christopher, A., Oluwagbenga, D., Aruna O., Adekiya, C.C., Khadijat, O., Suleiman, F.O., Okunlola, C.O., Aremu, I.O., Owolabi, C. and Temidayo A. J. 2019. Zinc Sulphate and Boron-Based Foliar Fertilizer Effect on Growth, Yield, Minerals, and Heavy Metal Composition of Groundnut (*Arachis hypogaea L*) Grown on an Alfisol. *International J. Agronomy*. 3:123-127.
- Duboon, S. S., Deshmukh, M. R. Jyotishi, A. & Jam H. C. (2005). Effect of sources and levels of sulphur on seed and oil yield of sesame (*Sesamum indicum L.*) under different agro-climatic situation-situations of india-India *Journal of Oilseed Research* 21 (1) 199-200.
- Dokkara Prem Kumar, Tanuj Kumar Mandal, A. Zaman and Arunabha Pal (2019) Studies on the Effect of Phosphorus Levels on Yield Attributes and Yield of Groundnut in South Odisha Condition, *International Journal of Agriculture, Environment and Biotechnology* 12(4): 339-344.
- Domonoske, Camila (April 20, 2014). "A Legume With Many Names: The Story Of'Goober'". NPR. National Public Radio. Archived from the original on June 6, 2020.
- Kumar R., J.S. Bohra, Narendra K. and Amitesh K.S. (2015) Fodder yield, nutrient uptake and quality of maize (*Zea mays L.*) as influenced by NPKS and Zn fertilization. *Res. on Crops* 16 (2): 243-249.41
- Khan Muhammad Meer, Abdul Manaf, Fayyaz-ulHassan, Muhammad Sheeraz Ahmed (2017). Micronutrients (Zn, B and Mn) Effects on Sesame (*Sesamum indicum L.*). *Int. J Agron. Agri. Res* 10(5):92-101.
- Malik, M.A., Saleem, M.F., Cheema, M.A., Shamim, A. 2003. Influence of different nitrogen levels on the productivity of sesame (*Sesamum indicum L.*) under varying planting patterns. *International Journal of Agriculture and Biology* 5(4), 490-492.

Formatted: Font color: Red

Meena J. K. (2013). Response of seccome (Sesamum dicum L.) varities to varying level levels of sulphur of sulphur and is sources (arter's thesis) Swami Keshwanand Rajasthan Agricultural University, Bikaner.

Mouri SJ, MAR Sarkar, MR Uddin, UK Sarker, MS ~~kaysar~~Kaysar, MMI Hoque (2018)Effectof varietyand phosphorus on theyield components andyield of groundnut29(2):117-126.

Moola Ram, RC Meena and MM Sundria (2021) Enhancing sesame productivity and profitability through zinc and iron application in western Rajasthan ~~The~~ Pharma Innovation Journal 2021; SP-10(10): 924-928.

Priyanka Evangilin N., B. Ramana Murthy, G. Mohan Naidu and B. Aparna (2020) Statistical Model for Forecasting Area, Production and Productivity of Sesame Crop (Sesamum indicum L.) in Andhra Pradesh, India, Int.J.Curr.Microbiol.App.Sci; 9(7): 1156-1166.

Olsen, S.R., Cole, C.V., Watanabe, F.S.& Dean.L.A.(1954). Estimation ofavailablephosphorusinsoilsbyextractionwithNaHCO₃,USDACir.939.U.S.Washingto n.

Pinky Yadav, SS Yadav, Kamal Garg, Sonal Athnere and Seema Yadav (2020) Effect of sulphur and zinc fertilization on productivity and economics of sesame (Sesamum indicum L.) in semi-arid conditions of Rajasthan

Formatted: Font color: Red

Ramakrishna, B. (2013). Growth and yield of sesame (Sesamum indicum L.) as Influenced by Sources ~~and~~and levels of sulphur (Master thesis) University of Agriculture Science Dharmad Karnataka (India) lit pkshikoch~~granth~~ Granthac in handle 1 \$6993 Accessed on March 21.2021.

Ray Hansen (August 2011). "Sesame profile". Agricultural Marketing Resource Center.

Raja, A.; Hattab, K.O.; Gurusamy, L.; Vembu, G.; Suganya, S. Sulphur application on growth and yield and quality of sesame varieties. *Int. J. Agric. Res.* 2007, 2, 599–606.

Vaiyapuri, V.; Amudha, A.; Sriramachandrasedkharan, M.; Imayavaramban, V. Effect of sulphur levels and organics on growth and yield of sesame. *Adv. Plant Sci.* 2004, 17, 681–685.

Sarkar, R.; Saha, A. Analysis of growth and productivity of sesame (*Sesamum indicum*) in relation to nitrogen, sulphur and boron. *Indian J. Plant Physiol.* 2005, 10, 333.

Raja, A.; Hattab, K.O.; Gurusamy, L.; Suganya, S. Sulphur levels on nutrient uptake and yield of sesame varieties and nutrient availability. *Int. J. Soil Sci.* 2007, 2, 278–285.

Shah, M.A.; Manaf, A.; Hussain, M.; Farooq, S.; Zafar-ul-Hye, M. Sulphur fertilization improves the sesame productivity and economic returns under rainfed conditions. *Int. J. Agric. Biol.* 2013, 15, 1301–1306

Singh SM, Singh D (2000). Response of summer groundnut and succeeding maize to sulphurand phosphorus fertilizer. *Indian JAgric.Sci.*70 (10):657-660.

Sujatha, V., Saritha, R., Haseena Bhanu, S.K., Sirisha, A.B.M. and Gangadhara Rao, S.V.S. (2021). Effect of sulphur on growth, yield and economics of sesame (Sesamum indicum). *Internat. J. ~~agrie~~Agric. Sci.*, 17 (AAEBSSD): 233-236,

Subbiah,B.V.andAsija,G.L.(1956)ARapidProcedurefortheEstimationofAvailableNitrogen in Soils. *Current Science*, 25,259-260.

Sarkar, R. K., & Saha, A. (2005). Analysis of growth and productivity of sesame (*Sesamum indicum* L.) in relation to nitrogen, sulphur and boron. *Indian Journal of Plant Physiology*, 10(4): 333-337

Sales, J. M.; Resurreccion, A. V. (2014). "Resveratrol in peanuts". *Critical Reviews in Food Science and Nutrition*. 54 (6): 734–70.

Tsonev, T.; Lidon, F.J.C 2012. Zinc in plants—An overview. *Emir. J. Food Agric*, 24, 322–333.

Toth, S.J.; Prine, A.L. (1949) Estimation of Cation-Exchange Capacity and Exchangeable Ca, K, and Na Contents of Soils by Flame Photometer Techniques, Vol 67 (6), 439-446.

Tathe AS, Patil GD, Khilari J 2008. Effect of sulphur and zinc on groundnut invertisols. *An Asian J of Soil Science*; 3(1):178-180.

Panda, A., Bhale, V.M., Bhattacharjee, S. and Kadam, S, R. (2015-16). Effect of Different Nutrient Management Practices and Zinc Fertilization on Various Growth and Development Stages of Maize (*Zea mays* L.) under Dryland Condition. *International Journal of Current Microbiology and Applied Sciences*. 8 (6).

Rao, Tejeswara, K., Rao, Upendra, A. and Sekhar, D (2013). Effect of sources and levels of groundnut. *Journal Academic & Industrial Research*, 2 (5): 268-270.

Warkad, S. V. (2018). Response of Kharif Sesamum (*Sesamum indicum* L.) to different sources and levels of sulphur (Master's thesis) Vasant Rao Naik Marathwada Krishi Vidyapeeth, Parbhani.