

# Exploring the Potential of Proximal Remote Sensing in Plant Stress Phenotyping: A Comprehensive Review

## ABSTRACT

The global challenge of feeding the world demands attention due to the projected population increase to 10.9 billion by 2050. Abiotic and biotic stressors, such as heat, drought, diseases, and pests, further compound the difficulties faced in achieving sufficient agricultural output. Early detection of crop stress is vital to mitigate yield loss and find appropriate agrotechnical solutions. However, the complex interactions between abiotic and biotic stressors and their impact on plant growth and yield present challenges in plant phenotyping and breeding. This review discusses recent advances in remote sensing technologies which offer promising solutions to overcome these challenges. Low-cost, reliable sensors and technologies facilitate data collection and interpretation, paving the way for proximal sensing and high-throughput phenotyping platforms. These automated platforms, equipped with imaging devices, enable non-destructive data collection and monitoring of plant properties over time. Optical methods like hyperspectral sensors, RGB imaging, remote sensing, and chlorophyll fluorescence contribute to early identification of plant stress causes, facilitating the development of control strategies. By providing accurate and timely information on crop stress, these technologies offer essential support in enhancing agricultural productivity and ensuring food security for a growing global population.

**Keywords:** Remote sensing technologies, proximal sensing, high-throughput phenotyping, imaging devices, hyperspectral sensors, agricultural productivity, food security.

## 1. INTRODUCTION-

Addressing the challenge presented in the 2nd Sustainable Development Goal (SDG) by Food and Agriculture Organisation [1], which pertains to nourishing the global population, stands out as one of the most formidable tasks confronting society in the present era. By 2050, it is anticipated that there will be 10.9 billion people on the planet. As a result, depending on the region, the food supply needs to be increased by 50% to 75% [2]. Due to short-term supply variations, climate change may endanger the stability of entire food systems [3,4]. On a regional scale, the potential consequences might not be immediately evident, but it is probable that areas already vulnerable to hunger and malnutrition will experience a deterioration in food security due to the impact of climate change [4]. As a result, abiotic (such as heat and drought) and/or biotic (such as diseases and pests) conditions and their combinations will also increase in frequency [5], which will result in decreased agricultural output if management does not act quickly and effectively [6]. Early detection of crop stress is therefore crucial to be able to respond with appropriate agrotechnical solutions and thereby minimise yield loss before permanent damage occurs.

Crops often experience an increased number of abiotic and biotic stress combinations as a result of global warming and probable accompanying climate irregularities, which negatively impact their growth and yield [7,8]. Plant stress is characterised as a major departure from the ideal conditions for plant growth that may have detrimental consequences on the plant and restrict its ability to grow and develop normally. Almost every component of a plant can be affected by plant stress, though often only one or a few plant structures are affected, depending on the age of the plant and the source of the stress [9]. Crop plants experience abiotic stress and biotic stress, which are two different forms of environmental stresses. Abiotic stress is responsible for significant crop losses globally and encompasses factors like radiation, salinity, floods, droughts, extreme temperatures, and heavy metal exposure. On the other hand, biotic stresses involve the invasion of various pathogens such as fungi, bacteria, oomycetes, nematodes, and herbivores that negatively impact crop health. [10]. Abiotic stresses, such as heat and drought, have been demonstrated to be more harmful to crop yield when they occur simultaneously than when they do so individually at various stages of crop growth [11,12]. In response to stress, plants experience a number of morphological, physiological, biochemical, and molecular changes that have a negative impact on growth and productivity. Plant phenotyping is currently the main barrier to plant breeding and crop management. Obtaining large-scale plant phenotyping data quickly with high dimensionality, density, and accuracy from a single molecule to a whole organism is now the biggest problem. Although some bottlenecks have been greatly reduced by modern phenomics technology, several problems remain regarding how to accurately and quickly enhance throughput and accuracy while simultaneously defining and extracting complex features [13].

Many of these issues can be resolved because to recent advances in machine learning and remote sensing. As more and more low-cost, reliable sensors and technologies are used for data collection, storage, and interpretation, proximal sensing is in the early stages of a potentially revolutionary development. [14]. Crop genetic improvement has been hampered by the lack of field-based high-throughput phenotyping methods

[15,16], but recent reviews [17,18,19] have highlighted the opportunities now provided by sensor technology and the digital age. High-throughput phenotyping platforms have been created to automate data collection on numerous plants in order to facilitate the gathering of phenotypic measurements [20,21,22]. These platforms frequently have imaging devices installed, which collect data without causing any damage to the plant and track changes in plant properties over time [23].

Earlier, crop monitoring used to be done visually, either from the ground or occasionally from the air, by inspecting the crops. These days, there are optical methods available, including as hyperspectral sensors, RGB imaging, remote sensing, and chlorophyll fluorescence, that can be used to identify plant stress causes early on, enabling for the development of control methods and the minimization of damage brought on by stress. [24]. Plants respond to severe biotic and abiotic circumstances by altering their physiology and metabolism through pulses of gene expression. In order to enhance agricultural development and increase yields, it became essential to identify and adopt novel technologies and approaches. [25]

## **2. STRESS DEFINITIONS AND PHYSIOLOGY**

According to Jones and Jones [26], plants are sessile organisms subject to biotic and abiotic forces, or the physical, chemical, and biological environment defining their habitat. The success or survival of a plant species can be affected by both biotic and abiotic interactions. Plant physiologists have therefore focused on understanding the effects of various stress factors on plants [27], which is crucial for crop sciences (28). According to Lichtenthaler [29], plant stress is generally understood to be "any unfavourable condition or substance that affects or blocks a plant's metabolism, growth, or development." External factors that negatively impact plant growth, development, or productivity rates, crop yields, etc. are referred to as stress in plants. Stress in plants typically results from certain abrupt alterations in the environment. However, exposure to a specific stress results in adaptation to that stress in stress-tolerant plant species in a time-dependent manner. [30]. However, some plants, such as desert plants (Ephemerals), may completely avoid the stress [31].

The reaction is triggered by biochemical factors, such as phytohormones or enzymes, and it results in a cycle of damage and repair, activating the metabolism of the plant and, often, producing minor or imperceptible phenotypic adaptations (reversible and adaptive strain tolerance). "Phenotypic plasticity" is the term used to characterise a plant's capacity to alter the expression of its phenotype in response to environmental factors. According to Kranner et al. [32], the effects of a single stress factor, several stress factors, as well as the combination of biotic and abiotic stresses, can frequently cause very comparable physiological responses in plants. Additionally, the timing of stress effects must be taken into account. For instance, alterations in the flux rates of photosynthesis, respiration, and transpiration may be a sign of brief environmental changes [33]. Long-term stress that occurs during distinct developmental stages can have a variety of effects on growth. To avoid severe crop loss, detecting physiological changes in plants before they become apparent is essential. [34]

When plants are exposed to biotic and abiotic stress, their metabolism is disrupted, implying physiological costs [35, 36, 37, 38] and eventually lowering fitness and production. One of the most crucial aspects is abiotic stress, which has a significant impact on development and, as a result, causes significant losses in the field. In most plant species, the ensuing growth decreases can reach >50% [39]. Additionally, biotic stress is a problem that adds to the harm caused by pathogen or herbivore attack and puts a lot of pressure on plants. Plants undergo long-term adaptations to change their phenotypic manifestations as the severity and length of the stress exposure grow. According to high or low light conditions, this may influence things like leaf size and thickness, stomatal density, or chloroplast function [40]. It should be emphasised that the effects of any stress factor typically lead to a complex interaction between the genes and environment of the plant, frequently resulting in various strains.

## **3. PLANT STRESS AND THEIR COMMON SYMPTOMS**

Multiple published reports have documented crop losses resulting from abiotic stresses [41]. In an ever-changing environment, various detrimental factors, including heat, cold, drought, and salinity, among others, can adversely impact agricultural land and crop productivity. Heat stress refers to prolonged exposure to temperatures beyond a critical threshold, causing irreparable harm to plant well-being [42]. It disrupts plant cell balance, hindering growth and potentially leading to plant mortality [43]. This poses a significant global threat to crop production [44]. Similarly, cold temperatures also impose limitations on the growing seasons of numerous plant species. Chilling stress, apart from causing noticeable phenotypic changes, also leads to significant biochemical and physiological alterations [45,46]. Drought, a major environmental stress, has a severe negative impact on crop yield [47].

Key effects of drought include poor germination [48,49], reduced nutrient availability and photosynthesis [50], decreased leaf number and size [51], and reduced fresh and dry weight of plants [52]. The plant's response to drought is influenced by factors such as growth stages, plant species, age, and the severity and duration of the drought stress [53]. Salinity is recognized as a highly detrimental environmental stress that significantly diminishes crop productivity and quality worldwide [54]. It affects plant growth and development, increases intracellular osmotic pressure, and can lead to toxic levels of sodium accumulation. Heavy metal stress inactivates enzymes and disrupts essential metallic ion substitution reactions, affecting membrane integrity, photosynthesis, and respiration [55]. Heavy metals induce oxidative stress by stimulating the production of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), superoxide radicals (O<sup>-</sup> 2), and hydroxyl radicals (OH) [56]. Excessive ultraviolet (UV) sunlight exposure disrupts plant growth, leading to ROS overproduction and potential cellular imbalance [57]. Waterlogging alters soil properties, causing hypoxia or anoxia, significantly reducing plant growth and survival [58].

Plants face the risk of infection by various pathogens, such as bacteria, fungi, viruses, and nematodes, along with attacks by herbivorous pests [59]. Biotic agents directly induce plant responses; for instance, herbivory results in immediate consequences like water loss, reduction in leaf area, and necrosis [60]. Biotic pathogens

initiate local effects and trigger systemic plant responses, including signal pathways reducing stomatal conductance [61] and increased production of volatile organic compounds (VOCs) as defense mechanisms [62]. Fungal pathogens exhibit diverse effects. Some pathogens initially stimulate photosynthesis, benefiting from increased sugar production during early infection stages [63]. However, in later stages, they can decrease photosynthetic capacity, leading to visible foliar lesions [64]. This fungal diversity gives rise to various plant diseases such as anthracnose, leaf spot, rust, or wilt [65]. These diseases can manifest as changes in the plant canopy's optical reflectance properties, detectable through radiometric changes or computer vision approaches for disease classification at the leaf level [66].




Fig. 1. An overview of major abiotic and biotic stresses .Taken from Umar et al. [67]

#### 4. PROXIMAL REMOTE SENSING TECHNIQUES FOR STRESS PHENOTYPING

Understanding stress-related phenotypic traits and their interrelationships requires precise knowledge of phenotyping in both controlled and field conditions. Advanced techniques, such as visible light imaging, hyperspectral imaging, infrared imaging, fluorescence imaging, and X-ray computed tomography, enable nondestructive acquisition of plant phenotype data using visible to near-infrared light sources [68]. These imaging-based, high-throughput platforms integrated with advanced software systems are valuable tools in plant biology [69]. By focusing on the interaction between light and plants, these imaging techniques provide accurate measurements of quantitative phenotypic traits. Implementing high-dimensional phenotyping requires uniform experimental protocols, calibrated imaging sensors, and precise data analysis. The following section outlines the imaging devices currently used for high-throughput phenotyping of crop plants.

##### 4.1. RGB IMAGING

In order to measure the morphological characteristics of plants, the RGB camera approach is the most extensively used system [70]. This is because it is both affordable and simple to deploy. In contrast to consumer cameras, RGB cameras have an infrared blocking filter (VIS camera), which only picks up light with a wavelength between 400 and 700 nm. To determine the colour of each pixel, the VIS camera uses red, green, and blue colour sensors. It is possible to collect morphological or colour data by using the pixel values of plants that the image processing system has detected [71]. In order to minimise variables and enable in-depth examination of each plant's response to stress, fixed equipment is employed in laboratory research. These studies also allow for the capture of picture data throughout the experiment and the detailed examination of how each plant reacts to stress.

Ge et al. [72] extracted plant pixels from RGB images to correlate with shoot fresh weight, dry weight, and leaf area. Neilson et al. [73] estimated plant biomass and compared it with actual plant size. Positive correlations were found between leaf area and shoot biomass in a water-limiting experiment. Leaf greenness was estimated using the RGB to HIS color system. Field studies use sensors attached to vehicles or aircraft for large-scale analysis. UAV-based RGB sensors provide high-resolution color information for vegetation indices. Bhandari et al. [74] and Francesconi et al. [75] computed canopy features using UAVs for monitoring drought effects on wheat. RGB cameras on fixed facilities reduce data errors from vehicle movement. Becker and Schmidhler [76] used an RGB sensor in a drought treatment facility to estimate yield and growth rate correlation. RGB images enable pixel-level analysis of biomass changes under drought stress. Biomass inference using pixel counts correlates well in various crops [77]. RGB images can predict plant growth changes and analyze specific parts. Color analysis detects leaf wilting and chlorophyll deficiency due to drought stress [78]. RGB sensors on UAVs provide limited upper images for vegetation index analysis. Phenotypic information helps identify candidate genes for specific traits Campbell et al. [77] used the HTP platform to quantify daily shoot biomass and soil water content and simulate shoot growth. Following that, a genome-enabled growth model was combined to identify a number of candidate genes. Using digital image processing, a multi-plant multi-disease detection system was developed in [79], where disease identification was carried out using colour transformations, colour histograms, and a pairwise-based classification system. a smartphone-based image processing software [80] for diagnosing plant diseases, where the created system was tested for diseases in vineyards using images of grape leaves. Through thresholding of the RGB pixel values, the analysis begins by separating region lesions from background and healthy tissue. Spot count, grey

level, and area were among the traits that were extracted, along with colour histograms. 90% of probable diseases were successfully identified by the application.

#### 4.2 NEAR-INFRARED IMAGING

The green parts of plants consistently had the maximum reflectivity at near infrared wavelengths between 700 and 1400 nm in many studies, While unhealthy plants reflect more red light than healthy plants, soil reflects very little near-infrared light. Additionally, freshly absorbed water by leaves scatters near-infrared wavelengths. In order to confirm how plants respond to drought stress, near-infrared imaging (NIR) is used [81]. These infrared imaging systems offer large-field images with excellent spatial resolution and accurate measurements while simultaneously operating in a variety of environmental situations [82].

The plant water content estimated by NIR is also used to identify candidate gene regions. El-Hendawy [83] et al. used a combination of The absorption rate is maximum in the spectral region between 1400 and 1450 nm, and it is strongly associated with the moisture content of plants. With the aid of a partial least squares regression (PLSR) model and an NIR picture [84] were able to locate water absorption bands. NIR can offer a reliable method for non-intrusive, real-time monitoring of the water content of leaves on crop plants. Using NIR imaging, Chen et al. [85] looked into the dynamics of water content and the drought responses of 18 distinct barley cultivars. After experiencing drought stress, plants had a quick decline in NIR signal, which recovered after re-watering.

This technology may immediately identify drought stress in plants since it uses NIR to measure reflectance in a particular wavelength band, which permits speedy study of plant water content. Additionally, by combining NIR imaging with visible imaging, or visible to short-wave infrared (VSWIR; 0.4-2.5 m), this method offers deeper insight into plant health under various stress conditions because it provides well-defined spectral features for pigments, leaf water content, and biochemicals like lignin and cellulose. Further, infrared thermography has also been used to study stomatal responses under salinity and drought by visualizing differences in canopy temperature [86].

#### 4.3 HYPER SPECTRAL IMAGING

Hyperspectral sensors capture numerous bands per pixel across the visible, NIR, and SWIR regions [87]. While the data complexity requires advanced computational resources and storage capacity, high-resolution images with narrow spatial coverage can differentiate responses to different stresses. Hyperspectral imaging enables the detection of early drought stress symptoms, imperceptible to the naked eye. By employing suitable spectral analysis techniques, plant responses to drought stress can be easily evaluated. In laboratory settings, stationary facilities can generate normalized difference vegetation index (NDVI) values by utilizing different hyperspectral wavelength ranges that are segmented based on plant structure.

While imaging hyperspectral sensors combine spectral and spatial resolutions, non-imaging hyperspectral sensors just detect the average spectral information over the area under study [88]. The visible (400–700 nm), near-infrared (700–1000 nm), and short-wave infrared (1000–2500 nm) wavelength ranges were the subject of earlier experiments using hyperspectral sensors [89]. Since plants usually reduce the concentration of leaf chlorophyll, reflectance in the visible range can be correlated to leaf pigment content [90], whereas the near-infrared range is primarily influenced by leaf structure (such as leaf trichome density or leaf thickness) and leaf water content [91].

The chemical compositions of the leaf, such as lignin or cellulose [92], have an impact on the short-wave infrared range in addition to the leaf's water content. Plant phenotyping has not yet utilised the ultraviolet spectrum (200-380 nm). Many plant compounds, including flavonoids, amino acids, anthocyanins, and nucleosides, have UV-range absorption maxima [93, 94, 95]. This method performs well for determining the water content and canopy reflectance of each individual plant section. Ge et al. [96] divided the plant into stem and leaves using a different wavelength and employed hyperspectroscopy to calculate the NDVI of maize. [97,98] have used hyperspectral data to compute various spectral indices and look for relationships with grain yield and drought stress. Specific vegetation indexes can be identified by extracting appropriate wavelengths from hyperspectral data [99,100]. Additionally, for genomic estimated breeding, hyperspectral data have been utilised in conjunction with molecular markers to improve projections of grain yield [101]. Utilising the vegetation index, which is defined as a linear combination or ratio of reflectance at many single wavelengths, is a standard method for the hyperspectral based estimate of plant characteristics under drought stress. There are differences in the photosynthetic apparatus, water content, plant organs, and yield among the physiological and biochemical responses of vegetation to drought stress. In a nutshell hyperspectral imaging can use different vegetable indices to analyse changes in plants under drought stress.

HS imaging is a promising field for pre-symptomatic crop health monitoring. In [102], an early detection system for tobacco mosaic virus (TMV) used HS imaging in the Visible /NIR spectral range (380 nm to 1023 nm). Spectral and textural features at selected wavelengths enabled a four-class classification (healthy, 2 DAI, 4 DAI, and 6 DAI) using machine-learning-based classifiers and the successive projections algorithm (SPA). Similarly, in [103], a HS imaging system detected tomato spotted wilt virus (TSWV) disease in capsicum plants with over 90% accuracy using support vector machine (SVM), classifiers on three types of features: full spectrum, spectral vegetative indices (VIs), and data-driven probabilistic topic model-generated features from both Visible -NIR and Short Wave -NIR hypercubes.

**Table -1 The most commonly utilized spectral indices for crop stress monitoring include pRED, pGREEN, and pBLUE, which represent the spectral reflectance of the red, green, and blue bands, respectively. Additionally, pNIR stands for the reflectance of the near-infrared band and pSWIR refers to the reflectance of the shortwave-infrared band. These indices play a crucial role in assessing and tracking crop stress levels.**

Name	Abbreviation	Formula	Description with related traits and challenges	References
normalized	NDVI	$(\rho_{NIR} - \rho_{RED})/(\rho_{NIR} +$	Assess green vegetation	[104]

difference vegetation index		$\rho_{RED}$	using a normalized ratio within the range of -1 to 1.	
normalized difference water index	NDWI	$\frac{\rho_{NIR} - \rho_{SWIR}}{\rho_{NIR} + \rho_{SWIR}}$	Quantifies variations in leaf water content by utilizing Near-Infrared (NIR) and Shortwave Infrared (SWIR) spectral bands.	[105]
difference vegetation index Jordan	DVI	NIR – Red	Highly responsive to vegetation quantity; characterized by its simplicity as a ratio; however, it does not account for discrepancies between reflectance and radiance due to atmospheric conditions or shadows.	[106]
green normalized difference vegetation index	GNDVI	$\frac{(\rho_{NIR} - \rho_{GREEN})}{(\rho_{NIR} + \rho_{GREEN})}$	An adaptation of NDVI (Normalized Difference Vegetation Index) with increased sensitivity to chlorophyll content.	[107]
photochemical reflectance index	PRI	$\frac{(\rho_{531} - \rho_{570})}{(\rho_{531} + \rho_{570})}$	Serves as an indicator of leaf and plant canopy photosynthetic efficiency.	[108]
structure insensitive pigment index	SIPI	$\frac{(\rho_{800} - \rho_{445})}{(\rho_{800} + \rho_{680})}$	Acts as an indicator of increased canopy stress, particularly related to carotenoid pigment levels.	[109]
moisture stress index	MSI	$\frac{(\rho_{1599})}{(\rho_{819})}$	Highly responsive to rising leaf water content and widely utilized for canopy stress analysis and productivity prediction.	[110]
Leaf water content index Ceccato et al. [144]	LWCI	$\frac{\log(1 - (\rho_{NIR} - \rho_{MIDIR}))}{-\log(1 - (\rho_{NIR} - \rho_{MIDIR}))}$	Assesses the moisture content of the leaf canopy.	[111]
modified red edge NDVI Sims & Gamon [35]	mRENDVI	$\frac{(\rho_{750} - \rho_{705})}{(\rho_{750} + \rho_{705} - 2 * \rho_{445})}$	Leverages the sensitivity of the vegetation red-edge to detect subtle changes in canopy foliage content, gap fraction, and senescence.	[112]

#### 4.4 FLUORESCENCE IMAGING

Fluorescence imaging is employed to estimate the rate of photosynthesis and monitor the impact of plant pathogens [113,114] as well as to detect early stress responses to both abiotic and biotic factors before any decline in growth becomes measurable [115, 116, 117, 118]. In this imaging technique, blue wavelength light with a wavelength of less than 500 nm is directed at the plants, causing them to emit fluorescence light in the red region of the spectrum, specifically at 600-750 nm. These differences in fluorescence are captured through photography and converted into false-color signals using computer software for further analysis [119]. The primary impact of abiotic stresses is on the chlorophyll content, making chlorophyll fluorescence a commonly used tool in phenomics to observe the effects of various environmental factors on genes and the plant's capacity to sustain photosynthesis under such conditions [119]. By measuring chlorophyll fluorescence, which refers to the emission of excessive energy by the plant in the form of fluorescence, fluorescence sensors can evaluate the photosynthetic efficiency of the crop being examined [120]. A visible or ultraviolet (UV) light is used to stimulate the plant, and the resulting fluorescence is captured using charge-coupled device (CCD) cameras [121].

When plants are exposed to UV light in the 340-360 nm range, two types of fluorescence are produced: red + far-red fluorescence and blue-green fluorescence. A multicolor fluorescence imaging principle is utilized to collect fluorescence emitted in four spectral bands, each represented by a specific wavelength—blue (F440), green (F520), red (F690), and near-infrared (F740) [122]. Fluorescence and chlorophyll content serve as crucial indicators of the metabolic states of plants. Additionally, fluorescence imaging can be utilized to study various aspects such as stomatal movement [123], phloem loading and unloading [124], the correlation between spatiotemporal variation of photosynthesis and growth limitation [125], and plant metabolite content [126] under stressful conditions [127,128,129]. Fluorescence imaging has been employed for different purposes, including stress detection at the primary level [130,131,132] and the characterization of heterogeneity in leaf photosynthetic performance [133]. McAusland et al. [134] proposed an experimental method for observing photosynthesis and photoprotection in crops across diverse light environments through

fluorescence analysis. Similarly, in field conditions, the photosynthetic efficiency of a target plant can be assessed by measuring its chlorophyll fluorescence using a portable fluorescence meter in environments with drought stress or adequate watering [135, 136]. Chlorophyll fluorescence is useful for monitoring linear electron transport, which is closely linked to CO<sub>2</sub> uptake during photosynthesis [137].

Several studies have focused on ChlF-based imaging, which includes the detection of sweet potato feathery mottle virus (SPFMV) and sweet potato chlorotic stunt virus (SPCSV) in sweet potato using thermal imaging and ChlF [138]. This study highlighted the operating efficiency of PS-II and photochemical quenching as the most sensitive parameters for quantifying virus effects, surpassing measures like maximum quantum efficiency, non-photochemical quenching, and leaf temperature. Additionally, an early detection system was developed by combining HS- and ChlF-imaging [139]. By integrating both techniques into one device, the classification errors were reduced to less than 5%, leading to more accurate results.

#### 4.5 THERMAL IMAGING

Thermal imaging is used to measure leaf surface temperatures to study plant water relations, and specifically for stomatal conductance, because a major determinant of the leaf temperature is the rate of evaporation or transpiration from a leaf. Abiotic or biotic stresses often result in decreased rates of photosynthesis and transpiration [140,141]; and, the remote sensing of the leaf temperature by thermal imaging can be a reliable way to detect changes in the physiological status of plants in response to different biotic and/or abiotic stresses. This method uses radiation emitted by an object to generate an image, which increases as the temperature of the object increases above absolute zero. Thermal sensors can use visualized image data to detect changes in plant temperature caused by transpiration due to stomatal closure. Therefore, thermal imaging can measure temperature-related features such as water content, transpiration rate, and stomatal conductance through model-based estimation [142]. The canopy temperature has been used successfully in breeding programs for drought-prone environments. In plant phenotyping, thermal imaging offers canopy temperatures to detect differences in stomatal conductance as a measure of the plant response to the water status and transpiration rate [143, 144], both in the field and in the greenhouse. Canopy temperature differences were compared with the surrounding air (for example, the canopy temperature depression, or CTD) as measured by thermal infrared imaging, and these results have been used as a selection criterion in breeding programs for drought resistance [145]. In addition, thermal imaging has been used for many crops, from small cereal grains to maize [146] and fruit trees [147]. It has also been used in combination with spectral imaging for the enhanced estimation of leaf water content [148]. Romano et al. [149] used thermal imaging in maize under reproductive stage drought stress and identified its potential for accelerating phenotyping and screening in maize water stress breeding programs.

A field experiment using a thermal sensor analyzed temperature differences between plants and plots under drought stress. Significant correlations were found between seed yield and canopy temperature, confirming the impact of drought stress on biomass and yield. Relevant genomic regions and extreme genotypes for canopy temperature were identified [150,151]. Aerial thermal infrared image analysis assessed canopy temperature and association mapping identified 52 SNPs significantly associated with canopy temperature. Canopy temperature was used to estimate the crop water stress index (CWSI), an indicator for selecting drought-resistant varieties [152]. Thermal imaging provides valuable information on stomatal conductance, allowing quick evaluation of temperature increase and assessment of changes in plant transpiration rates and drought stress levels. Its use, along with thermal cameras, improves crop phenotyping for drought adaptation [151,153].

Thermal imaging rapidly diagnosed crop diseases [154]. For tomato mosaic disease and wheat leaf rust, the maximum temperature difference (MTD) ranged from 0.2°C to 1.7°C and 0.4°C to 2°C, respectively. MTD increased as the disease progressed, detectable 5 to 7 days before visible symptoms. In apples, IR thermography sensed scab disease caused by *Venturia inaequalia* [155]. MTD linearly correlated with infection size ( $R^2 = 0.85$ ), later decreasing due to leaf senescence.

#### 4.6 NON-IMAGING SPECTROSCOPIC METHODS

Non-imaging spectroscopic methods comprise VIS/IR reflectance and transmittance spectroscopy, along with Raman spectroscopy. VIS/IR spectroscopy is a type of hyperspectral imaging that records only spectral information (not spatial/pixel data) within the VIS/IR wavelength range. Plant health is assessed by measuring either the reflectance of the leaf/tissue surface or the transmitted light through the leaf tissue and correlating it with relevant indicators. A VIS-NIR reflectance spectroscopy system [156] for detecting HLB or citrus greening disease in citrus trees used two portable halogen lamps and a field-portable SVC HR 1024 spectro radiometer. The system collected reflectance data from 350 nm to 2500 nm (989 data points) and included a laser pointer for target area designation. The mobile platform allowed in-field operations, and the best-performing classification algorithm was quadratic discriminant analysis (QDA), with a 95% average accuracy. Multiple recent studies have used spectroscopic methods for plant disease detection. For instance, research has been conducted into the early detection of potato late blight using leaf reflectance measurements with a spectroradiometer [157]. Another study applied NIR spectroscopy for the detection of bitter pit disorder in honeycrisp apples, recording spectra for 40 apples at three different times post-harvest [158]. Non-imaging VIS/IR spectroscopy (VIS/IR-spec) has unique advantages, such as simpler and cheaper setup involving a light source and a spectroradiometer, compared to hyperspectral (HS) imaging. Data analysis is less complex since image pre-processing is unnecessary. However, VIS/IR-spec's lack of spatial information complicates its field application due to potential environmental interference. Raman spectroscopy (RS) is another analytical spectroscopic technique that reveals molecular structure information by analyzing molecular vibrations from inelastic photon collisions. A recent study explored the use of a handheld Raman spectrometer to detect and identify fungal pathogens in maize kernels [159].

**Table 2 Applications and limitations of common sensors used for proximal remote sensing**

Sensor Type	Applications	Limitations
RGB Cameras	Used for imaging canopy cover and canopy color. They provide insights into chlorophyll concentration using greenness indices and enable canopy architecture estimation with 3D stereo reconstruction.	Lack spectral calibration, leading to relative measurements. Susceptible to shadows and changes in light conditions, hindering automated image processing.
Spectral sensors	Assess biochemical composition of leaves and canopies, including pigment concentration and water content. Contribute to canopy architecture assessment with indices like NDVI.	Require frequent calibration and susceptible to changes in light conditions, necessitating white reference calibration. Canopy structure and camera/sun geometries affect measurements, posing data management challenges.
Fluorescence	Suitable for assessing the photosynthetic status of plants and indirectly measuring biotic and abiotic stress factors.	Field-level canopy fluorescence measurements can be challenging due to low signal-to-noise ratio, but techniques like laser-induced fluorescence transients (LIFT) and solar-induced fluorescence extend the range for remote sensing.
Thermal sensors	Employed to measure stomatal conductance and detect water stress induced by biotic or abiotic factors.	Changes in ambient conditions cause canopy temperature fluctuations, making comparisons over time difficult without references. Distinguishing soil temperature from plant temperature in sparse canopies is problematic, hindering automated image processing. Calibration and atmospheric correction often required for accurate results.
X-ray Computed Tomography (CT)	It provides high-resolution, three-dimensional architecture	low automation and low throughput, high cost
Magnetic Resonance Imaging (MRI)	Gives information about water status, transportation, and root architecture, Provides three-dimensional architecture	low throughput and high cost
Positron Emission Tomography (PET)	Gives information about translocation and transport of elements, shows movement and path of positron through the plant	low throughput, high cost

#### 4.7 OTHER IMAGING TECHNIQUES

Functional imaging and optical 3D structural tomography are two recent technological developments that have shifted more and more towards in vivo live imaging of plants. Functional imaging focuses on physiological changes to assess photosynthetic performance under stress, such as ChlF imaging and positron emission tomography (PET). Positron emission tomography (PET) is an imaging technique that quantitatively and nondestructively assesses the 3D spatial distribution and kinetics of radio-tagged biomolecules in a living subject. The elements required for plant growth are  $^{11}\text{C}$ ,  $^{13}\text{N}$ , and  $^{15}\text{O}$ , which are often utilised positron-emitting radionuclides. [160] The transport of  $^{11}\text{C}$ -labeled photo-assimilates can be repeatedly observed in 3D by PET when  $\text{CO}_2$  is consumed during photosynthesis. [161].

Nuclear magnetic resonance is used in nuclear magnetic resonance imaging (MRI), which produces images and recognises nuclear resonance signals coming from  $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{14}\text{N}$ , and  $^{15}\text{N}$ . In seeds [162], whole root systems growing in or near natural soil [163], and entire plants, 3D datasets of plant structures can be acquired using an MRI. It may then assess water spread and movement through xylem and phloem in crops such as castor beans, tomatoes, tobacco, and poplars [164]. It is possible to monitor dynamic changes in plant structure and function by combining MRI and PET technology.

X-ray CT is a computer-processed X-ray technique that can create 3D representations of an object's internal structures from a collection of 2D radiographic images taken around a single axis of rotation. It has been effectively applied for a variety of purposes, including the analysis of soil structural heterogeneity [165] and the visualisation of plant structures [166]. The costs and duration of the scans are the limitations of CT, though.

With the help of fluorescent sensors that are genetically programmed, Förster resonance energy transfer (FRET) is a cutting-edge noninvasive technique for high-resolution imaging of tiny molecules in live tissue. It has been successfully utilised to detect calcium and zinc dynamics in roots during sugar transport [167]. It permits the discovery of numerous paths and dynamic processes of the target molecule. FRET is a superb tool for advanced phenotyping that can solve a number of fundamental queries about plant growth and development. Wheat plants were phenotyped under both control and salt stress conditions using the 3D laser scanner PlantEye. From the overhead data cloud, the system calculated characteristics including 3D leaf area, plant height, and leaf number. In wheat under salt stress, correlations between the manually measured characteristics (leaf area, fresh and dry biomass) and the PlantEye-scanned trait (3D leaf area) were observed [168]. Other sensors may be able to offer 3D structural data. LiDAR is conceivably the most commonly utilised type of sensor for 3D canopy reconstruction [169,170]. Such laser devices have been utilised to determine the plant area density profiles of a wheat canopy and for quick LAI mapping [171,172].

## 5. CONCLUSION

In conclusion, proximal remote sensing offers a powerful and effective approach for plant stress phenotyping. The ability to collect real-time, accurate, and non-destructive data can significantly contribute to our understanding of plant health, leading to improved agricultural practices, increased crop resilience, and better food security. To fully leverage the potential of proximal remote sensing, ongoing research, technological advancements, and collaborative efforts between scientists, farmers, and technology providers are crucial. While proximal remote sensing has shown great promise, there are still challenges and limitations to address. These may include the need for specialized equipment, data processing complexity, and potential interference from environmental factors. Additionally, access to data and affordability can be barriers in some regions.

## References-

1. Anonymous. Sustainable food and home food and agriculture. 2021. Available: [https://zbook.org/sustainable-food-and-home-food-and-agriculture\\_MjUzODMw.html](https://zbook.org/sustainable-food-and-home-food-and-agriculture_MjUzODMw.html).
2. Prosekov AY and Ivanova SA. Food security: the challenge of the present. *Geoforum*. 2018; 91: 73–77.
3. Henry RJ, Rangan P, Furtado A. Functional cereals for production in new and variable climates. *Curr. Opin. Plant Biol.* 2016 ; 30: 11–18.
4. Wheeler T, Von Braun J. Climate change impacts on global food security. *Science*. 2013 ; 341:508–513.
5. Govender M, Govender PJ, Weiersbye IM, Witkowski E, Ahmed F. Review of commonly used remote sensing and ground-based technologies to measure plant water stress. *Water SA*. 2009 ; 35(5)
6. Atzberger C. Advances in remote sensing of agriculture: context description, existing operational monitoring systems and major information needs. *Remote Sens*. 2013; 5: 949–981.
7. Lokers R, Knapen R, Janssen S, van Randen Y, Jansen J. Analysis of big data technologies for use in agro-environmental science. *Environ Model Softw*. 2016.1;84:494-504.
8. Pandey P, Sinha R, Mysore KS, and Senthil KM. Impact of concurrent drought stress and pathogen infection on plants in Combined Stresses in Plants. Cham (Springer). 2015; 203–222.
9. Chen D, Neumann K, Friedel S, Kilian B, Chen M, Altmann T, Klukas C. Dissecting the phenotypic components of crop plant growth and drought responses based on high-throughput image analysis. *Plant Cell*. 2014;26:4636–4655.
10. Gull A, Ahmad L A, Ul Islam Wani N. Biotic and Abiotic Stresses in Plants. In: *Abiotic and Biotic Stress in Plants*. IntechOpen; 2019. Available from: <http://dx.doi.org/10.5772/intechopen.85832>.
11. Mittler R. Abiotic stress, the field environment and stress combination. *Trends Plant Sci*. 2006; 11: 15–19.
12. Prasad PVV, Pisipati SR, Momcilovic I, and Ristic Z. Independent and combined effects of high temperature and drought stress during grain filling on plant yield and chloroplast EF-Tu expression in spring wheat. *J. Agron. Crop Sci*. 2011; 197: 430–441.
13. Yang W, Feng H, Zhang X, Zhang J, Doonan JH, Batchelor WD, Xiong L and Yan J. Crop phenomics and high-throughput phenotyping: Past decades, current challenges, and future perspectives. *Mol. Plant*. 2020; 13: 187–214.
14. Lokers R, Knapen R, Janssen S, van Randen Y, Jansen J. Analysis of big data technologies for use in agro-environmental science. *Environ Model Softw*. 2016.1;84:494-504.
15. Tuberosa R. Phenotyping for drought tolerance of crops in the genomics era. *Front. Physiol*. 2012 ;3
16. Cobb JN, DeClerck G, Greenberg A, Clark R, McCouch S. Next-generation phenotyping: requirements and strategies for enhancing our understanding of genotype-phenotype relationships and its relevance to crop improvement. *Theor. Appl. Genet*. 2013 ;126: 867–887.
17. Araus JL and Cairns JE. Field high-throughput phenotyping: The new crop breeding frontier. *Trends Plant Sci*. 2014; 19: 52–61.
18. White JW, Andrade SP, Gore MA, Bronson KF, Coffelt TA, Conley M M, Feldmann KA, French AN, Heun JT, Hunsaker DJ. Field-based phenomics for plant genetics research. *Field Crops Res*. 2012; 133: 101–112.
19. Cabrera BL, Crossa J, Von ZJ, Serret MD, Araus JL. High-throughput phenotyping and genomic selection: The frontiers of crop breeding converge. *J. Integr. Plant Biol*. 2012 ; 54: 312–320.
20. Granier C, Aguirrezabal L, Chenu K, Cookson S J, Dauzat M, Hamard P, Thioux JJ, Rolland G, Sandrine BC, Lebaudy A, Muller B, Simonneau T, Tardieu F (2006). Phenopsis, an automated platform for

- reproducible phenotyping of plant responses to soil water deficit in *Arabidopsis thaliana* permitted the identification of an accession with low sensitivity to soil water deficit. *New Phytol.* 2006; 169(3):623-35.
21. Busemeyer L, Mentrup D, Möller K, Wunder E, Alheit K, Hahn V, Maurer HP, Reif JC, Würschum T, Müller J, et al. BreedVision — A Multi-Sensor Platform for Non-Destructive Field-Based Phenotyping in Plant Breeding. *Sensors.* 2013; 13(3):2830-2847.
  22. Virlet N, Sabermanesh K, Sadeghi T P, and Hawkesford M J. Field Scanalyzer: an automated robotic field phenotyping platform for detailed crop monitoring. *Funct. Plant Biol.* 2017 ; 44 :143–153.
  23. Humplík J F, Lazár D, Fürst T, Husièková A, and Spíchal, L. Automated phenotyping of plant shoots using imaging methods for analysis of plant stress responses – a review. *Plant Methods.* 2015 ;11:29.
  24. Prasad and Lakshman. Monitoring biotic and abiotic stress in crop plants through remote sensing. *Chron Bio Manag.* 2021; 5(2) :33-36.
  25. Galieni A, D'Ascenzo N, Stagnari F, Pagnani G, Xie Q and Pisante M. Past and future of plant stress detection: an overview from remote sensing to positron emission tomography. *Front. Plant Sci.* 2021;11:609155.
  26. Jones HG, Jones MB. Plants under stress. Introduction: some terminology and common mechanisms. *Soc Exp Biol.* 1989.
  27. Larcher W. *Physiological Plant Ecology: ecophysiology and Stress Physiology of Functional Groups.* Springer-Verlag Berl Heidelberg. 2003.
  28. Lichtenthaler HK. Vegetation stress: an introduction to the stress concept in plants. *J Plant Physiol.* 1996;148:4-14.
  29. Blum A. Stress, strain, signaling, and adaptation – not just a matter of definition. *J Exp Bot.* 2016;67:562-565
  30. Verma S, Nizam S, Verma PK. Biotic and abiotic stress signalling in plants. In: *Stress Signaling in Plants: Genomics and Proteomics Perspective.* 2013;1:25-49
  31. Zhu JK. Salt and drought stress signal transduction in plants. *Annu Rev Plant Biol.* 2002;53:247-273.
  32. Kranner I, Minibayeva FV, Beckett RP, Seal CE. What is stress? Concepts, definitions and applications in seed science. *New Phytol.* 2010;188:655-673.
  33. Damm A, Paul LE, Haghighi E, Simmer C, Morsdorf F, Schneider F, Van der T C., Migliavacca M, Rascher U. Remote sensing of plant-water relations: an overview and future perspectives. *J. Plant Physiol.* 2018;227: 3–19.
  34. Pachori S, Thakur S, Barela A, Tomar A, Nagre SP, Anand KJ, Patel T, Sharma SK. Remote Sensing for Crop Management: A Comprehensive Review. *Biol Forum Int J.* 2023;15(2):382-387.
  35. Heil M and Bostock RM. Induced systemic resistance (ISR) against pathogens in the context of induced plant defences. *Ann Bot.* 2002;89:503-512.
  36. Swarbrick PJ, Schulze-Lefert P, Scholes JD. Metabolic consequences of susceptibility and resistance in barley leaves challenged with powdery mildew. *Plant Cell Environ.* 2006;29:1061-1076.
  37. Bolton MV. Primary metabolism and plant defense-Fuel for the fire. *Mol Plant Microbe Interact.* 2009;22:487-497.
  38. Massad TJ, Dyer LA, Vega CG. Cost of defense and a test of the carbon-nutrient balance and growth-differentiation balance hypotheses for two co-occurring classes of plant defense. *PLoS One.* 2012;7:7554.
  39. Wang W, Vinocur B, Altman A. Plant responses to drought; salinity and extreme temperatures: Towards genetic engineering for stress tolerance. *Planta.* 2003;218:1-14.
  40. Sarto MVM, Sarto JRW, Rampim L, Rosset JS, Bassegio D, Costa PFd, Inagaki AM. Wheat phenology and yield under drought: a review. *Aust J Crop Sci.* 2017;11:941-946
  41. Wu G, Zhang CA, Chu LY, Shao HB. Responses of higher plants to abiotic stresses and agricultural sustainable development. *J Plant Interact.* 2007;2:135-147.
  42. Wahid A, Gelani S, Ashraf M, Foolad MR. Heat tolerance in plants: An overview. *Environ Exp Bot.* 2007;61:199-223.
  43. Kotak S, Larkindale J, Lee U, von Koskull-Doring P, Vierling E, Scharf KD. Complexity of the heat stress response in plants. *Curr Opin Plant Biol.* 2007;10:310-316.
  44. Hall AE. *Crop Responses to Environment.* CRC press; 2000.
  45. Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K. Effects of abiotic stress on plants: A systems biology perspective. *BMC Plant Biol.* 2011;11:163.
  46. Rucker K, Kvien C, Holbrook C, Hook J. Identification of peanut genotypes with improved drought avoidance traits. *Peanut Sci.* 1995;22:14-18.
  47. Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A, Sadia S, Nasim W, Adkins S, Saud S. Crop production under drought and heat stress: Plant responses and management options. *Front Plant Sci.* 2017;8:1147.
  48. Yadav S, Modi P, Dave A, Vijapura A, Patel D, Patel M. Effect of Abiotic Stress on Crops. In: Hasanuzzaman M, Filho M, Fujita M, Nogueira T, editors. *Sustainable Crop Production.* IntechOpen; Rijeka, Croatia: 2020.
  49. Shahid MJ, Ali S, Shabir G, Siddique M, Rizwan M, Seleiman MF, Afzal M. Comparing the performance of four macrophytes in bacterial assisted floating treatment wetlands for the removal of trace metals (Fe, Mn, Ni, Pb, and Cr) from polluted river water.
  50. Manickavelu A, Nadarajan N, Ganesh S, Gnanamalar R, Babu RC. Drought tolerance in rice: Morphological and molecular genetic consideration. *Plant Growth Regul.* 2006;50:121-138.
  51. Rucker K, Kvien C, Holbrook C, Hook J. Identification of peanut genotypes with improved drought avoidance traits. *Peanut Sci.* 1995;22:14-18.

52. Zhao TJ, Sun S, Liu Y, Liu JM, Liu Q, Yan YB, Zhou HM. Regulating the drought-responsive element (DRE)-mediated signaling pathway by synergic functions of trans-active and trans-inactive DRE binding factors in *Brassica napus*. *J Biol Chem*. 2006;281:10752-10759.
53. Gray SB, Brady SM. Plant developmental responses to climate change. *Dev Biol*. 2016;419:64-77.
54. Arora NK. Impact of climate change on agriculture production and its sustainable solutions. *Environ Sustain*. 2019;2:95-96
55. Hossain MA, Piyatida P, da Silva JAT, Fujita M. Molecular mechanism of heavy metal toxicity and tolerance in plants: Central role of glutathione in detoxification of reactive oxygen species and methylglyoxal and in heavy metal chelation. *J Bot*. 2012;2012:872875.
56. Ghorri NH, Ghorri T, Hayat MQ, Imadi SR, Gul A, Altay V, Ozturk M. Heavy metal stress and responses in plants. *Int J Env Sci Technol*. 2019;16:1807-1828.
57. Singh S, Fatima A, Tiwari S, Prasad SM. Plant responses to radiation stress and its adaptive mechanisms. In: *Plant Life Under Changing Environment*. Academic Press, Cambridge, MA, USA; 2020. p. 105-122.
58. Tewari S, Mishra A. Flooding stress in plants and approaches to overcome. In: *Plant Metabolites and Regulation Under Environmental Stress*. Academic Press, Cambridge, MA, USA; 2018. p. 355-366.
59. Atkinson NJ, Urwin PE. The interaction of plant biotic and abiotic stresses: from genes to the field. *Journal of Experimental Botany*. 2012;63:3523-3543.
60. Meza-Canales ID, Meldau S, Zavala JA, Baldwin IT. Herbivore perception decreases photosynthetic carbon assimilation and reduces stomatal conductance by engaging 12-oxo-phytodienoic acid, mitogen-activated protein kinase 4 and cytokinin perception. *Plant Cell Environ*. 2017;40:1039-1056.
61. Aldea M, Hamilton JG, Resti JP, Zangerl AR, Berenbaum MR, Elucia EHd. Indirect effects of insect herbivory on leaf gas exchange in soybean. *Plant Cell Environ*. 2005;28:402-411.
62. Holopainen JK, Gershenzon J. Multiple stress factors and the emission of plant VOCs. *Trends Plant Sci*. 2010;15:176-184.
63. Chang Q, Liu J, Wang Q, Han L, Liu J, Li M, Huang L, Yang J, Kang Z. The effect of *Puccinia striiformis* f. sp. *tritici* on the levels of water-soluble carbohydrates and the photosynthetic rate in wheat leaves. *Physiol Mol Plant Pathol*. 2013;84:131-137.
64. Vergara-Diaz O, Kefauver SC, Elazab A, Nieto-Taladriz MT, Araus JL. Grain yield losses in yellow-rusted durum wheat estimated using digital and conventional parameters under field conditions. *Crop J*. 2015;3:200-210.
65. Jain A, Sarsaiya S, Wu Q, Lu Y, Shi J. A review of plant leaf fungal diseases and its environment speciation. *Bioengineered*. 2019;10:409-424.
66. Singh S, Fatima A, Tiwari S, Prasad SM. Plant responses to radiation stress and its adaptive mechanisms. In: *Plant Life Under Changing Environment*. Academic Press, Cambridge, MA, USA; 2020. p. 105-122.
67. Umar BO, Amudalat RL, Shehu AA, Lukman BA, Khadijat AA, Ramat BM, et al. Stresses in Plants: Biotic and Abiotic. In: *Current Trends in Wheat Research*. IntechOpen; 2022.
68. Rahaman MM, Chen D, Gillani Z, Klukas C, Chen M. Advanced phenotyping and phenotype data analysis for the study of plant growth and development. *Front Plant Sci*. 2015;6:619
69. Paproki A, Sirault X, Berry S, Furbank R, Fripp J. A novel mesh processing based technique for 3D plant analysis. *BMC Plant Biol*. 2012;12:63-71. Petrellis N. A smart phone image processing application for plant disease diagnosis. In: *Proceedings of the 2017 6th International Conference on Modern Circuits and Systems Technologies (MOCASST)*, Thessaloniki, Greece, 4–6 May 2017; pp. 1–4.
70. Rosero A, Pérez J-L, Rosero D, Burgos-Paz W, Martínez R, Morelo J, Pastrana I, Burbano E, Morales A. Morphometric and colourimetric tools to dissect morphological diversity: An application in sweet potato [*Ipomoea batatas* (L.) Lam.]. *Genet Resour Crop Evol*. 2019;66:1257-1278.
71. Faragó D, Sass L, Valkai I, András N, Szabados L. PlantSize offers an affordable, non-destructive method to measure plant size and color in vitro. *Front Plant Sci*. 2018;9:219.
72. Ge Y, Atefi A, Zhang H, Miao C, Ramamurthy RK, Sigmon B, Yang J, Schnable JC. High-throughput analysis of leaf physiological and chemical traits with VIS-NIR-SWIR spectroscopy: A case study with a maize diversity panel. *Plant Methods*. 2019;15:1–12.
73. Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K. Effects of abiotic stress on plants: A systems biology perspective. *BMC Plant Biol*. 2011;11:163.
74. Bhandari M, Baker S, Rudd JC, Ibrahim AM, Chang A, Xue Q, Jung J, Landivar J, Auvermann B. Assessing the Effect of Drought on Winter Wheat Growth Using Unmanned Aerial System (UAS)-Based Phenotyping. *Remote Sens*. 2021;13(6):1144.
75. Francesconi S, Harfouche A, Maesano M, Balestra GM. UAV-Based Thermal, RGB Imaging and Gene Expression Analysis Allowed Detection of Fusarium Head Blight and Gave New Insights Into the Physiological Responses to the Disease in Durum Wheat. *Front Plant Sci*. 2021;12:551.
76. Becker E, Schmidhalter U. Evaluation of yield and drought using active and passive spectral sensing systems at the reproductive stage in wheat. *Front Plant Sci*. 2017;8:379.
77. Campbell MT, Grondin A, Walia H, Morota G. Leveraging genome-enabled growth models to study shoot growth responses to water deficit in rice. *J Exp Bot*. 2020;71(19):5669-5679.
78. Deng XP, Shan L, Inanaga S, Inoue M. Water-saving approaches for improving wheat production. *J Sci Food Agric*. 2005;85:1379-1388.
79. Barbedo JGA, Koenigkan LV, Santos TT. Identifying multiple plant diseases using digital image processing. *Biosyst Eng*. 2016;147:104-116.

80. Petrellis N. A smartphone image processing application for plant disease diagnosis. In: Proceedings of the 2017 6th International Conference on Modern Circuits and Systems Technologies (MOCASST); 2017 May 4-6; Thessaloniki, Greece. p. 1-4
81. Khan Z, Rahimi-Eichi V, Haefele S, Garnett T, Miklavcic SJ. Estimation of vegetation indices for high-throughput phenotyping of wheat using aerial imaging. *Plant Methods*. 2018;14:1-11.
82. Li L, Zhang Q, Huang D. A review of imaging techniques for plant phenotyping. *Sensors*. 2014;14:20078-20111
83. El-Hendawy SE, Hassan WM, Al-Suhaibani NA, Schmidhalter U. Spectral assessment of drought tolerance indices and grain yield in advanced spring wheat lines grown under full and limited water irrigation. *Agric Water Manag*. 2017;182:1-12.
84. Das B, Sahoo RN, Pargal S, Krishna G, Verma R, Viswanathan C, Sehgal VK, Gupta VK. Evaluation of different water absorption bands, indices and multivariate models for water-deficit stress monitoring in rice using visible-near infrared spectroscopy. *Spectrochim Acta Part A Mol Biomol Spectrosc*. 2021;247:119104.
85. Chen D, Neumann K, Friedel S, Kilian B, Chen M, Altmann T, Klukas C. Dissecting the phenotypic components of crop plant growth and drought responses based on high-throughput image analysis. *Plant Cell*. 2014;26:4636-4655.
86. Rahaman MM, Chen D, Gillani Z, Klukas C, Chen M. Advanced phenotyping and phenotype data analysis for the study of plant growth and development. *Front Plant Sci*. 2015;6:619.
87. Ge Y, Atefi A, Zhang H, Miao C, Ramamurthy RK, Sigmon B, Yang J, Schnable JC. High-throughput analysis of leaf physiological and chemical traits with VIS-NIR-SWIR spectroscopy: A case study with a maize diversity panel. *Plant Methods*. 2019;15:1-12.
88. Thomas S, Kuska MT, Bohnenkamp D, Brugger A, Alisaac E, Wahabzada M, Behmann J, Mahlein AK. Benefits of hyperspectral imaging for plant disease detection and plant protection: A technical perspective. *J Plant Dis Prot*. 2018;125:5-20.
89. Mahlein AK, Kuska MT, Behmann J, Polder G, Walter A. Hyperspectral sensors and imaging technologies in phytopathology: State of the art. *Annu Rev Phytopathol*. 2018;56:535-558.
90. Carter GA, Knapp AK. Leaf optical properties in higher plants: Linking spectral characteristics to stress and chlorophyll concentration. *Am J Bot*. 2001;88:677-684.
91. Slaton MR, Raymond Hunt E Jr, Smith WK. Estimating near-infrared leaf reflectance from leaf structural characteristics. *Am J Bot*. 2001;88:278-284.
92. Curran PJ. Remote sensing of foliar chemistry. *Remote Sens Environ*. 1989;30:271-278.
93. Fischer J. Specific detection of nucleotides, creatine phosphate, and their derivatives from tissue samples in a simple, isocratic, recycling, low-volume system. *LC GC Int*. 1995;8:254.
94. Giusti MM, Polit MF, Ayvaz H, Tay D, Manrique I. Characterization and quantitation of anthocyanins and other phenolics in native Andean potatoes. *J Agric Food Chem*. 2014;62:4408-4416.
95. Talrose V, Yermakov A, Usov A, Goncharova A, Leskin A, Messineva N, Trusova N, Efimkina M. UV/visible spectra. In: NIST Chemistry WebBook, NIST Standard Reference Database; National Institute of Standards and Technology: Gaithersburg, MD, USA, 2009; p. 20899.
96. Ge Y, Bai G, Stoerger V, Schnable JC. Temporal dynamics of maize plant growth, water use, and leaf water content using automated high throughput RGB and hyperspectral imaging. *Comput Electron Agric*. 2016;127:625-632.
97. Rischbeck P, Elsayed S, Mistele B, Barmeier G, Heil K, Schmidhalter U. Data fusion of spectral, thermal and canopy height parameters for improved yield prediction of drought stressed spring barley. *Eur J Agron*. 2016;78:44-59.
98. Tattaris M, Reynolds MP, Chapman SC. A direct comparison of remote sensing approaches for high-throughput phenotyping in plant breeding. *Front Plant Sci*. 2016;7:1131.
99. Elsayed S, Rischbeck P, Schmidhalter U. Comparing the performance of active and passive reflectance sensors to assess the normalized relative canopy temperature and grain yield of drought-stressed barley cultivars. *Field Crops Res*. 2015;177:148-160.
100. Rutkoski J, Poland J, Mondal S, Autrique E, Pérez LG, Crossa J, Reynolds M, Singh R. Canopy temperature and vegetation indices from high-throughput phenotyping improve accuracy of pedigree and genomic selection for grain yield in wheat. *G3 Genes Genomes Genet*. 2016;6:2799-2808.
101. Trachsel S, Dhliwayo T, Gonzalez Perez L, Mendoza Lugo JA, Trachsel M. Estimation of physiological genomic estimated breeding values (PGEBV) combining full hyperspectral and marker data across environments for grain yield under combined heat and drought stress in tropical maize (*Zea mays* L.). *PLoS ONE*. 2019;14
102. Zhu H, Chu B, Zhang C, Liu F, Jiang L, He Y. Hyperspectral Imaging for Presymptomatic Detection of Tobacco Disease with Successive Projections Algorithm and Machine-learning Classifiers. *Sci Rep*. 2017;7:4125.
103. Moghadam P, Ward D, Goan E, Jayawardena S, Sikka P, Hernandez E. Plant Disease Detection Using Hyperspectral Imaging. In: Proceedings of the 2017 International Conference on Digital Image Computing: Techniques and Applications (DICTA), Sydney, NSW, Australia, 29 November-1 December 2017; pp. 1-8.
104. Rouse J, Haas RH, Schell JA, Deering D. Monitoring vegetation systems in the great plains with ERTS. In: Proceedings of the Earth Resources Technology Satellite Symposium, pp. 309-317. Washington, DC: NASA; 1973.
105. Gao B-C. NDWI—A normalized difference water index for remote sensing of vegetation liquid water from space. *Remote Sens Environ*. 1996;58.
106. Jordan CF. Derivation of leaf-area index from quality of light on the forest floor. *Ecology*. 1969;50:663-666.
107. Agapiou A, Hadjimitsis DG, Alexakis DD. Evaluation of broadband and narrowband vegetation indices for the identification of archaeological crop marks. *Remote Sens*. 2012;4:3892-3919.

108. Gamon JA, Peñuelas J, Field CB. A narrowwaveband spectral index that tracks diurnal changes in photosynthetic efficiency. *Remote Sens Environ.* 1992;41:35–44.
109. Peñuelas J, Filella I, Lloret P, Munˆ Oz F, Vilajeliu M. Reflectance assessment of mite effects on apple trees. *Int J Remote Sens.* 1995;16:2727–2733.
110. Behmann J, Steinrücken J, Plümer L. Detection of early plant stress responses in hyperspectral images. *ISPRS J Photogramm Remote Sens.* 2014;93:98–111.
111. Ceccato P, Gobron N, Flasse S, Pinty B, Tarantola S. Designing a spectral index to estimate vegetation water content from remote sensing data: Part 1: theoretical approach. *Remote Sens Environ.* 2002;82:188–197.
112. Sims DA, Gamon JA. Relationships between leaf pigment content and spectral reflectance across a wide range of species, leaf structures and developmental stages. *Remote Sens Environ.* 2002;81:337–354.
113. Balachandran S, Hurry V, Kelley S, Osmond C, Robinson S, Rohozinski J, Seaton G, Sims D. Concepts of plant biotic stress. Some insights into the stress physiology of virus-infected plants, from the perspective of photosynthesis. *Physiol Plant.* 1997;100:203–213.
114. Lohaus G, Heldt H, Osmond C. Infection with phloem limited abutilon mosaic virus causes localized carbohydrate accumulation in leaves of abutilon striatum: Relationships to symptom development and effects on chlorophyll fluorescence quenching during photosynthetic induction. *Plant Biol.* 2000;2:161–167.
115. Baker NR. Chlorophyll fluorescence: A probe of photosynthesis in vivo. *Annu Rev Plant Biol.* 2008;59:89–113.
116. Chaerle L, Hagenbeek D, De Bruyne E, Van Der Straeten D. Chlorophyll fluorescence imaging for disease-resistance screening of sugar beet. *Plant Cell Tissue Organ Cult.* 2007;91:97–106.
117. Konishi A, Eguchi A, Hosoi F, Omasa K. 3D monitoring spatio-temporal effects of herbicide on a whole plant using combined range and chlorophyll a fluorescence imaging. *Funct Plant Biol.* 2009;36:874–879.
118. Lenk S, Chaerle L, Pfündel EE, Langsdorf G, Hagenbeek D, Lichtenthaler HK, van der Straeten D, Buschmann C. Multispectral fluorescence and reflectance imaging at the leaf level and its possible applications. *J Exp Bot.* 2007;58:807–814.
119. Weirman A, 2010. Plant Phenomics Teacher Resource. [http://www.plantphenomics.org.au/files/teacher/Final\\_Phenomics\\_for\\_word\\_with\\_image.doc](http://www.plantphenomics.org.au/files/teacher/Final_Phenomics_for_word_with_image.doc).
120. Rajendran K, Tester M, Roy SJ. Quantifying the three main components of salinity tolerance in cereals. *Plant Cell Environ.* 2009; 32: 237–249
121. Chaerle L, Hagenbeek D, De Bruyne E, Van Der Straeten D. Chlorophyll fluorescence imaging for disease-resistance screening of sugar beet. *Plant Cell Tissue Organ Cult.* 2007;91:97–106.
122. Buschmann, C.; Langsdorf, G.; Lichtenthaler, H.K. Fluorescence: The blue, green, red and far-red fluorescence signatures of plant tissues their multicolour fluorescence imaging and application for agrofood assessment. *Opt. Monit. Fresh Process. Agric. Crop.* 2008, 32, 272–319.
123. Cardon ZG, Mott KA, Berry JA. Dynamics of patchy stomatal movements, and their contribution to steady-state and oscillating stomatal conductance calculated using gas-exchange techniques. *Plant Cell Environ.* 1994;17:995-1007.
124. Siebke K, Weis E. Assimilation images of leaves of *Glechoma hederacea*: analysis of non-synchronous stomata related oscillations. *Planta.* 1995;196:155-65.
125. Walter A, Rascher U, Osmond B. Transitions in photosynthetic parameters of midvein and interveinal regions of leaves and their importance during leaf growth and development. *Plant Biol (Stuttg).* 2004;6:184-91.
126. Li L, Zhang Q, Huang D. A review of imaging techniques for plant phenotyping. *Sensors.* 2014;14:20078–20111.
127. Chaerle L, Van Der Straeten D. Seeing is believing: imaging techniques to monitor plant health. *Biochim Biophys Acta.* 2001;1519:153-66.
128. Rascher U, Hutt MT, Siebke K, Osmond B, Beck F, Lüttge U. Spatio-temporal variations of metabolism in a plant circadian rhythm: the biological clock as an assembly of coupled individual oscillators. *Proc Natl Acad Sci U S A.* 2001;98:11801-5.
129. Osmond B, Ananyev G, Berry J, Langdon C, Kolber Z, Lin G, Monson R, Nichol C, Rascher U, Schurr U, Smith S. Changing the way we think about global change research: scaling up in experimental ecosystem science. *Glob Chang Biol.* 2004;10:393-407.
130. Atkinson NJ, Urwin PE. The interaction of plant biotic and abiotic stresses: from genes to the field. *Journal of Experimental Botany.* 2012;63:3523-3543.
131. Konishi A, Eguchi A, Hosoi F, Omasa K. 3D monitoring spatio-temporal effects of herbicide on a whole plant using combined range and chlorophyll a fluorescence imaging. *Funct Plant Biol.* 2009;36:874–879.
132. Chen D, Ming C, Thomas A, Christian K. Bridging genomics and phenomics: approaches in integrative bioinformatics. Springer, Berlin, Heidelberg; 2014. p. 299-33.
133. Baker NR. Chlorophyll fluorescence: A probe of photosynthesis in vivo. *Annu Rev Plant Biol.* 2008;59:89–113.
134. McAusland L, Atkinson JA, Lawson T, Murchie EH. High throughput procedure utilising chlorophyll fluorescence imaging to phenotype dynamic photosynthesis and photoprotection in leaves under controlled gaseous conditions. *Plant Methods.* 2019;15:1-15.
135. Begovic L, Galic V, Abicic I, Loncaric Z, Mlinaric S. Implications of intra-seasonal climate variations on chlorophyll a fluorescence and biomass in winter barley breeding program. *Photosynthetica.* 2020;58:995-1008.
136. Hao D, Chao M, Yin Z, Yu D. Genome-wide association analysis detecting significant single nucleotide polymorphisms for chlorophyll and chlorophyll fluorescence parameters in soybean (*Glycine max*) landraces. *Euphytica.* 2012;186:919-31.

137. Tietz S, Hall CC, Cruz JA, Kramer DM. NPQ (T): A Chlorophyll Fluorescence Parameter for Rapid Estimation and Imaging of Non-Photochemical Quenching of Excitons in Photosystem-II-Associated Antenna Complexes. *Plant Cell Environ.* 2017.
138. Wang L, Poque S, Valkonen JP. Phenotyping viral infection in sweetpotato using a high-throughput chlorophyll fluorescence and thermal imaging platform. *Plant Methods.* 2019;15:116.
139. Bauriegel E, Herppich WB. Hyperspectral and Chlorophyll Fluorescence Imaging for Early Detection of Plant Diseases, with Special Reference to *Fusarium spec.* Infections on Wheat. *Agriculture.* 2014;4:32-57.
140. Nilsson HE. Remote sensing and image analysis in plant pathology. *Can J Plant Pathol.* 1995;17:154-66.
141. Chaerle L, van der Straeten D. Imaging techniques and the early detection of plant stress. *Trends Plant Sci.* 2000;5:495-501.
142. Ishimwe R, Abutaleb K, Ahmed F. Applications of thermal imaging in agriculture—A review. *Adv Remote Sens.* 2014;3:128.
143. Munns R, James RA, Sirault XR, Furbank RT, Jones HG. New phenotyping methods for screening wheat and barley for beneficial responses to water deficit. *J Exp Bot.* 2010;61:3499-3507.
144. Jones HG, Serraj R, Loveys BR, Xiong L, Wheaton A, Price AH. Thermal infrared imaging of crop canopies for the remote diagnosis and quantification of plant responses to water stress in the field. *Funct Plant Biol.* 2009;36:978-989.
145. Blum A, Mayer J, Gozlan G. Infrared thermal sensing of plant canopies as a screening technique for dehydration avoidance in wheat. *Field Crops Res.* 1982;5:137-146.
146. Colombo G, Vaz-de-Melo A, Taubinger M, de Faria E, Tavares RDC. Phenotyping of maize genotypes under aluminum stress and its correlation with grain yield. *Rev Agrar.* 2014;7:60-71.
147. Gonzalez-Dugo V, Zarco-Tejada P, Nicolás E, Nortes P, Alarcón J, Intrigliolo D, Fereres E. Using high resolution uav thermal imagery to assess the variability in the water status of five fruit tree species within a commercial orchard. *Prec Agric.* 2013;14:660-678.
148. Zia S, Romano G, Spreer W, Sanchez C, Cairns J, Araus J, Müller J. Infrared thermal imaging as a rapid tool for identifying water-stress tolerant maize genotypes of different phenology. *J Agron Crop Sci.* 2013;199:75-84.
149. Romano G, Zia S, Spreer W, Sanchez C, Cairns J, Araus JL, Müller J. Use of thermography for high throughput phenotyping of tropical maize adaptation in water stress. *Comput Electron Agric.* 2011;79:67-74.
150. Ge Y, Atefi A, Zhang H, Miao C, Ramamurthy RK, Sigmon B, Yang J, Schnable JC. High-throughput analysis of leaf physiological and chemical traits with VIS-NIR-SWIR spectroscopy: A case study with a maize diversity panel. *Plant Methods.* 2019;15:1–12.
151. Kaler AS, Ray JD, Schapaugh WT, Asebedo AR, King CA, Gbur E, Purcell LC. Association mapping identifies loci for canopy temperature under drought in diverse soybean genotypes. *Euphytica.* 2018;214:1-18.
152. Rischbeck P, Elsayed S, Mistele B, Bartheimer G, Heil K, Schmidhalter U. Data fusion of spectral, thermal and canopy height parameters for improved yield prediction of drought stressed spring barley. *Eur J Agron.* 2016;78:44-59.
153. Jones H. *Plants and Microclimate: A Quantitative Approach to Environmental Plant Physiology*; Cambridge University Press: Cambridge, UK, 2013; pp. 1–407.
154. Zhu W, Chen H, Ciechanowska I, Spaner D. Application of infrared thermal imaging for the rapid diagnosis of crop disease. *IFAC-PapersOnLine.* 2018;51:424-430.
155. Oerke E, Fröhling P, Steiner U. Thermographic assessment of scab disease on apple leaves. *Precis Agric.* 2011;12:699-715.
156. Sankaran S, Mishra A, Maja JM, Ehsani R. Visible-near infrared spectroscopy for detection of Huanglongbing in citrus orchards. *Comput Electron Agric.* 2011;77:127-134.
157. Gold KM, Townsend PA, Herrmann I, Gevens AJ. Investigating potato late blight physiological differences across potato cultivars with spectroscopy and machine learning. *Plant Sci.* 2020;295:110316.
158. Kafle GK, Khot LR, Jarolmasjed S, Yongsheng S, Lewis K. Robustness of near infrared spectroscopy based spectral features for non-destructive bitter pit detection in honeycrisp apples. *Postharvest Biol Technol.* 2016;120:188-192.
159. Farber C, Kurouski D. Detection and Identification of Plant Pathogens on Maize Kernels with a Hand-Held Raman Spectrometer. *Anal Chem.* 2018;90:3009-3012.
160. Komarov S, Tai YC. Positron Emission Tomography (PET) for Molecular Plant Imaging. *Methods Mol Biol.* 2022;2539:97-118.
161. Bühler J, Huber G, Schmid F, Blümli P. Analytical model for long-distance tracer-transport in plants. *J Theor Biol.* 2011;270:70-79.
162. Melkus G, Rolletschek H, Fuchs J, Radchuk V, Grafahrend-Belau E, Sreenivasulu N, Rutten T, Weier D, Heinzl N, Schreiber F. Dynamic <sup>13</sup>C/<sup>1</sup>H NMR imaging uncovers sugar allocation in the living seed. *Plant Biotechnol J.* 2011;9:1022-1037.
163. Moradi A, Oswald S, Nordmeyer-Massner J, Pruessmann K, Robinson B, Schulin R. Analysis of nickel concentration profiles around the roots of the hyperaccumulator plant *berkheya coddii* using mri and numerical simulations. *Plant Soil.* 2010;328:291-302.
164. Windt CW, Vergeldt FJ, De Jager PA, Van As H. MRI of long distance water transport: a comparison of the phloem and xylem flow characteristics and dynamics in poplar, castor bean, tomato and tobacco. *Plant Cell Environ.* 2006;29:1715-1729.
165. Pierret A, Capowiez Y, Belzunces L, Moran C. 3D reconstruction and quantification of macropores using X-ray computed tomography and image analysis. *Geoderma.* 2002;106:247-271.
166. Stuppy WH, Maisano JA, Colbert MW, Rudall PJ, Rowe TB. Three-Dimensional analysis of plant structure using high-resolution x-ray computed tomography. *Trends Plant Sci.* 2003;8:2-6.
167. Jones AM, Danielson JA, Manoj Kumar SN, Lanquar V, Grossmann G, Frommer WB. Abscisic acid dynamics in roots detected with genetically encoded FRET sensors. *Elife.* 2014;3:e01741.

168. Maphosa L, Thoday-Kennedy E, Vakani J, Phelan A, Badenhorst P, Slater A, Spangenberg G, Kant S. Phenotyping wheat under salt stress conditions using a 3D laser scanner. *Isr J Plant Sci.* 2016;1:1–8.
169. Eitel JUH, Vierling LA, Long DS, Hunt ER. Early season remote sensing of wheat nitrogen status using a green scanning laser. *Agric For Meteorol.* 2011;151:1338–1345.
170. Sanz R, Rosell J, Llorens J, Gil E, Planas S. Relationship between tree row LIDAR-volume and leaf area density for fruit orchards and vineyards obtained with a LIDAR 3D Dynamic Measurement System. *Agric For Meteorol.* 2013;171-172:153-162.
171. Gebbers R, Ehlert D, Adamek R. Rapid Mapping of the Leaf Area Index in Agricultural Crops. *Agron J.* 2011;103:1532-1541.
172. Hosoi F, Omasa K. Estimating vertical plant area density profile and growth parameters of a wheat canopy at different growth stages using three-dimensional portable lidar imaging. *ISPRS J Photogramm Remote Sens.* 2009;64:151-158.

UNDER PEER REVIEW