

## Integrated nutrient management on semi *rabi* green gram

---

### ABSTRACT

An experiment was conducted to study the response of organic sources on the growth, yield, and quality of semi-rabi green gram. The experiment was conducted during semi rabi season 2021, at Agronomy Farm, B.A. College of Agriculture, Anand Agricultural University, Anand, Gujarat, comprising one chemical treatment with nine integrated nutrient management and *biofertilizer* treatments i.e. total 10 integrated organic manures treatments impose in semi rabi green gram. The layout of experiment was done according to Randomized complete block design with three replications. Growth attributes like plant height, number of branches, dry matter accumulation plant<sup>-1</sup> and dry weight of nodules plant<sup>-1</sup> of green gram as influenced by an application of 75% N through vermicompost + ST (5 ml kg seed<sup>-1</sup>) of *Rhizobium* + Bio NPK consortium 1 L ha<sup>-1</sup>. Significantly higher green gram yield attributing character and biological yield was observed significantly higher where application done by 75% N through vermicompost + ST (5 ml kg seed<sup>-1</sup>) of *Rhizobium* + Bio NPK consortium 1 L ha<sup>-1</sup>. Quality parameter and content is most important factors for green gram, an application of 75% N through vermicompost + ST (5 ml kg seed<sup>-1</sup>) of *Rhizobium* + Bio NPK consortium 1 L ha<sup>-1</sup> treatment reported significantly higher seed and straw NPK content and uptake.

**Keywords:** biofertilizer consortium, castor cake, farmyard manure, green gram, organic sources, seed, stover yield and vermicompost

### Introduction

India is a largest producer of Pulses (Joshi *et al.*, 2016 and Kavaya *et al.*, 2021) and its universally utilized as food legumes, which come in second position in terms of produce and

utilization after cereals. Being an essential component of the food of the great majority of vegetarians and the foundation of sustainable agricultural production. For livelihood activities, about 65-70% of the population depend on agricultural and its relative activities. Nearly agricultural contributes 17% of domestic growth. Pulses have a rich content in protein (25-26%), vitamins (3%), fibres, carbohydrate (51%) and minerals nutritional like iron, zinc and magnesium also having essential amino acids, which play a vital role in human health (Yadav *et al.*, 2017). Due to its many uses, green gram is a major pulse crop in India, where it is growing under a different agroclimatic conditions. (Banotra *et al.*, 2022).

Pulses, which are high in protein and play an active part in human food, are the second most significant section of Indian agriculture after grains. (Chaudhary *et al.*, 2018). In India, pulse farming is important role in agriculture (Kundal *et al.*, 2023). These crops are grown for domestic human consumption, mainly by vegetarian people due to nutritional factors. Pulses have capacity to fix atmospheric nitrogen (average 30-50 kg ha<sup>-1</sup>) into soil and ultimately to improve soil fertility, being recognized as natural fertilizer factory (Makwana *et al.*, 2020). Pulses are called “marvel of nature” and are cultivated in kharif, rabi as well as summer season. They are pointed out as fundamental sources of protein, presenting 14% of total protein in the typical Indian diet. As a third important pulse crops, can be used as vegetables, fodder and green manuring in India (Saravanan *et al.*, 2013). Pulse seed is more palatable, nutritive, digestible and not flatulent than other pulse grown in country (Angmo *et al.*, 2022). In current agriculture system, chemical products can demonstrate a lack of sustainability and generate harmful effects such as soil erosion, associated plant nutrient losses, surface and ground water pollution from agrochemical and sediment; also affecting the low farm income from higher cost (Joshi *et al.*, 2016). One of the most essential agronomic practices for boosting crop output and preserving

soil fertility is nutrient management (Patel *et al.*, 2019). A sustainable farming system must have pulse. They improve soil health and fix nitrogen into the soil, which reduces the requirement for water (Dubey *et al.*, 2023).

“Green gram is a staple meal that has a protein content of roughly 25%, which is about three times that of grains (Sengupta and Tamang, 2015). Overall, the area planted with pulses has grown over time, however the productivity level is not very satisfying because of continuous cropping and negligence about inorganic fertilizers. The crop is often grown using only conventional fertilizers, without the use of biofertilizers or any other supplemental sources of nutrients (Pushpa *et al.*, 2022). Integrated plant nutrition management promotes economic yield as well as the physicochemical aspects of the soil. (Tripathi and Tiwari, 2022). When crop residues are applied along with chemical fertilizers, the chemical characteristics, specifically the organic carbon content and nutrient availability, improve. The use of fertilizer, organic manures, and biofertilizers in tandem, also known as integrated nutrient management, is the most effective technique for satisfying plant nutrient demands, resolving the issue of nutrient mining, and preserving soil health. (Tripathi and Tiwari, 2022). The important component of enhanced yield is nutrient balance. Excessive and unbalanced fertilizer usage has resulted in soil mining, which has reduced agricultural output and, ultimately, soil health (Mishra *et al.*, 2023).

Organic manure involves both micro and macro nutrients, and it enhances soil fertility, soil physical, chemical, and biological qualities; it also maintains soil microbial population balance. Furthermore, reduce soil erosion and increase plant nutrient availability through the mineralization process. (Kishor *et al.*, 2021 and Kumar *et al.*, 2011). It is widely acknowledged that the combination of organic manures and biofertilizers plays a critical role in boosting soil production. (Kishor *et al.*, 2021). Biofertilizers made from microalgae have

been found to be an excellent alternative to toxic chemical fertilizers in the agriculture sector (Dineshkumar *et al.*, 2020). Biofertilizers are primarily responsible for boosting nutrient availability by using various microorganisms; they are also beneficial in enhancing atmospheric nitrogen fixation in soil and for increasing the availability of nutrients in soil that are easily accessible to plants. The use of organic manures and biofertilizer is crucial for soil structure and nutrient management. They improve nutrient availability and soil health by converting a fixed form of nutrient into a soluble form, increasing soil fertility and productivity (Perli *et al.*, 2022).

The need for an inexpensive alternative to inorganic fertilizer has grown significantly in popularity due to the high price of inorganic fertilizer, growing environmental awareness, and the energy crisis. Given this, the current study on integrated nutrient management is more beneficial for enhancing the physical, chemical, and biological aspects of the soil as well as the production of healthy foods, increasing income, and lowering cultivation costs.

Organic manures and biofertilizer have the ability to promote development and provide immunity in the plant system by lowering the soil microbial population. Excessive and inefficient application of inorganic fertilizers reduced crop quality and harmed the soil. As a result, total yields increase. This may be accomplished by substituting different nutrients using organic methods. This research was designed to examine integrated nutrient management affects the development, production, quality and economics of green gramme under semi-rabi conditions.

## **METHODOLOGY**

The present research investigation was carried out in the semi *rabi* season (2021) at the Agronomy Farm, B. A. College of Agriculture, Anand Agricultural University, Anand, Gujarat, India. The experimental site is located at 22°-35' N latitude 72°-55' longitude and an altitude of 45.1 meters above mean sea level in the Anand district of Gujarat. Metrological data are

presented in Fig. 1. The soil of top 15 cm depth of experimental site was loamy sand in texture, low in organic carbon (0.30%) and available nitrogen ( $210 \text{ kg ha}^{-1}$ ), medium in available phosphorus ( $40 \text{ kg ha}^{-1}$ ) and available potash ( $285 \text{ kg ha}^{-1}$ ) with 8.05 soil pH. The details of the ten treatments viz., (T<sub>1</sub>) 100% RDF (20:40:0 kg NPK  $\text{ha}^{-1}$ ), (T<sub>2</sub>) 100% N through farm yard manure (FYM), (T<sub>3</sub>) 100% N through vermicompost, (T<sub>4</sub>) 100% N through castor cake, (T<sub>5</sub>) 75% N through FYM + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$ , (T<sub>6</sub>) 75% N through vermicompost + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$ , (T<sub>7</sub>) 75% N through castor cake + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$ , (T<sub>8</sub>) 50% N through FYM + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$ , (T<sub>9</sub>) 50% N through vermicompost + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$  and (T<sub>10</sub>) 50% N through castor cake + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$ . The inputs nutrient was analyzed before sowing (FYM, vermicompost, castor cake containing 0.32, 1.21 and 3.24 % N, respectively) while initial microbial count  $117.60 \times 10^7$  cfu/g observed in soil. Tested organic manures were applied 10 days before planting and integrated into the soil, *rhizobium* was used as a seed treatment, and Bio NPK consortium was applied by soaking and irrigation. At the maturity five randomly selected plants from each plot were harvested separately to study the growth and yield attributing parameters. The crop was harvested, threshed and cleaned manually and seed and stover yield per plot have been recorded. The results from the field experiment were statistically analyzed to compare treatments using analysis of variance techniques (ANOVA) for RBD design and the result were interpreted at 5% level of significance (Gomez and Gomez, 1984).

## RESULTS AND DISCUSSION

### *Response on growth attributes*

Perusal of data depicted in Table 1 indicated that response of integrated nutrient management treatment on green gram was found to be significant. Application of 75% N through vermicompost + seed treatment ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$  ( $T_6$ ) reported significantly higher plant height (52.32 and 54.33 cm at 60 DAS (days after sowing) and at harvest, respectively) as well as higher number of branched per plant (5.20) at harvest as compared to treatments. Manure treatment may have encouraged improved root proliferation, increased phosphorous solubility, and hence enhanced biological nitrogen fixation, nutrient absorption, and during the plant growth nutrient availability of all nutrients during the growth phase. As a result, the plant grew taller. (Verma *et al.*,2022). Vermicompost and FYM has improved nutrient availability due to speedy mineralization and improved nutrient availability to plants, advancing plant growth and development. The use of organic manures and biofertilizers enhanced green gram growth. The maximum number of branches per plant might be due to synergistic result; it is well recognized that organic manures increase chlorophyll concentration, cell division, shoot growth, and photosynthetic rate (Priyadarshini *et al.*,2021). According to the findings, adding manure to the soil increased nutrient availability over time and had a positive effect on plant height. A more balanced diet might have resulted in faster development and growth. Nitrogen is one of the most significant components influencing the leaf area index, which may have aided in agricultural photosynthesis and productivity. The response of organic manures on leaf area index might be due to (Rajeshwari,2011 and Prajapati, 2014) supply of macro and micronutrients, vitamins, and growth hormones like as gibberellins, which increased leaf area and hence photo assimilates. Another reason might be due to the practice of keeping soil fertility and plant nutrient delivery at an optimal level in order to sustain goal

productivity by optimizing the benefits from all viable organic and biological components except inorganic is referred to as integrated organic nutrient management (Pandey *et al.*, 2019 and Das 2018).

Presented Table-1 indicated that application organic manures with biofertilizer combination reported its significant response on dry matter accumulation and dry weight of root nodules per plant. Higher dry matter accumulation ( $31.87 \text{ g plant}^{-1}$ ) and dry weight of root nodules per plant at 45 DAS ( $45.67 \text{ g}$ ) were observed higher in treatment T<sub>6</sub> (75% N through vermicompost + seed treatment ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1\text{L ha}^{-1}$ ) than other treatments. Organic manure application may have increased organic matter content and microbial activity and ultimately improved soil conditions for nutrient availability, whereas the addition of biofertilizers may have enhanced the biochemical processes of nitrogen fixation and phosphorus solubilization, boosting nutrient availability and promoting crop development. Nutrient availability improved due to higher microbial availability and activities ultimately improved by using biofertilizers. Biofertilizers have improved soil nutrient availability by improving microbial activity and releasing nutrients from soil, which aids in adequate nutrient absorption and utilization by plants (Gohain and Kikon 2017; Kaur 2016). In general, proper provision of plant nutrients as per crop requirement via solubilization of insoluble phosphorus and potassium, as well as favorable environmental conditions in the rhizosphere, which supported the crop's longer vegetative and reproductive phases. (Singh *et al.*, 2021).

**Table 1:** Response of treatments on growth parameters of semi *rabi* green gram

Treatment	Plant height (cm)		Branches/plant	Dry matter accumulation/plant (g plant <sup>-1</sup> )	Dry weight of root nodules/plant
	60 DAS	At harvest	At harvest		45 DAS
T <sub>1</sub>	48.87	51.15	4.90	28.33	23.00
T <sub>2</sub>	44.67	46.78	3.73	24.67	29.73
T <sub>3</sub>	45.60	47.74	3.93	25.67	32.20
T <sub>4</sub>	39.52	41.35	2.40	24.33	23.77
T <sub>5</sub>	50.27	52.62	5.13	30.33	44.40
T <sub>6</sub>	52.32	54.33	5.20	31.87	45.67
T <sub>7</sub>	49.33	51.65	4.93	29.27	44.13
T <sub>8</sub>	47.04	49.23	4.33	26.33	43.40
T <sub>9</sub>	47.93	50.18	4.53	27.50	43.87
T <sub>10</sub>	46.57	48.74	4.13	26.00	42.20
<b>SEm ±</b>	<b>1.98</b>	<b>2.06</b>	<b>0.35</b>	<b>1.29</b>	<b>1.54</b>
<b>CD (p=0.05)</b>	<b>5.89</b>	<b>6.12</b>	<b>1.03</b>	<b>3.83</b>	<b>4.56</b>

***Effect of treatment on yield attributes and yield***


The yields varied significantly because of diverse organic nutrient sources and their combinations. (Table 2 and Fig.1). Dry matter accumulation and dry weight of root nodules per plant of green gram as affected by different organic manure treatments depicted in Table 2. An application of 75% N through vermicompost + ST (5 ml kg<sup>-1</sup> seed) of *Rhizobium* + Bio NPK consortium 1L ha<sup>-1</sup> (T<sub>6</sub>) reported significantly higher pods per plant (33.20) and length of pod (8.60 cm) it was found to be significantly superior over all the treatments. Significantly maximum seed (1481 kg ha<sup>-1</sup>) and straw (2008 kg ha<sup>-1</sup>) yield than rest of treatments (Fig.1). As compared to RDF treatment (T<sub>1</sub>), treatment T<sub>6</sub> (75% N through vermicompost + seed treatment (5 ml kg<sup>-1</sup> seed) of *Rhizobium* + Bio NPK consortium 1L ha<sup>-1</sup>) reported 12.48% higher seed yield. The improved seed yield and yield-attributing characteristics of green gram due to organic soil application may be increased nutrient availability to plants, in addition to increased water holding capacity and other physical properties, which may be due to the formation of more root nodules, vigorous root development, and better nitrogen fixation (Shete *et al.*, 2010), the

development of plant growth leading to higher photosynthesis activity and translocation resulted in improvement of yield attributes and finally higher seed yield (Ravikumar *et al.*, 2012). Another explanation for increased economic output might be the reaction of yield-attributing variables such as number of pods per plant, length of pods, number of seeds per pod, and test weight to improved yield resulting in a greater economic yield. Nutrient availability may also have increased photosynthate synthesis and improved translocation within plants led in enhanced growth of photosynthate sink source ratio. Improvement in higher biological yields recorded could be attributed to better plant growth and dry matter accumulation as recorded under these treatments (Gohain and Kikon 2017).

**Table 2: Response of treatments on yield attributes and yield of semi *rabi* green gram**

Treatment	Pods/plant	Pod length (cm)	Test weight (g)	Protein content in seed (%)	Net return (₹/ha)	B:C ratio
T <sub>1</sub>	29.60	7.60	45.25	21.42	40583	2.46
T <sub>2</sub>	27.60	5.80	44.61	20.38	27367	1.87
T <sub>3</sub>	29.27	6.00	44.88	20.60	27134	1.77
T <sub>4</sub>	22.80	5.47	44.17	20.17	19932	1.61
T <sub>5</sub>	31.40	8.00	45.36	21.77	50613	2.66
T <sub>6</sub>	33.20	8.60	45.41	22.48	51231	2.54
T <sub>7</sub>	30.40	7.80	45.32	21.52	45802	2.46
T <sub>8</sub>	28.60	7.20	45.11	21.10	40307	2.39
T <sub>9</sub>	29.20	7.40	45.17	21.23	41878	2.36
T <sub>10</sub>	28.20	7.00	44.95	20.94	35944	2.22
SEm ±	<b>1.34</b>	<b>0.33</b>	<b>0.92</b>	<b>0.27</b>	-	-
CD (p=0.05)	<b>3.99</b>	<b>0.99</b>	<b>NS</b>	<b>0.80</b>	-	-

**Selling price of green gram:** Seed @ ₹ 50/kg for RDF, @ ₹ 55/kg seed from organic sources, and straw @ ₹ 1.5/kg



**Fig 1: Effect of integrated organic manures treatments on seed and straw yield**

## QUALITY PARAMETERS


### *NPK content, uptake and protein content*

The evaluation of seed quality criteria of the resulting seed crop is a critical component in determining the planting value and the performance of seeds as a result of treatments. The use of organic manures in conjunction with biofertilizers altered seed nutrient content, absorption, and protein content of green gram.

Different organic manures on nutrient content and uptake by seed and straw were found to be significant (Fig 2). Application of 75% N through vermicompost + seed treatment ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$  ( $T_6$ ) reported higher nitrogen content in seed and straw (3.6 and 0.65 %, respectively) but it remains statistically at par with treatment  $T_5$  (75% N through FYM + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$ ). An application of 75% N through vermicompost + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1 \text{ L ha}^{-1}$  ( $T_6$ ) reported higher NPK uptake by seed (53.27, 7.38 and  $9.08 \text{ NPK kg ha}^{-1}$ )

and straw (12.91, 3.16 and 32.78 NPK kg ha<sup>-1</sup>). Data presented in Table 2 revealed that green gram seed protein content (22.48%) reported higher by application of 75% N through vermicompost + ST (5 ml kg<sup>-1</sup> seed) of *Rhizobium* + Bio NPK consortium 1L ha<sup>-1</sup> (T<sub>6</sub>). Higher nitrogen concentration in biological products might be attributed to organic manure, which contains all macro and trace nutrients and is gradually and slowly released, helping to crop nutrition balance. It also improves the soil environment for root development and proliferation. As a result, plants will absorb more nutrients, resulting in higher nutrient content in seed and stover in green gram (Shariff *et al.* (2017) and Singh *et al.* (2017). In general, the use of organic manure alone or combinations with *Rhizobium* and PSB, was found to improve NPK absorption by both seed and stover, with the combined treatment resulting in a greater uptake (Fig. 2).

ascribed to improved soil fertility and greater nutrient availability and higher nutrient absorption by crop might be due to combined application of organic manures and biofertilizers (Gohain and Kikon, 2017). The significantly protein content by application of integrated organic manures treatments might be due to nitrogen levels are continuously available at all stages of the crop growth; culminating in slow release of nitrogen through organic sources and resulting in later increased protein content in seed. Then, resulting later stage nitrogen availability through



increased protein content in seed.

**Fig 2: Effect of different organic sources on NPK uptake by seed and straw**

**ECONOMICS**

The Perusal of data show in Table 2 indicated that net returns in green gram were influenced significantly due to various treatments of organic manures. Results presented in Table-2 indicated that the higher net return (₹ 51231/ha) and B:C ratio (2.66) were obtained with application of 75% N through vermicompost + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1\text{L ha}^{-1}$  and 75% N through FYM + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1\text{L ha}^{-1}$  respectively, which could be due to the higher seed yield during the growing season, higher selling price of organic treated seed and low cost of FYM. The lowest B:C ratio of 1.61 was recorded under 100% N through castor cake.

**CONCLUSION**

In order to increase seed yield and quality of green gram as well as maintain soil health, the use of organic sources for fulfilling nutrient requirement to sustain crop productivity and safeguard soil human-environment health is recommended. Thus, it can be concluded that the application of 75% N through vermicompost + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1\text{L ha}^{-1}$  gave significantly higher seed yield ( $1480 \text{ kg ha}^{-1}$ ), net return (₹ 51231/ha) and B:C ratio (2.66) with the application of 75% N through FYM + ST ( $5 \text{ ml kg}^{-1}$  seed) of *Rhizobium* + Bio NPK consortium  $1\text{L ha}^{-1}$ .

## REFERENCES

- Angmo, P., Mandal A.K., Phogat M., Kumar S. and Rai A.P., 2022. Effect of phosphorus and zinc interaction of performance of green gram [(*Vigna radiata* (L.). Wilczek)]. Legume Research; 45(10): 1283-1286.
- Banotra, Monika, Sharma B.C., Kumar Rakesh, Mahajan A., 2022. Influence of Differential Substitution of Nutrients through Organics on Growth, Yield and Nutrient Content of Green Gram (*Vigna radiata*) in Irrigated Sub-tropical Region of J and K. Legume Research. 45(7): 866-871. doi: 10.18805/LR-4823.
- Chaudhary, M., Singh, S., Babu, S and Prasad, M., 2018. Effect of integrated nutrient management on productivity, nutrient acquisition and economics of black gram (*Phaseolus mungo* L.) in an inception of eastern Uttar Pradesh. Legume Research;41(5):759-762.
- Das, S.K. and Avasthe, R.K., 2018. Soil organic nutrients management through integrated approach: A policy for Environment & Ecology. Aug. 20;4(1): 1-8.
- Dineshkumar R., Duraimurugan M., Sharmiladevi N., Priya Lakshmi L., Ahamed A., Rasheeq, Arumugam A. and Sampathkumar P., 2020. Microalgal liquid biofertilizer and bio stimulant effect on green gram (*Vigna radiata* L) an experimental cultivation. *Biomass Conversion and Biorefinery* volume 12, 3007–3027.
- Dubey, L.R., Rathava R.S., Saini L.H. and Saini A.K., 2023. Analysis of marketing status and buying behavior of *kharif* green gram seeds in banaskantha district of North Gujarat. *The Pharma Innovation Journal*;12(3); 1429-1434.
- Gohine R.T. and Kikon N., 2017. Optimization of organic nutrient sources for green gram (*Vigna radiata* L. Welczek) under rainfed conditions. *Indian J. Agric. Res.*, 51(5): 443-447.

- Gomez, K.W. and Gomez, A.A., 1984. Statistical procedures for Agricultural Research. Willy-Inter Science publication, New York, USA, pp.60-65.
- Joshi, D. Gediya, K.M., Gupta, S and Birari, M.M., 2016: Effect of organic manures on soil and quality parameters of cowpea [*Vigna unguiculata* (L.) Walp] under middle Gujarat. *Agric. Sci., Digest*; 36(3): 216-2019.
- Kaur, H., 2016. Effect of biofertilizers and organic fertilizers on soil health, growth and yield of green pea (*Pisum sativum* L.). M.Sc Thesis, Punjab Agricultural University, Ludhiana.
- Kavya P., Singh, S., Hinduja, N., Towari, D. and Sruthi, S., 2021. Effect of foliar application of micronutrients on growth and yield of green gram. *Legume Research*; 44(12):1460-1464.
- Kishor K., Kumar V., Upadhaya B. and Borpatragohain B.2021. Effect of integrated nutrient management on growth yield and economics of summer mungbean (*Vigna radiata* L.). *The pharma innovation journal*;10(8):978-983.
- Kumar, A.B.M., Gowda N.C.N., Shetty G.R., Karthik M.N., 2011. Effect of organic manures and inorganic fertilizers on available NPK, microbial density of the soil and nutrient uptake of brinjal. *Research journal of Agricultural Sciences*; 2(2): 304-307.
- Kaundal, M., Kaur,N., Seth, M. and Singh, V., 2023. Effect of integrated nutrient management through phosphorus and vermicompost on growth, phenology and yield of chickpea (*Cicer arietinum* L.). *Journal of Crop and Weed*, 19(1): 48-52
- Makwana,S. N., Patel, G. G., Patel, H. K., Shiyal, V. N. and Patel, B. K., 2020. Effect of Inorganic and Organic Nutrients on Growth and Yield of Summer Green Gram (*Vigna radiata* L. wilczek). *Int.J.Curr.Microbiol.App.Sci.*; Special Issue-11: 730-738.
- Mishra, S. and Chaturvedi, P., 2023. Effect of integrated nutrient management of growth and yield of green gram (*Vigna radiation* L.) *The Pharma Innovation*; 12(1):671-673.

- Pandey, O.P., Shahi, S., Dubey, A. and Maurya S., 2019. Effect of integrated management of growth and yield attributes of green gram (*Vigna radiata* L.). *J. of pharmacognosy and phytochemistry*; 8(3): 2347-2352.
- Patel, H.B., Shah, K.A. and Patel, K.H., 2019. Response of green gram to different level of phosphorus and organic liquid fertilizer on yield, quality and nutrient content and uptake. *Int.J.Curr.Microbiol.App.Sci.*; 8(4): 1234-1242.
- Perli, V.H., Tomar S.S., Kasi R Gummadala, Deepsikha and Madhav Kaushik ,2022. Influence of organic manures and Bio-fertilizers on the growth and yield of green gram (*Vigna radiata* L.). *The Pharma Innovation Journal* 11(6): 141-144.
- Prajapati, S., 2014. Effect of integrated nutrient management on growth, yield, quality and economics of summer green gram [*Vigna radiata*]. (Master's thesis, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, India).
- Priyadarshini, A.S., Singh V., Tiwari D., Karthik B. and Mahesh K., 2021. Influence of Spacing and Organic manures on Growth, Yield and Economics of Mung Bean (*Vigna radiata* L.). *Biological Forum* 13(1): 617-621.
- Pushpa, K. Pruthviraj, N. and Vinay, M.G., 2022. Integrated nutrient management for improving growth and yield of cowpea. *Journal of Crop and Weed*, 18(3): 26-30.
- Rajeshwari, H. J., 2011. Integrated nutrient management in black gram (*Vigna mungo* L.) in northern transition zone of Karnataka. (Master's thesis, University of Agricultural Sciences, Dharwad, India).
- Ravikumar, M., Ventesh J., Nagarjuna K.S., 2012. Effect of nitrogen fixing bacteria in combination with organic manures and fertilizers on dry matter production and iptake of N, P and K in *Coleus forskohili*. *Green gram* 3: 637-641.

- Verma, R., Singh M., Dawson J and Pathan M. I. K., 2022. Effect of bio-fertilizers and organic manures on growth and yield of green gram (*Vigna radiata* L.). *The Pharma Innovation Journal* 11(4): 1599-1602.
- Saravanan, P.S., Singh, K., Ignesh, A., 2013. Effect of organic manures and chemical fertilizers on yield and macro nutrient concentration of green gram. *International Journal of Pharmaceutical Science Innovation*; 2(1): 18-20.
- Sengupta, K. and Tamang D., 2015. Response of green gram to foliar application of nutrients and brassinolide. *Journal Crop and Weed*, 11(1):43-45.
- Shariff, A. F., Sajjan, A. S., Babalad, H. B., Nagaraj, L. B., and Palankar, S. G., 2017. Effect of organics on seed yield and quality of green gram (*Vigna radiata* L.). *Legume Research- An International Journal*, 40 (2), 388-392.
- Shete P.G., Tanki J.D., Adhav S.L., Kushare Y.M., 2010. Response of *rabi* green gram (*Vigna radiata* L.) to land configuration and inorganic fertilizer with and without FYM. *Crop Res.*, 39: 43-46.
- Singh, M., Mishra, J. S., and Bhatt, B. P., 2017. Effect of integrated nutrient management on production potential and quality of summer mung bean (*Vigna radiata* L.). *Journal of Krishi Vigyan*, 5 (2), 39-45.
- Singh, R. K., Singh, S. R. K. Singh, Tripathi U, Srivastava V. and Rai H. K., 2021. Enhance the productivity and nutrient use efficiency of green gram (*Vigna radiata*) through inorganic nutrients and biofertilizers. *Indian Journal of Agricultural Sciences* 91(3): 417-420.
- Tripathi, A. and Tiwari, A.S., 2022. Integrated nutrient management of growth, yield and quality of green gram (*Vigna radiata* L.). *The pharma Innovation journal*;11(7): 2480-2482.

Yadav, K.V., Singh, D.P., Sharma, S.K. and Kishor, K. 2017. Use of phosphorus for maximization of summer green gram [*Vigna radiata* (L.) Wilszeck] productivity under sub humid condition of Rajasthan, India. Journal of Pharmacognosy and Phytochemistry. 6(4): 01-03.

UNDER PEER REVIEW