

Stability analyses of pearl millet genotypes (*Pennisetum glaucum*) and its related traits using the Freeman and Perkins model

Abstract

In the present study, twenty-nine genotypes were carried out to determine the stability of morphological characters viz., dry fodder yield, Days to 50% flowering, and Days to maturity of pearl millet in eight different environments viz., ARS Mandor, Bikaner, RARI Jaipur, Jamnagar, Hisar, Gwalior, Ranchi and Jammu Kashmir using Freeman and Perkins model. Data was collected from ICAR- All India Coordinated Research Projects (AICRP) of pearl millet, Agricultural University, Mandor, Rajasthan, India for one year (2019). The combined analysis of variance showed significant differences among genotypes, environments and Genotype by Environment (G x E) interaction for all morphological characters under study. Mean, regression coefficient and deviation from regression were used as stability measures. Genotype G23 exhibited stability across all studied traits, while G10 and G29 demonstrated stability specifically for dry fodder yield and days to 50% flowering. Furthermore, genotype G22 displayed stability for dry fodder yield and days to maturity in all environments.

Keywords: pearl millet, genotype, environment and G x E interaction.

Introduction

Pearl millet (*Pennisetum glaucum*) is a vital cereal crop known for its adaptability to harsh growing conditions. It is cultivated globally, particularly in arid and semi-arid regions. Pearl millet ranks as the sixth most important cereal crop worldwide, due to its ability to thrive in hot and dry climates (Gaoh et al., 2023). In India, pearl millet holds great significance and is grown extensively in states such as Rajasthan, Gujarat, Haryana, and Uttar Pradesh. This crop is not only a staple food source but also serves as livestock fodder. With its high nutritional content and gluten-free nature, pearl millet has gained attention for its suitability in addressing nutritional challenges and dietary restrictions. Ongoing research and breeding programs aim to enhance pearl millet's productivity, nutritional value, and stress tolerance. These efforts contribute to achieving food security and sustainable agriculture in regions facing climate variability and resource constraints.

Stability analysis plays a crucial role in agricultural research, particularly in crop improvement programs. It involves assessing the performance and adaptability of different genotypes under varying environmental conditions. This analysis helps researchers and farmers identify genotypes that exhibit consistent performance across diverse environments, ensuring reliable crop production and minimizing yield fluctuations. "The stability is the consistency in the performance of genotypes over a wide range of environments" (Singh and Chaudhary, 1985). For farmers, stability analysis provides valuable insights into the performance of crop genotypes across different agro-climatic conditions. It helps them make informed decisions regarding genotype selection, thereby increasing their chances of obtaining stable and high-yielding crops. In such circumstances, the identification of genotypes that exhibit stable performance across diverse environments or locations becomes crucial. The genotypes x environment interaction, as defined by Allard and Bradshaw in 1964, plays a significant role in genotype development and evaluation. This interaction is particularly important as diverse

environments can pose challenges and potentially decrease the stability of genotypes (Eberhart and Russell, 1966). It is widely observed that the performance of different genotypes varies across different environments, indicating the presence of genotype-environment interaction. Several studies, such as those conducted by Yates and Cochran (1938), Finlay and Wilkinson (1963), Rowe and Andrew (1964), Eberhart and Russell (1966), Perkins and Jinks (1968), Breese (1969), and Baker (1969), have further noted that the relationship between genotype performance and environmental factors often exhibits linearity or near-linearity. However, there are two fundamental statistical concerns raised regarding the aforementioned studies. These concerns include the selection of appropriate sums of squares and degrees of freedom to account for regression components, as well as the choice of the environmental measurement used for regression analysis. To address these issues, Freeman and Perkins (1971) developed a model that resolves these statistical challenges.

The experiment exhibited significant results due to environmental variations using Freeman and Perkins model in four rabi seasons of 2009-10 to 2012-13 on eight genotypes with yield and yield components (Husan and Deb, 2017). Eberhart and Russell (1966) and Freeman and Perkins model were used to study the twenty-five genotypes of wheat across eight locations with seven multiple traits (Said et al, 2020). Sokoto et al. (2023) conducted a scientific study in the Sudan savanna zone of Nigeria to assess the influence of three factors such as intra-row spacing, organic and inorganic fertilizers, and millet variety on millet yield.

The aim of this study was to assess the stability for Dry fodder yield (kg/net plot), Days to 50% flowering (days) and Days to maturity (days) of pearl millet genotypes across different environments using the Freeman and Perkins model. Mean, regression coefficient and deviation from regression were used as stability measures.

Material and Methods

Data set:

The analysis of the multi-environment experiment of twenty-nine pearl millet genotypes namely, HHB272 (G₁), MPMH21(G₂), RHB177(G₃), HHB197(G₄), GHB538(G₅), HHB67(G₆), AHB1269(G₇), HHB299(G₈), AHB1200(G₉), PB1705(G₁₀), XMT1497(G₁₁), 86M01(G₁₂), GHB905(G₁₃), MPMH17(G₁₄), RHB173(G₁₅), HHB223(G₁₆), GHB744(G₁₇), GHB732(G₁₈), KBH108(G₁₉), 86M86(G₂₀), Kaveri Super Boss(G₂₁), MP-7792(G₂₂), Proagro9444(G₂₃), GHB558(G₂₄), Dhanshakti(G₂₅), ICMV221(G₂₆), PusaComposite701(G₂₇), Posa Composite383(G₂₈), JBV 2(G₂₉) conducted at eight different locations of India viz, ARS Mandor, Bikaner, RARI Jaipur, Jamnagar, Hisar, Gwalior, Ranchi, Jammu Kashmir for one year (2019). The Randomized Block Design (RBD) with three replication was used. The data were recorded on Dry fodder yield (kg/net plot), Days to 50% flowering (days) and Days to maturity (days). The net plot size was 14.4 m² in four locations viz., ARS Mandor, Bikaner, Jamnagar, Jammu Kashmir and 12 m² in other four locations viz., RARI Jaipur, Hisar, Gwalior, Ranchi. Freeman and Perkins model (1971) was used to determine the stability of pearl millet genotypes.

Freeman and Perkins model:

This method ensures that the sum of squares resulting from various components must be correctly divided where environment taken as independent component.

The statistical analysis was based on the model in equation (1) as follows,

$$Y_{ij} = \mu + d_i + \bar{B}Z_j + \bar{\delta}_{ij} + B_{di}Z_j + \delta_{dij} + e_{ij} \quad (1)$$

Z_j is the environmental index of *j*th environment, μ is the grand mean, B_i is regression coefficient of *i*th genotype for regression of Y_{ij} on Z_j, \bar{B} is the combined regression coefficient, B_{di} is the difference between regression coefficient of *i*th genotype and combined regression coefficient, δ_{ij} is the

Deviation of i^{th} genotype from its linear regression on j^{th} environment, $\bar{\delta}_j$ is the Deviation from the combined regression line of the mean of all genotypes in the j^{th} environment (i.e., $(e_{ij} - \bar{B}Z_j)$), δ_{dij} is the deviation of the i^{th} genotype from each linear regression on Z_j in the j^{th} environment minus $\bar{\delta}_j$ (i.e., $\delta_{ij} - \bar{\delta}_j$), Such that, $\sum d_i = \sum Z_j = \sum B_{di} = \sum \bar{\delta}_j = \sum \delta_{dij} = 0$; ($i = 1, 2, \dots, g$ and $j = 1, 2, \dots, e$). The regression stability parameters of model (b^{Fi}) and $[\bar{S}_{di(F)}^2]$ were calculated as in equation 2 and 3 respectively.

$$b^{Fi} = \sum_i \sum_j \bar{y}_{ij} / \sum_i \sum_j Z_{ij}^2 \text{ and } Z_j = \bar{y}_j - \bar{y}.. \quad (2)$$

Where, \bar{y}_j is the mean of genotypes in j^{th} environment

$$[\bar{S}_{di(F)}^2] = [\sum_j \delta_{ij}^2 / (E - 2)] - (S_e^2 / r) s \quad (3)$$

$$\sum_j \delta_{ij}^2 = \sigma_{vi}^2 - b^{Fi} \sum_j \bar{y}_{ij} Z_j$$

$$\widehat{\sigma_{vi}^2} = \sum_j y_{ij}^2 = \sum_j y_{i.}^2 / E$$

Where, $\bar{y}..$ is the general mean, \bar{S}_e^2 is the Pooled EMS, $\widehat{\sigma_{vi}^2}$ is the S.S due to i^{th} genotype

Result and Discussion

The pooled analysis of variance (Table 1) based on eight different environments showed significant differences ($p \leq 0.01$) among genotypes, environments for dry fodder yield (kg/net plot), days to 50% flowering (days) and days to maturity (days). The genotype x environment interaction also showed highly significant differences for all studied characters under study. That indicates genotypes differed not only genetically, but also that some of them displayed diverse reactions to the dynamic environments. Then, various stability measures may be used to find stable genotypes. Similar results of significant differences among genotypes and environments were also reported by Alemu *et al.* (2018) and Haydar *et al.* (2018).

Table 1:- Pooled analysis of variance of different attributes of Pearl millet.

Sources	df	Mean sum of square		
		Dry fodder yield	Day to 50% flowering	Days to maturity
Genotypes	28	21.77**	240.04**	139.36**
Environments	7	999.34**	1774.63**	5185.92**
Rep. in Env.	16	3.75	8.49	4.56
Gen. x Env.	196	8.86**	11.46**	17.10**
Error	464	0.83	2.46	4.27

*,** significant at 5% and 1% prob. level, resp.

The partitioning analysis of variance model of Freeman and Perkins was conducted for traits under study showed in Table 2. It was noticed that the mean squares due to genotype showed significance for dry fodder yield (kg/net plot), days to 50% flowering (days) and days to maturity (days). Moreover, significant variations were observed in all three studied characters. Significant combined regression indicated that environments were well measured. Residual-1 item, when compared to the error was significant. Suggesting that the environmental index adequately captured to the error was significant. Suggesting that the environmental index adequately captured the index of additive environmental effect. G x E interaction was significant for a studied character, indicating that pearl millet genotypes interacted with the environment differently. The significant interaction of genotypes with environments warrants further computations of stability parameters. The heterogeneity of regression was found to be significant. On the other hand, residual-2 was significant, which indicates that the studied characters showed linear performance in the environments in which it was grown. Similar finding were reported by Sowmya *et al.* (2018).

Table 2: - Analysis of variance of pearl millet genotypes using Freeman and Perkins model (1971).

Sources	df	Mean sum of square		
		Dry fodder yield	Days to 50% flowering	Days to maturity
Genotypes (G)	28	12.97**	168.70**	94.05**
Environment (E)	7	648.89**	1180.33**	3497.00**
Combined regression	1	4430.86**	8109.88**	24413.07**
Residual (1)	6	18.56**	25.41**	10.99**
G x E interaction	196	6.10**	8.71**	13.63**
Heterogeneity of reg.	28	6.15**	18.26**	15.42**
Residual (2)	168	6.09**	7.12**	13.34**
Error between rep.	232	0.83	2.78	4.32

*, ** significant at 5% and 1% probability levels, respectively.

Stability analysis

Dry Fodder Yield

Three stability measures viz., mean, regression coefficient and mean square deviation were estimated for all the genotypes and its morphological characters under study for analysis presented in Table 3. Based on Freeman and Perkin's model, a genotype is said to be stable if it has regression coefficient near to unity and mean square deviation near or equal to zero. High value of regression ($b^{Fi} > 1$) indicates that the genotype/ variety is more responsive for input rich (favourable) environment, while low value of regression ($b^{Fi} < 1$), is an indication that the variety may be adopted in poor (unfavourable) environment.

Mean values of dry fodder yield ranged from 5.76 kg/net plot (G_{17}) to 8.93 kg/net plot (G_{10}) and the regression coefficient of both genotypes was quite equal but G_{17} has highest mean square deviation (S_{di}^2) among genotypes after G_9 . The genotypes G_7 , G_{10} , G_{11} , G_{12} , G_{14} , G_{15} , G_{18} , G_{19} , G_{21} , G_{22} , G_{23} , G_{28} and G_{29} had higher mean values than the overall mean. G_2 , G_4 , G_6 and G_{19} genotypes have unity regression coefficients indicating that these can be adapted to all environments, and G_3 , G_{10} , G_{12} , G_{13} , G_{16} , G_{17} , G_{18} and G_{28} may be adopted in favourable environments, while the remaining may be adopted in unfavourable (poor) environments. Therefore, the genotypes G_{10} , G_{14} , G_{22} , G_{23} and G_{29} have high mean value than the overall mean, unity regression coefficient and non-significant deviation from regression. These results agreed with Thakur *et al.*, 2019

Days to 50% flowering

The mean value for days to 50% flowering (days) ranged from 34.83 (G_6) to 44.80 (G_{21}) and the regression coefficient of both genotypes have less than unity indicating that they may be responsive to poor environments but the G_{21} had the highest mean square deviation. The genotypes G_7 , G_8 , G_{10} , G_{11} , G_{12} , G_{15} , G_{18} , G_{19} , G_{20} , G_{21} , G_{22} , G_{23} , G_{27} , G_{28} and G_{29} had higher mean values than the overall mean. The genotypes G_7 , G_8 , G_{20} , G_{24} and G_{29} have regression coefficients equal to one indicating that these can be adapted to all environments, and G_4 , G_9 , G_{12} , G_{14} , G_{15} , G_{16} , G_{17} , G_{18} , G_{22} , G_{23} , G_{27} and G_{28} may be adapted in favourable environments, while the remaining may be adopted in unfavourable (poor) environments. Meanwhile, the genotypes G_6 , G_{10} , G_{12} , G_{15} , G_{17} , G_{20} , G_{23} , G_{24} and G_{29} have high mean values than the overall mean, unity regression coefficient and non-significant deviation from regression.

Days to maturity

The mean value for days to maturity (days) ranged from 61.67 (G_6) to 69.37 (G_{19}) and the regression coefficient of both genotypes have unity indicating that these may have adapted to all environments. The genotypes G_7 , G_{11} , G_{12} , G_{17} , G_{18} , G_{19} , G_{20} , G_{21} , G_{22} , G_{23} , G_{27} and G_{29} had higher mean values than overall mean. G_1 , G_4 , G_5 , G_9 , G_{24} and G_{25} genotypes have regression coefficient equal

to one indicating that these genotypes adapted to all environments, and genotypes G3, G6, G7, G14, G15, G16, G17, G18, G19, G22, G23 and G26 have ($b^{Fi} > 1$) indicating genotypes may be adapted to favourable environments, while the remaining may be adapted in unfavourable environments. Therefore, the genotypes G1, G2, G4, G5, G7, G9, G22 and G23 found the most stable genotypes among all genotypes with mean square deviation near to zero.

Table 3: - Estimates of stability measures for pearl millet genotypes in different environments of India.

Varieties	Dry fodder yield			Days to 50% flowering			Days to maturity		
	mean	b^{Fi}	S_{di}^2	mean	b^{Fi}	S_{di}^2	mean	b^{Fi}	S_{di}^2
G ₁	6.33	0.89	2.65**	36.33	0.77	-1.51	62.43	1.07	-1.53
G ₂	6.38	1.00	2.36**	37.30	0.50	-0.63	63.03	0.84	0.51
G ₃	6.46	1.05	0.56	35.10	0.56	-1.38	61.83	1.12	5.85**
G ₄	6.72	1.02	1.83**	37.77	1.19	2.15*	62.63	1.00	3.10*
G ₅	6.16	0.93	0.73*	36.67	0.67	1.40*	61.93	1.03	1.31
G ₆	6.49	1.04	-0.29	34.83	0.79	0.01	61.67	1.09	4.60*
G ₇	7.05	0.77	3.15**	41.00	1.04	2.20*	65.57	1.12	0.66
G ₈	6.45	0.91	0.92*	39.70	1.01	2.84*	64.40	0.96	-1.98
G ₉	6.41	0.50	5.26**	39.27	1.08	-1.99	64.20	1.01	0.42
G ₁₀	8.93	1.25	0.37	40.60	0.91	0.13	64.00	0.83	0.14
G ₁₁	7.71	0.90	1.16**	40.17	0.56	-0.53	64.43	0.67	2.22
G ₁₂	7.80	1.11	2.21**	41.50	1.11	-1.38	65.10	0.94	-2.38
G ₁₃	6.76	1.18	0.44	38.57	0.82	1.80*	63.97	0.98	-2.25
G ₁₄	7.08	0.98	0.41	37.77	1.16	-1.78	63.67	1.14	5.11**
G ₁₅	7.11	0.83	0.65*	39.50	1.21	-0.23	64.07	1.12	-0.47
G ₁₆	6.39	1.13	2.67**	37.40	1.16	-0.72	62.43	1.09	-2.85
G ₁₇	5.76	1.24	3.94**	41.00	1.18	0.54	66.17	1.14	1.60
G ₁₈	7.16	1.18	1.42**	40.57	1.22	-2.00	65.20	1.15	-1.15
G ₁₉	8.44	1.04	3.44**	44.47	0.95	2.10*	69.37	1.08	-1.93
G ₂₀	6.83	0.57	3.78**	43.37	1.00	-1.54	67.00	0.90	1.43
G ₂₁	8.23	0.88	3.39**	44.80	0.91	5.79**	68.73	0.87	1.76
G ₂₂	7.28	0.85	0.38	41.27	1.19	-2.52	66.43	1.24	0.69
G ₂₃	7.04	0.75	0.50	40.50	1.30	-0.01	65.13	1.30	-0.09
G ₂₄	6.80	0.95	1.31**	39.37	1.05	0.39	64.70	1.05	3.17*
G ₂₅	5.78	0.91	0.66*	36.77	0.64	4.29**	61.70	1.02	2.44
G ₂₆	6.40	0.74	1.66**	36.87	0.70	1.12	63.20	1.10	0.51
G ₂₇	6.69	0.84	-0.22	40.17	1.27	-1.10	64.73	0.89	1.48
G ₂₈	7.84	1.18	1.07*	41.63	1.47	-2.03	65.13	0.90	-0.39
G ₂₉	7.46	0.71	0.43	40.63	1.05	-1.84	64.83	0.84	-2.99
Average	6.96			39.48			64.40		

*, ** significant at 5% and 1% probability level, respectively.

Conclusion

The results were found highly significant among genotypes, environments and genotype x environment interaction for all the morphological characters viz., dry fodder yield, days to 50% flowering and days to maturity. The genotypes G2, G4, G6 and G19 were responsive to all environments for dry fodder yield, G7, G8, G20, G24 and G29 genotypes were responsive to all environments for days to 50% flowering while, G1, G4, G5, G9, G24 and G25 genotypes may be adopted to all environments for days to maturity. The genotypes G10, G14, G22, G23 and G29 were found stable for dry fodder, G6, G10, G12, G15, G17, G20, G23, G24 and G29 genotypes were stable

for days to 50% flowering and G1, G2, G4, G5, G7, G9, G22 and G23 genotypes were stable for days to maturity. The genotype G23 was observed stable for all the morphological characters.

References

- Alemu, G., Mohammed, H. and Asnake, D. (2018). Analysis of genotype by environment interaction for agronomic traits of bread wheat (*Triticum aestivum* L) genotype in Ethiopia. *Journal Agricultural Research*, 3(8), 1-11. <https://doi.org/10.23880/OAJAR-16000191>
- Allard, R.W. and Bradshaw, A.D. (1964). Implications of genotype-environment interactions in applied plant breeding. *Crop Science*, 4, 503-508.
- Baker, IT. (1969). Genotype-environment interactions in yield of wheat. *Canadian Journal Plant Science*, 49(6), 743-751.
- Breese, E. L. (1969). The measurement and significance of genotype-environment interactions in grasses. *Heredity*, 24, 27-44.
- Eberhart, S.A. and Russell, W.A. (1966). Stability parameters for comparing varieties. *Crop Science*, 6, 36-40.
- Finlay, K. W., and Wilkinson, G. N. (1963). The analysis of adaptation in a plant breeding programme. *Aust. J. Agric. Res.*, 14, 742-754.
- Freeman, G.H. and Perkins, J.M. (1971). Environmental and Genotype-environmental components of variability VIII. Relations between genotypes grown in different environments and measures of these environments. *Heredity*, 27: 15-23.
- Gaoh BS, Gangashetty PI, Mohammed R, Ango IK, Dzidzienyo DK, Tongoona P, Govindaraj M. (2023). Combining ability studies of grain Fe and Zn contents of pearl millet (*Pennisetum glaucum* L.) in West Africa. *Frontiers in Plant Science*. 1-17.
- Haydar, F.M. A., M.N. Uddin, U.K. Roy, G.M. Uddin and A.B. Siddique. (2018). Stability analysis of yield performance in wheat (*Triticum aestivum* L.). *Plant Environ. Dev.* 7(2): 10-15.
- Husan, M.T. and Deb, A.C. (2017) Stability Analysis of Yield and Yield Components in Chickpea (*Cicer Arietinum* L.). *Horticulture International Journal*, 1(1), 4-14.
- Perkins, Jean M., and Jinks, J. L. (1968). Environmental and genotype-environmental components of variability. III. Multiple lines and crosses. *Heredity*, 23, 3 39-356.
- Rowe, P. K. and Andrew, IT. A. (1964). Phenotypic stability for a systematic series of corn genotypes. *Crop Sri.*, 4, 563-564.
- Said, A.A., Motawea, M.H. and Morsy, H (2020). A comparison of bread wheat genotypes in various environments using different stability models. *International Journal of Agricultural Science*, 2(2), 340-355.
- Singh RK and Chaudhary BD. (1985). *Biometrical methods in Quantitative Genetic Analysis*. Kalyani Publications, New Delhi.
- Sokoto MB, Aigbojie EO, Dikko AU, Singh A, Aliyu U, Lukman SA, Sharif RM. (2023) Productivity of Millet (*Pennisetum glaucum* (L) R. BR.) as Influenced by Intra-row Spacing, Low Levels of Combined Organic and In-organic Fertilizers and Variety in Sokoto, Sudan Savannah, Nigeria. *International Journal of Plant & Soil Science*, 35(17),132-48.

Sowmya HH, Kamatar MY, Shantha kumar G, Brunda SM, Shadakshari TV, Showkath Babu BM, Sanjeev Singh Rajput (2018). Stability analysis of maize hybrids using Eberhart and Russel Model. *International Journal of Current Microbiology and Applied Sciences*, 7(2), 3336-3343.

Thakur, P., Prasad, L.C., Prasad, R., Omprakash, Chandra, K. and Rashmi, K. (2019). Stability analysis for yield and related traits over four environments in wheat (*Triticum aestivum* L.). *Plant Archieve*, 19(2), 3541-3545.

Yates, P. and Cochran, W. G. (1938). The analysis of groups of experiments. *Journal agricultural Science*, **28**, 556-580.