

In-silico screening of lung cancer inhibiting potential of the chemical Constituents of n-hexane extract of *Elaisiguineenses*

Abstract

Palm oil is an edible vegetable oil that is rich in saturated fat, it is derived from the mesocarp of the oil palm fruit especially the African oil palm fruit (*Elaisiguineenses*), this study investigates the possible inhibition of lung cancer cells using the bioactive constituents of *Elaisiguineenses*. The chemical constituents of the oil were extracted in n-hexane, analyzed with Fourier transform infrared spectrometer (FTIR) and gas chromatography-mass spectrometry (GC-MS) studies. Absorption, distribution, metabolism and elimination and toxicity (ADMET) properties were predicted using ADMETSAR online software while the molecular docking was achieved using the AutodockVina found in PYRX software. The ADMET properties of the constituents showed that they are good drug leads for lung cancer treatment. The docking result revealed that 1,4,10,13-tetraoxa-7,16-dithiacyclooctadecane gave a good binding affinity ($-5.8 \text{ kcal mol}^{-1}$) that is close to that of the control (Osimertinib) ($-7.8 \text{ kcal mol}^{-1}$). The obtained result showed that palm oil may be a promising lead candidate for the treatment of lung cancer.

Keywords: lung cancer, *Elaisiguineenses*, osimertinib, molecular docking, PYRX,

1. Introduction

Cancer is one of the most deadly diseases known in developing African countries especially Nigeria (1-2). Cancer has limited chances for curing it, especially when it goes beyond the early stage. It occurs as a result of biochemical alteration in the genetic and epigenetic mechanisms while targeting many tracts in growth and malignant progression towards incurable lethal disease (3). Most cancer treatment methods especially chemotherapeutic agents offer some level of toxicity to the patient showing severe side effects. Natural anticancer products have proven to be reliable when it comes to the management of the disease (4-6). Lung cancer is one of the major life threatening diseases in Nigeria today, this may be attributed to the careless life style of people living in this area. Majority of the young men smoke

tobacco and hard drugs. Molecular docking is a method used to model the interaction between natural compounds and target protein (7-9), The binding site area between the amino acid residues plays an important role in binding with drugs or ligands (10). Molecular docking correlates the computational and experimental approaches and has been of great assistance in the identification and development of promising and novel compounds (11-12). West Africa Palm oil (*Elaeis guineensis*) is one of the most beneficial food products in the world today. It's used for preparing meals and also as an ingredient in ice cream, margarine, and many other foods products. Palm oil is made from the fruit of the African oil palm. It is used traditionally for the control of rheumatism headaches, pains, arterial thrombosis, cancer, cardiovascular diseases and an atherosclerosis because of the very rich phytonutrients it contains (13-14). The biomass extracted from palm oil industries which include oil palm frond, oil palm leaves, oil palm trunks, mesocarp fibers, empty fruit bunches, palm kernel shells, and palm oil mill are poorly utilized (15-16). This work tries to identify some natural lung cancer inhibitors from palm oil through molecular docking method.

2.0 Materials and Methods

2.1 Collection of plant Materials

Fresh palm fruits were harvested from a local village in Imo State Nigeria and were identified by Prof. Martins Mbagwu of the Plant Science and Biotechnology department Imo State University, Owerri. After proper identification and authentication, the fruits were ground with care so that the hard nut does not crack. The nut was separated from the fiber and the fiber was subsequently dipped in n-hexane and allowed to stand for 24 hours after which the oil was filtered and stored in an air tight container for further use (17).

2.2 FTIR Analysis

Fourier transform infrared analysis was conducted on the extract to identify the functional groups therein. The analysis was carried out at a frequency range of 4000-400 cm^{-1} . The extracted oil was encapsulated in 200 mg of KBr salt pellet, using a mortar and pestle, compressed into a thin pellet, the sample scans were set at 30 to calibrate it. Resolution was set to 8.2 and the Background Scans were set at 16 (18)

2.3 Gas-Mass Chromatography Analysis

The extracted oil sample was subjected to GC-MS experiment using an Agilent technology gas-mass chromatography instrument (model: 19091S-433UI) USA. The conditions for the gas chromatography part were set with the parameter as follows: A HP-5ms capillary standard nonpolar column Ultra Inert 0 °C—325 °C (350 °C): 30 m x 250 µm x 0.25 µm was used. The temperature range was 50 – 325°C, held at 50 °C for 2 min, the rate was 5 °C/*min* and at 180 °C for 0 minutes increased to 20 °C/*min* at 270 °C for 5 min, the carrier gas used was helium at a rate of 1 mL per minute. The results obtained were analyzed by comparing with the spectra program imbibed in the National Institute of Standards and Technology (NIST) mass spectral library (19).

2.4 Ligands preparation and identification

The 3D structures of the compounds identified in the GC-MS experiment were downloaded from the [pubchem online database](#). The obtained compounds were minimized at PyRx virtual screening tool with the aid of universal force field with 200 steps and changed to AutoDock ligands and used for the molecular docking procedures (20).

2.5. Absorption, distribution, metabolism and elimination and toxicity (ADMET) screening

The identified compounds were submitted to ADMETSar2 server for drug-like properties, pharmacokinetics, and pharmacodynamics parameter examination (21).

2.6. Identification and preparation of molecular targets

The 3D structure of epidermal growth factor receptor protein target was downloaded from protein data bank ([PDB database](#)) with PDB ID: 4zau. The protein was prepared using discovery studio where the interfering crystallographic water molecules and co-crystallized ligand were removed, the active site was discovered using pymol software. Further preparation was done by addition of polar hydrogen. The prepared protein was saved as protein data bank file and used for the molecular docking [analysis](#)

2.7 Docking Analysis

Docking studies of the compounds on the prepared protein was achieved with the Autodockvina imbibed in PYRX software. Depending on several scoring functions, the PYRX software allows one to virtually screen a library of compounds and detects the strongest binders. The ligand binding site of Osimertinib was chosen to be the active site of the 4zau protein. The results from this study showed that all the docked compounds bind either at the active site or very close to the site

(22).The **Selected** residues involved in covalent interaction around the active site of the chosen protein are Leu718, Val 726, Ala 743, Rhu 723, Thr 790, Pro 794 and Cys 797. The docked results were subsequently visualized in discovery studies for further analysis.




Figure 1. 3 D view of (a) 4zau protein and (b) Osimertinib(Atom legend: white = H; gray = C; red = O; blue = N.)

3.0 Results

3.1 FTIR Result

Ftir analysis was used to identify the functional groups present in the **n**-Hexane extract *Elaisiguineenses*. The presence of some functional groups was observed. The major functional groups are summarized in Table 1 while the spectrum is shown in Figure 2. The functional groups observed is similar to those reported elsewhere **to enhance drug activity (23)**

Table 1 Major functional groups found in n-Hexane extract of *Elaisiguineenses*

Absorption peak (cm^{-1})	Functional group	Appearance
3008.0	O-H of carboxylic acid	Weak Broad

2922.2	Asymmetric sp^3 C-H	Strong sharp
2855.1	Symmetric Sp^3 C-H	H Strong sharp
1744.4, 1710.8	C = O	Strong sharp
1481.1	C – H	Sharp




Figure 2. FTIR spectra of the n-hexane extract of *Elaisiguineenses*

3.2 GC-MS Result

The n-Hexane extracted oil was injected into an Agilent Gas chromatography-mass spectrophotometer (model 19091S-433UI) for examination and analysis. The chromatogram is presented in Figure 3 while Table 2 gives the compounds identified from the GC-MS analysis. N-(2-[(2-(dimethylamino)ethyl)(methyl)amino]-4-methoxy-5-[4-(1-methyl-1H-indol-3-yl)pyrimidin-2-yl]amino}phenyl)prop-2-enamide (**Osimertinib**) with Pubchem Id 71496458, molecular formula $C_{28}H_{33}N_7O_2$ and weight 499.6 g/mol was used

as control drug to identify compounds with good binding affinities for the studied cancer protein




Figure 3 Chromatogram of the n-hexane extract of *Elaisiguineenses*

Table 2 **phytochemical** identified in the n-hexane extract of *Elaisiguineenses* oil

S. No.	Compound	Area Peak	Pubchem Id	Structure	Molecular formula	Molar weight (g/mol)
1.	cis-Vaccenic acid	28.5039	5282761		$C_{18}H_{34}O_2$	282.50
2.	13-Octadecenal, (Z)-	3.1848	5364497		$C_{18}H_{34}O$	266.50
3.	Palmitic acid	6.6054	985		$C_{16}H_{32}O_2$	256.42
5	Oleic Acid	27.5039	445639		$C_{18}H_{34}O_2$	282.50
6	trans-13-Octadecenoic acid	4.1679	6161490		$C_{18}H_{34}O_2$	282.50
7	9-Octadecenal, (Z)-	4.8017	5364492		$C_{18}H_{34}O$	266.50
8	Hexadecenoic acid, Z-11-	8.983	5312414		$C_{16}H_{30}O_2$	254.41
9	1,4,10,13-tetraoxa-7,16-dithiacyclooctadecane	9.0417	67527		$C_{12}H_{24}O_4S_2$	296.5
10	Z-2-Octadecen-1-ol	7.2167	5365011		$C_{18}H_{36}O$	268.5

3.3. ADMET behavior

The **Absorption**, distribution, metabolism and elimination and toxicity prediction which show their pharmacokinetics and pharmacodynamics properties are summarized in Table 3. **pfizer's rule** of five by Lipinski (24) was used to evaluate the drug-likeness of the identified compounds. The rule helps to determine if a compound with given pharmacological activity has chemical and physical properties that would likely make it an orally active drugs in humans. In compliance to the rule, none of the compound has more than 5 hydrogen bond donor, none has more than 10 hydrogen bond acceptors, Table 2 showed that all have molecular mass of (**< 500**). According to Lipinski, an orally active drug should not violate more than one of the rules. The result revealed that all the compounds studied were in compliance with the rule, the higher human intestinal absorption (HIA) results indicated that the compounds may be better consumed from the gastrointestinal tract upon oral administration, none of the compounds was found to be carcinogenic. With the observed results the compounds may be classified as good drug candidates.

Table 3. ADMET characteristics of the docked compounds











S/N	Compound	HI	AO	WS	H-bond acceptor	H-bond donor	Rota table bonds	Car cinogeni ety
1.	Osimertinib	+	2.89 3	- 3.549	8	2	10	-
2.	cis-Vaccenic acid	+	1.23 8	+ -4.04	1	1	15	-
3.	13-Octadecenal, (Z)-	+	1.53 5	- 2.949	1	0	15	-
4.	Palmitic acid	+	1.16	+ - 3.502	1	1	14	-
5.	Oleic Acid	+	1.24 6	+ -4.04	1	1	15	-
6.	trans-13-Octadecenoic acid	+	1.22 8	+ -3.791	1	1	15	-
7.	9-Octadecenal, (Z)-	+	1.22 8	- 3.791	1	1	15	-

8. Hexadecenoic acid, Z-11-	+	1.26 8	+	-3.791	1	13	-
9. 1,4,10,13-tetraoxa-7,16-dithiacyclooctadecane	+	1.85 3	+	-1.002	6	0	0
10. Z-2-Octadecen-1-ol	+	1.29 7	+	-2.349	1	1	15

HIA=Human intestine Absorption, AOT=Acute oral Toxicity, BBB=Blood Brain Barrier, WS=Water Solubility

3.4 Molecular docking result

Molecular docking was performed on the chosen protein using the identified compounds and a control (Osimertinib), the 3D and 2D protein–ligand interactions of the compounds are presented in Figure 4 while the amino acid residues involved in the interactions with the compounds and the control drug Osimertinib are shown in Table 4. The main forces involved in the interactions were Van der Waals, Conventional hydrogen bond, alkyl, unfavorable acceptor-acceptor. The forces of interaction and amino acid residues involved in each interaction are presented in Table 5, The results from the docking studies showed that all the compounds either bound exactly at active site or close to the active site and their binding scores were equally close to that of the control. 1,4,10,13-tetraoxa-7,16-dithiacyclooctadecane gave the closest binding score (-5.8 kcal mol⁻¹) to the control.

5	Oleic Acid	 <p>H-Bonds Donor Acceptor</p>	 <p>Interactions van der Waals Unfavorable Acceptor-Acceptor Alkyl</p>
6	trans-13-Octadecenoic acid	 <p>H-Bonds Donor Acceptor</p>	 <p>Interactions van der Waals Conventional Hydrogen Bond Alkyl</p>
7	9-Octadecenal, (Z)-	 <p>H-Bonds Donor Acceptor</p>	 <p>Interactions van der Waals Alkyl</p>
8	Hexadecenoic acid, Z-11-	 <p>H-Bonds Donor Acceptor</p>	 <p>Interactions van der Waals Conventional Hydrogen Bond Unfavorable Donor-Donor Alkyl</p>
9	1,4,10,13-tetraoxa-7,16-dithiacycloctadecane	 <p>H-Bonds Donor Acceptor</p>	 <p>Interactions van der Waals Carbon Hydrogen Bond Alkyl</p>




Figure 4 the 3D and 2D protein–ligand interactions

Table 4. **binding scores**, forces of interactions and amino acid residues involved in the docking studies

S. No.	Compound	Pubchem ID	BA	Types of interaction	amino acids involved
1.	Osimertinib	71496458	-7.6	Van der Waals, carbon hydrogen, pi-anion, pi-sigma, alkyl, pi-alkyl.	Leu 718, Val 726, Ala 743, Glu 762, Leu 792, Leu 844 Asp 855
2.	cis-Vaccenic acid	5282761	-5.1	Van der Waals, Conventional hydrogen bond, alkyl.	Val 726, Ala 743, Lys 745, Thr 790, Leu 792, Met 793, Thu 844,
3.	13-Octadecenal, (Z)-	5364497	-4.6	Van der Waals, Conventional hydrogen bond, alkyl.	Leu 718, Val 726, Ala 743, Lys 745, Leu 844, Thr 854
4.	n-Hexadecanoic acid (Palmitic acid)	985	-4.9	Van der Waals, Conventional hydrogen bond, alkyl.	Leu 718, Val 726, Ala 743 Lys 745, Leu 788, Thr 790, Leu 792, Leu 844, Met 793
5	Oleic Acid	445639	-5.0	Van der Waals,	Val 726, Ala 743, Lys 745, Thr 790,

				unfavorable acceptor-acceptor	Leu 844, Thr 854
6	trans-13-Octadecenoic acid	6161490	-5.3	Van der Waals, Conventional hydrogen bond, alkyl	Leu 718, Val 726, Ala 743, Leu 788, Thr 790, Leu 844
7	9-Octadecenal, (Z)-	5364492	-4.2	Van der Waals, alkyl	Leu 718, Val 726, Ala 743, Lys 745, Leu 792, Met 766, Leu 844
8	Hexadecenoic acid, Z-11-	5312414	-5.4	Van der Waals, Conventional hydrogen bond, alkyl, unfavorable acceptor-acceptor	Leu 718, Val 726, Ala 743, Cys 775, Thr 790, Leu 844, Lys 745
9	1,4,10,13-tetraoxa-7,16-dithiacyclooctadecane	67527	-5.8	Van der Waals, Conventional hydrogen bond, alkyl	Leu 718, Val 726, Met 793, Leu 844
10	Z-2-Octadecen-1-ol	5365011	-4.7	Van der Waals, Conventional hydrogen bond, alkyl	Leu 718, Val 726, Ala 743, Lys 793, Leu 788, Thr 854, Leu 844, Met 793

BA= Binding Affinity in (kcalmol⁻¹)

Conclusion

The chemical constituents of *Elaisiguineenses* were extracted in n-hexane, analyzed with FTIR and GC-MS. The FTIR result showed that the compound contained some functional groups which had been associated with drug properties and the GC-MS analysis revealed that it contained many compounds with known biological and therapeutic activities. ADMET properties showed that the constituents have good pharmacological and pharmacodynamics properties in the human system. The molecular docking study of the identified compound against the epidermal growth factor receptor lung cancer protein showed that the constituents of the African palm oil (*Elaisiguineenses*) were promising candidates in the inhibition of the activity of this protein.

References

1. Vikas J. et al., Screening of Phytochemicals as Potential Inhibitors of Breast Cancer using Structure Based Multitargeted Molecular Docking Analysis, *Phytomedicine Plus*, 2022: 2667-0313
<https://doi.org/10.1016/j.phyplu.2022.100227>.
2. Drescher C W, Bograd, A J, Chang S C, Weerasinghe R K, Vita A, Bell R, Cancer Case Trends Following the Onset of the COVID-19 Pandemic: A Community-Based Observational Study with Extended Follow-Up. *Cancer* 2022 ;128: 1475–1482.
3. Davis E J, Beebe-Dimmer J L, Yee C L, Cooney K A. Risk of Second Primary tumors in men diagnosed with prostate cancer: A Population-Based Cohort Study. *Cancer* 2014; 120: 2735–2741.
4. Taghizadeh MS, Niazi A, Moghadam A, Afsharifar A (2022) Experimental, molecular docking and molecular dynamic studies of natural products targeting overexpressed receptors in breast cancer. *PLoS ONE* 17; 5: e0267961. <https://doi.org/10.1371/journal.pone.0267961>.
5. Al-Amiery AA, Al-Majedy YK, Kadhum AAH, Mohamad AB (2015) Novel macromolecules derived from coumarin: synthesis and antioxidant activity. *Scientific Reports* 2015; 1 5:1–7. <https://doi.org/10.1038/srep11825>.
6. Sanket R. et al., Computational Exploration of Anti-cancer Potential of Flavonoids against Cyclin-Dependent Kinase 8: An InSilico Molecular Docking and Dynamic Approach, *ACS Omega* 2023; 8: 391–409, <https://doi.org/10.1021/acsomega.2c04837>.
7. Larionova I, Cherdyntseva N, Liu T, Patysheva M, Rakina M, Kzhyshkowska J, Interaction of tumor-associated macrophages and cancer chemotherapy. *Oncoimmunology*, 2019; 8: 1–15. <https://doi.org/10.1080/2162402X.2019.1596004>.

8. Lyskov S, Gray J., TheRosettaDock server for local protein-protein docking. *Nucleic Acids Research*. 2008;36: 233–238.
<https://doi.org/10.1093/nar/gkn216>.
9. Mace PD, Riedl SJ, Salvesen GS, Caspase enzymology and activation mechanisms. In: *Methods in Enzymology*. Elsevier Inc., 2014; 161–178.
<https://doi.org/10.1016/B978-0-12-417158-9.00007-8>.
10. Arwansyah I, Ambarsari L, Sumaryada TI, Simulasi Docking SenyawaKurkumindanAnalognyaSebagai Inhibitor Reseptor Androgen padaKankerProstat. *Current Biochemistry*2014; 1: 11–19.
<https://doi.org/10.29244/cb.1.1.11-19>.
11. Gong N, Wang L, An L and Xu Y, Exploring the active ingredients and potential mechanisms of action of sinomeniumacutum in the treatment of rheumatoid arthritis based on systems biology and network pharmacology. *Front. Molecular Bioscience*2023; 10:1065171. doi: 10.3389/fmolb.2023.1065171.
12. Awik P. D. et al. Anti-cancer potency by induced apoptosis by molecular docking P53, caspase, cyclin D1, cytotoxicity analysis and phagocytosis activity of trisindoline 1,3 and 4, *Saudi Pharmaceutical Journal* 2022;30:1345–1359.
13. Kamal-Eldin A, Appelqvist LÅ. The chemistry and antioxidant properties of tocopherols and tocotrienols. *Lipids*. 1996;3,17:671-701
14. Fattore E, Fanelli R, Palm oil and palmitic acid: a review on cardiovascular effects and carcinogenicity. *International journal of food sciences and nutrition*. 2013 ;164(5):648-59.
15. Romes N B, Hamid M A, Hashim S E, Wahab R A, Statistical modelling of ultrasonic-aided extraction of *Elaeisguineensis* leaves for better-quality yield and total phenolic content. *Indonesia Journal of Chemistry*, 2018;19(3):811–826.
16. Romes NB, Wahab RA, Hamid M A, Proximate analysis and bioactivity study on acoustically isolated *Elaeisguineensis* leaves extract. In: *AIP Conference Proceedings*. AIP Publishing LLC, Berlin, 2019; 20001
17. Duru CE Mineral and phytochemical evaluation of Zea mays husk. *SciAfr*7:e00224:2020;<https://doi.org/10.1016/j.sciaf.2019.e00224>.
18. Ikpa CCB and Maduka TOD. Antimicrobial Properties of Methanol Extract of Dacryodesedulis Seed and Determination of Phytochemical Composition Using FTIR and GCMS. *Chemistry Africa*. 2020: 3(4):927-935.
19. Iwu I C, Oze RN, Onu U L, Amarachi N, Ukaoma A, Phytochemical and gc/ms analysis of the rhizome of *Zingiberofficinale* plant grown in eastern part of Nigeria, *African Journal of Biology and Medical Research*, 2018;1: 1,43-54

20. Duru CE, Duru IA, Adegboyega AE, In silico identification of compounds from *Nigella sativa* seed oil as potential inhibitors of SARS-CoV-2 targets, *Bulletin of national research center*, 2021;45:57, <https://doi.org/10.1186/s42269-021-00517-x>.
21. Aprilita RY et al., **molecular docking** analysis of natural products from *Centella asiatica* for inhibition of renin, *Rasayan journal of chemistry*, 2023; 16: 2, 557-564.
22. Yosaatmadja Y. et al Binding mode of the breakthrough inhibitor AZD9291 to epidermal growth factor receptor revealed, *Journal of Structural Biology*, 2015; 1047-8477, <http://dx.doi.org/10.1016/j.jsb.2015.10.018>
23. Ikpa C B, Ikezu U J, Maduwuba M C, In Silico Docking Analysis of Anti-malaria and Anti-typhoid Potentials of Phytochemical Constituents of Ethanol Extract of *Dryopteris dilatata*, *Tropical Journal of Natural Product Research*, 2022; 6(5):772-782.
24. Lipinski CA (2016) Rule of five in 2015 and beyond: Target and ligand structural limitations, ligand chemistry structure and drug discovery project decisions, *Advanced Drug Delivery Review* 2015;101:34–41