

Original Research Article

Mycelium performance of medicinally important fungus (*Cordyceps militaris*) on different pH and temperature

ABSTRACT

Medicinal Mushrooms are most valuable fungi in the world. Artificial cultivation of these fungi is trending now days in the different part of world. Many factors are influence with fungus mycelium during its growth period. So the present study was undertaken with the aim of finding out most favourable pH and temperature for mycelia growth and dry matter weight of *Cordyceps militaris*. In this study a range of pH (5.5-8.5) and temperature (16 - 28°C) was used. The Optimum pH for mycelium growth and dry matter weight of *Cordyceps militaris* was 7.0 while optimum temperature for mycelium growth and dry matter weight was 22 °C.

KEYWORDS: *Cordyceps*, Mushroom, pH, temperature, mycelial growth, dry matter weight.

Comment [I1]: Change this key words to normal words don't write it italic

1. INTRODUCTION

There are at least 12,000 species of fungi that can be considered as mushrooms with at least 2000 species showing various degrees of edibility [3]. *Cordyceps militaris* (an entomopathogenic fungus), is one of the most important medicinal mushrooms, belonging to the class Ascomycetes, has been used popularly as a crude drug and a folk tonic food in East Asia [26]. The name comes from the Latin words: *cord* and *ceps*, meaning "club" and "head", respectively. It is named "soft gold" in China [25]. It possesses many kinds of active components (such as cordycepin, polysaccharides, ergosterol, and mannitol), and due to its several physiological activities, it is currently used for multiple medicinal purposes [18]. It is widely distributed in North America, South America, Europe, and Asia [17], from sub-tropical to temperate regions around the world. The main active constituent of *Cordyceps* fruiting bodies is cordycepin, which was first extracted from *C. militaris* and then found to be present in *Cordyceps sinensis* [2] and *Cordyceps kyushuensis* [14]. Cordycepin has a broad spectrum of biological activity, plays an important role in the treatment of respiratory and cerebrovascular diseases, enhancement of body immunomodulatory function and regulation of liver and renal metabolism [11]. Moreover, it also has been used as an anti-cancer [6], anti-tumor [20], anti-fungal [13], anti-hyperlipidemia [8], antioxidant [21], and anti-leukemia [24]. Cordycepin is also a Phase I/II clinical stage drug candidate for the treatment of refractory Acute Lymphoblastic Leukemia (ALL) patients who express enzyme terminal deoxynucleotidyl transferase (TdT). The natural fruiting bodies of *Cordyceps* are very rare and costly to collect. Fruiting body production *in vitro* is not repeatable and cordycepin content of natural *Cordyceps* is much lower than that of cultured mycelia [7]. In recent years *C. militaris* is extensively cultivated in liquid as well as solid media [4] and is the most successfully cultivated *Cordyceps* species [23]. Cultivation of *C. militaris* mycelium using artificial media [16] gave higher cordycepin yields. However, only a single *C. militaris* strain was employed and cordycepin production may vary with different strains.

Comment [I2]: Added paragraph about effect of pH and temperature on mushroom as introduction for your works

Keeping in view the above nutritional and medicinal importance of *C. militaris* mushroom and possibilities cultivating of medicinal mushrooms in the rural as well as urban areas of the country, present study has been done with the aim to examine the effect of different temperature and pH on the growth of *Cordyceps militaris*.

2. MATERIAL AND METHODS

2.1. Experimental site

The experiments were conducted in Mushroom Laboratory Department Plant of Pathology, S. V. P. University of Agriculture and Technology, Meerut, U.P. (India) during year 2018-20, which is situated on the Western side of the Delhi-Dehradun high way (NH-58) at distance of 10.0 km away in the north of Meerut city. The district Meerut is situated between 29° 01'N latitude and 77° 45'E longitude at an altitude of 237 meters above the mean sea level

2.2. Establishment of pure culture

Culture of *Cordyceps militaris* were purified and maintained by single hyphal tip method. For this purpose, the cultures were grown in sterilized Petri plate on Potato Dextrose Agar Medium (PDA) for 8-10 days. Single branched hyphae from the periphery of the growing colony were marked under low power (10x) in the compound microscope and transferred to PDA slants. These tubes were incubated at 21-24°C for about a week, again sub cultured on PDA and then stored in a refrigerator at 5-10°C for further use [5]. Optimizations of Culture Conditions on radial Growth of *Cordyceps militaris* given below:

Comment [I3]: 4-8 not 5-10

2.3. Effect of different pH on mycelial growth and dry matter weight.

For the studies of suitable pH, the culture of *Cordyceps militaris* were incubated at seven different pH media viz. 5.5, 6.0, 6.5, 7.0, 7.5, 8.0 and 8.5. Required pH of the culture media (PDA) was adjusted with N/10 solutions of NaOH or HCl, used before adding of agar, it was measured by a digital pH meter. After sterilization at 121°C (1.1 kg/cm² pressure) for 20 minutes in autoclave, the test media (PDA) were poured into the Petri plates (90 mm @ 20 ml/plate). The plates were inoculated centrally with a 9 mm diameter disc of 7 days old culture of *Cordyceps militaris* and incubated at 23±2°C in B.O.D with three replications of each treatment. The observations of radial growth were taken at each 48 hrs in *Cordyceps militaris* until the colony covered the full plate. The PD broth media was poured into 150 ml conical flasks @ 50 ml per flask and after adjusting of pH, broth sterilized in autoclave at 121°C (1.1 kg/cm² pressure) for 20 minutes and allowed to cool. The 9 mm diameter disc cut by cork borer from the periphery of 7 days old culture of *Cordyceps militaris* were inoculated in the flask with the help of sterilized inoculation needle. The flasks were incubated at 23±2°C in B.O.D with three replications of each treatment. The culture was then filtered with Whatman filter paper No. 1 and the mycelium was dried at 60°C in oven for 48 hours before measuring the dry weight of mycelial mat, on an electronic balance. The dry weight of mycelial mat was recorded after 10 days of inoculation [12].

Comment [I4]: Delete @ in all article

Comment [I5]: Write all abbreviation in complete formula in all article

2.4. Effect of different temperature on mycelial growth and dry matter weight.

For studies of suitable temperature, the culture of *Cordyceps militaris* were incubated at seven different temperature viz. 16, 18, 20, 22, 24, 26 and 28°C in BOD. Petri plates containing 20 ml of sterilized PDA medium were inoculated at the centre with 9 mm diameter disc of 7 days old actively growing culture of *Cordyceps militaris* under aseptic conditions. Inoculated plates were incubated in different B.O.D and three replications for each treatment were maintained. Observations on radial growth were taken at each 48 hrs in *Cordyceps militaris* until the colony covered the first full plate.

The PD broth was poured into 150 ml conical flasks @ 50 ml per flask and sterilized in autoclave at 121°C (1.1 kg/cm² pressure) for 20 minutes and allowed to cool. The 9 mm diameter disc was cut by cork borer from the periphery of 7 days old culture of *Cordyceps militaris* and inoculated in the flask with the help of inoculation needle. The flasks were incubated at seven different temperature viz. 16, 18, 20, 22, 24, 26 and 28°C in different B.O.D with three replications of each treatment. The culture was then filtered with Whatman filter paper No. 1 and the mycelium was dried at 60°C in oven for 48 hours before measuring the dry weight of mycelium mat, on an electronic balance. The dry weight of mycelial mat was recorded after 10 days [1].

2.5. Statistical analysis

The suitable statistical design (CRD) was applied and the data thus obtained were analyzed statistically. Analysis of variance (ANOVA) technique and critical difference (CD) was calculated at five percent level of significance for comparison with other treatment [9,12].

3. RESULT AND DISCUSSION

3.1 Effect of different pH on mycelial growth and dry matter weight.

Experiment was conducted to find out the effect of range of pH (5.5-8.5) on the mycelial growth and dry matter weight of Caterpillar Fungus (*Cordyceps militaris*). Observations were recorded on 2nd, 4th and 6th day. The Results revealed that on 6th day, maximum mycelial growth (89.00 mm) with (14.83 mm/day) growth rate was found at pH 7.0 which was significantly superior than all other treatments while it was followed by pH 7.5 (84.00 mm) with (13.83 mm/day) growth rate. Minimum mycelial growth (62.33 mm) was recorded at pH 8.5 with (10.38 mm/day) growth rate which was significantly lower than all other treatments. It was followed by pH 5.5 (66.66 mm) with (11.11 mm/day) growth rate as shown in Table-1, Fig-1 and Plate-1. While maximum dry matter weight (3.55 mg/50ml) of Caterpillar Fungus was observed at pH 7.0 with (0.35 mg/day) dry matter growth rate which was significantly superior to all other treatments and followed by pH 7.5 (2.54 mg/50ml) dry matter weight with (0.25 mg/day) dry matter growth rate. Minimum dry matter weight (0.93 mg/50ml) was observed at pH 8.5 with (0.09 mg/day) dry matter growth rate which was statically lower than all other treatments. It was followed by pH 5.5 (1.50 mg/50ml) dry matter weight with (0.15mg/day) dry matter growth rate as shown in Table-1, Fig-1 and Plate-2.

The results were in accordance with the findings of Sehgal and Sagar [22] studied the effect of different pH on the growth of *Cordyceps militaris*, the best solid medium was adjusted at different pH levels, viz. 4.0 to 10.0 and observed maximum mycelial growth was observed at pH 7.5. Lee *et al.* [15] also studied the influence of pH on *Cordyceps cardinalis* and observed that mycelial growth was highest when the pH of liquid medium was 7.0 and lowest at 4.0. Pathania *et al.* [19] tested the effect of different pH on the growth of *Cordyceps militaris*, the best solid medium, was adjusted at different pH levels (4.0 to 10.0). The best liquid medium was also adjusted at different pH levels (4.0 - 8.5). The inoculated petri plates and flasks were incubated for 10 days. The best mycelial growth of *Cordyceps militaris* was observed at pH 7.5 in solid and liquid medium respectively.

Comment [16]: Write why this pH suitable for growth mushroom mention causes

Table-01. Effect of different pH on mycelial growth (mm) and dry matter weight (mg) of *Cordyceps militaris*.

pH	Radial Growth (mm)			6 th days Growth rate (mm/day)	Dry Matter weight (mg/50ml)	Dry Matter Growth rate (mg/day)
	2 nd day	4 th day	6 th day			
5.5	31.00	54.66	66.66	11.11	1.50	0.15
6.0	31.00	56.66	67.33	11.22	1.53	0.15
6.5	33.66	64.33	79.33	13.22	2.10	0.21
7.0	34.66	68.66	89.00	14.83	3.55	0.35
7.5	33.66	65.66	84.00	13.83	2.54	0.25
8.0	32.00	62.00	72.33	12.05	1.88	0.18
8.5	30.33	53.33	62.33	10.38	0.93	0.09
CD at 5 %	2.25	4.33	4.25	-	0.41	-
SE(m)	0.73	1.57	1.90	-	0.10	-

*Average of three replications




Figure: 01. Effect of different pH on mycelial growth (mm) and dry matter weight (mg) of *Cordyceps militaris*.



4

Plate: 01. Effect of different pH on mycelial growth (mm) of *Cordyceps militaris*.
1. pH 5.5, 2. pH 6.0, 3. pH 6.5, 4. pH 7.0, 5. pH 7.5, 6. pH 8.0, 7. pH 8.5 .



7

Plate: 02. Effect of different pH on dry matter weight (mg) of *Cordyceps militaris*.
1. pH 5.5, 2. pH 6.0, 3. pH 6.5, 4. pH 7.0, 5. pH 7.5, 6. pH 8.0, 7. pH 8.5 .

3.2. Effect of different temperature on mycelial growth and dry matter weight.

A range of temperature (16-28 °C) was used to know its effect on the mycelial growth and dry matter weight of Caterpillar Fungus (*Cordyceps militaris*). Observations of effect of different temperature were recorded on 2nd, 4th and 6th day and shown in Table-2, Fig-2 and Plate-3. The Results revealed that on 6th day, maximum mycelial growth (89.66 mm) with (14.94 mm/day) growth rate was found at temperature 22 °C which was statically higher than all other treatments while it was followed by temperature 24 °C (84.00 mm) with (14.00 mm/day) growth rate. Minimum mycelial growth (52.00mm) was recorded at temperature 28 °C with (8.67 mm/day) growth rate which was significantly lower than all other treatments. It was followed by temperature 16 °C (65.66 mm) with (10.94 mm/day) growth rate. While maximum dry matter weight (1.46 mg/50ml) of Caterpillar Fungus (*Cordyceps militaris*) was also observed at temperature 22 °C with (0.14 mg/day) dry matter growth rate which was statically at par with temperature 20 °C (1.40mg/50ml) with (0.14 mg/day) dry matter growth rate and followed by temperature 24 °C (1.38mg/50ml) with (0.13 mg/day) dry matter growth rate. Minimum dry matter weight (1.18 mg/50ml) was observed at temperature 28 °C with (0.11 mg/day) dry matter growth rate which was statically at par with temperature 16 °C (1.27mg/50ml) with (0.12 mg/day) dry matter growth rate and followed by temperature 18 °C (1.35mg/50ml) with (0.13 mg/day) dry matter growth rate. The data are presented in Table-2, Fig-2 and Plate-4.

The results were in accordance with the findings of Hung, *et al.*[10] reported that, the temperature from 15 °C to 20 °C was suitable for mycelium growth of *Cordyceps militaris*. Pathania *et al.*[19] tested a range of temperature viz. 5 to 40 °C for growth of the fungus *Cordyceps militaris* in solid and liquid medium. [They found best mycelial growth of fungus *Cordyceps militaris* was found at 25 °C] both in solid and liquid media. Adnan *et al.*[1] also evaluated the optimum temperature for cordycepin production, fermentation was carried out in 250 mL conical flasks containing 100 mL of basal medium with initial pH 5.5 at 5 °C intervals in the range of 10–40 °C for 20 days under static condition, and found that 25 °C was the suitable temperature for maximum production of cordycepin.

Comment [17]: Mention why this temperature best

Table-02. Effect of different temperature on mycelial growth (mm) of *Cordyceps militaris*.

Temperature (°C)	Radial Growth (mm)			6 th days Growth rate (mm/day)	Dry Matter weight (mg/50ml)	Dry Matter Growth Rate (mg/day)
	2 nd day	4 th day	6 th day			
16	25.33	41.00	65.66	10.94	1.27	0.12
18	26.33	56.33	73.00	12.16	1.35	0.13
20	30.00	60.66	80.00	13.33	1.40	0.14
22	33.33	64.00	89.66	14.94	1.46	0.14
24	30.66	62.33	84.00	14.00	1.38	0.13
26	9.00	58.33	76.66	12.77	1.36	0.13
28	21.00	39.00	52.00	8.67	1.18	0.11
CD at 5 %	2.53	3.80	3.80	-	0.16	-
SE(m)	0.82	1.24	1.24	-	0.03	-

*Average of three replications

Figure: 02. Effect of different temperature on mycelial growth (mm) and dry matter weight (mg) of *Cordyceps militaris*.





Plate: 03. Effect of different temperature on dry matter weight (mg) of *Cordyceps militaris*. 1. 16 °C, 2. 18 °C, 3. 20 °C, 4. 22 °C, 5. 24 °C, 6. 26 °C, 7. 28 °C.



4

7

Plate: 04. Effect of different temperature on mycelial growth (mm) of *Cordyceps militaris*. 1. 16 °C, 2. 18 °C, 3. 20 °C, 4. 22 °C, 5. 24 °C, 6. 26 °C, 7. 28 °C.

4. CONCLUSION

Thus it can be concluded that maximal mycelium growth and dry matter weight of *Cordyceps militaris* can be achieved by cultivating fungus at pH 7.0 and in case temperature maximal mycelium growth and dry matter weight obtained at of 22°C and thus it's recommended for *Cordyceps militaris* culture conditions to be use.

REFERENCES

1. Adnan M, Ashraf SA, Kha S, Alshammari E, Awadelkareem AM. Effect of pH, temperature and incubation time on cordycepin production from *Cordyceps militaris* using solid-state fermentation on various substrates. *Journal of food*. 2017;15(4): 617–621.
2. Cunningham KG, Hutchinson SA, Manson W, Spring FS. Cordycepin, a metabolic product from cultures of *Cordyceps militaris* (Linn.) Link. Part I. Isolation characterisation, *J. Chem. Soc.* 1951;2299–3200.
3. Chang ST. Global impact of edible and medicinal mushroom human welfare in the 21st century: non green revolution. *Inter. J. of Medi. Mush.* 1999;1:1–7.
4. Das SK, Masuda M, Sakura IA, Sakakibara M. Medicinal uses of the mushroom *Cordyceps militaris*: current state and prospects. *Fitoterapia*. 2010;81:961-968.
5. Dlamini BE, Earnshaw DM, Masarirambi MT. Growth and yield response of Oyster mushroom (*Pleurotus ostreatus*) grown on locally available substrates. *Curr. Res. J. Biol. Sci.* 2012; 4 (5): 623-629.
6. De Silva DD, Rapior S, Fons F, Bahkali A H, Hyde KD. Medicinal mushrooms in supportive cancer therapies: an approach to anti-cancer effects and putative mechanisms of action. *Fungal Diversity*. 2012;55:1–35.
7. Guo C, Zhu J, Zhang C, Zhang L. Determination of adenosine and 3'-deoxyadenosine in *Cordyceps militaris* (L.) link by HPLC. *Zhongguo Zhong Yao Za Zhi*. 1998;23:236- 237.

Comment [18]: Old references replace

Comment [19]: Old references replace

8. Guo P, Kai Q, Gao J, Lian Z, Wu C, Wu, C, Zhu H. Cordycepin prevents hyperlipidemia in hamsters fed a high-fat diet via activation of AMP-activated protein kinase. *J. of Pharmacological Sci.*2010;13:395–403.
9. Gomez KA, Gomez AA. Statistical procedure for Agricultural research. 2nd edition. *J. Wiley and Sons*, New York. 1984:680.
10. Hung LT, Keawsompong S, Hanh V T, Sivichai S, Hywel-Jone NL. Effect of temperature on cordycepin production in *Cordyceps militaris*. *Thai J. of Agri. Sci.*2009;42(4): 219-225.
11. Koh JH, Yu KW, Suh HJ, Choi YM, Ahn TS. Activation of macrophages and the intestinal immune system by an orally administered decoction from cultured mycelia of *Cordyceps sinensis*. *Biosci. Biotechnol Biochem.*2002;66: 407- 411.
12. Kumar V, Mishra SK, Kaur M. Effect of different media, temperature and pH on radial mycelial growth of *Lentinula edodes* strain Le-17-04. *Journal of Pharmacognosy and Phytochemistry.* 2019;8(1): 345-348.
13. Kim JR, Yeon SH, Kim HS, Ahn YJ. Larvicidal activity against *Plutella xylostella* of cordycepin from the fruiting body of *Cordyceps militaris*. *Pest Manage. Scie.* 2002;58:713–717.
14. Ling JY, Sun YJ, Zhang PL, Zhang CK. Measurement of cordycepin and adenosine in stroma of *Cordyceps* sp. by capillary zone electrophoresis (CZE). *J. Biosci Bioeng.*2002;94:371-374.
15. Lee JO, Shrestha B, Kim TW, Sung GH, Sung JM. Stable formation of fruiting body in *Cordyceps bassiana*. *Mycobiology.* 2007;35:230-234.
16. Masuda M, Urabe E, Honda H, Sakurai A, Sakakibara M. Enhanced production of cordycepin by surface culture using the medicinal mushroom *Cordyceps militaris*. *Enzyme Microb. Tech.* 2007;40:1199-1205.
17. Mains EB. North American entomogenous species of *Cordyceps*. *Mycologi.*,1958;50(2):169-222.
18. Nag TB, Wang HX. Pharmacological actions of *Cordyceps*, a prized folk medicine. *J. Pharm. Pharmacol.*2005;57:1509-1519.
19. Pathania P, Joshi M, Sagar A. Morphological, physiological and molecular studies on wild collected *Cordyceps militaris* from North West Himalayas, India. *European J. of Biotech. and Biosci.*2015;3 (1): 53-62.
20. Pao HY, Pan BS, Leu SF, Huang BM. Cordycepin stimulated steroidogenesis in MA-10 mouse Leydig tumor cells through the protein kinase C Pathway. *Journal of Agri. and Food Chem.*2012;60:4905–4913.
21. Ramesh T, Yoo SK, Kim, SW, Hwang SY, Sohn SH, Kim IW, Kim SK. Cordycepin (3'-deoxyadenosine) attenuates age-related oxidative stress and ameliorates antioxidant capacity in rats. *Experi. Gerontol. I* 2012;47:979–987.
22. Sehgal AK, Sagar A. *In vitro* isolation and influence of nutritional conditions on the mycelial growth of the entomopathogenic and medicinal fungus *Cordyceps militaris*. *Plant Path. J.*2006;5(3): 315-321.
23. Sung JM. Production of fruiting body using cultures of entomopathogenic fungal species. *Korean J. of Mycology.* 1996;27:15-19.
24. Thomadaki H, Tsiapalis CM, Scorilas A. The effect of the polyadenylation inhibitor cordycepin on human Molt-4 and Daudi leukaemia and lymphoma cell lines. *Cancer Chemotherapy and Pharmacology.*2008;61:703–711.
25. Winkler D. Yartsa gunbu (*Cordyceps sinensis*) and the fungal commodification of Tibet's rural economy. *Econ. Bot.*2008;62:291–305.
26. Ying J, Mao X, Ma Q, D Wen H. Icons of medicinal mushroom from China. Beijing: *Science Press Beijing (in Chinese)*.1987:151-155.

Comment [I10]: Update references most it old