

Spatial and temporal variation of fish assemblage associated with aquatic macrophytes in three small lagoons of the South-eastern, Côte d'Ivoire

Abstract

Aquatic plants play a crucial role in an aquatic ecosystem partly because these plant communities provide suitable habitats and food items to other aquatic organisms especially fish. The purpose of the present study was to examine spatial variation in fish assemblage structure associated with aquatic macrophytes in three small lagoons of Côte d'Ivoire. Fish were monthly sampled from September 2014 to August 2016 using traps, gill nets, harpoons and hawks. The abiotic parameters namely temperature, transparency, pH, total dissolved solids, conductivity, dissolved oxygen, ammonium-nitrogen, nitrate, nitrite and phosphate were recorded for the characterization of the water mass in the lagoons. Significant variation in water parameters was observed among the sampling lagoons, except for nitrite and ammonium-nitrogen. However, no significant difference was obtained between Kodjoboue and Hebe lagoons. A total of 42, 35 and 37 species were recorded respectively in Ono, Kodjoboue and Hebe lagoons, with Perciformes (22 species), Siluriformes (8 species) and Osteoglossiformes (7 species) being the most diverse Orders. Habitats with highly amounts of macrophyte cover provide the optimal environment for many fish and hence, increase the taxonomic richness, abundance, diversity and equitability. The multivariate analysis showed that temperature, dissolved oxygen, nitrate, phosphate, pH and ammonium-nitrogen and macrophyte were key environmental variables and played important roles in determining species composition of fish assemblages.

Keywords: abundance, aquatic plants, diversity indices, fish, lagoons, Côte d'Ivoire

Introduction

Aquatic macrophytes play an important role in structuring fish assemblages [6 ; 20] by providing higher carrying capacity for food resources due to the availability of substrates for prey and higher productivity. They also increase the spatial complexity of the ecosystems, favouring the presence of numerous fish species [15 ; 23] and distribution of fish populations

Comment [u1]: The theme and the objective point to a relationship with macrophytes, while the methods and results presented point to physical chemical parameters of water. The proposal must be revised and better clarified

Formatted: Indent: First line: 0"

Comment [u2]: Alphabetical order

Comment [u3]: The term macrophytes is too generic and the reader has no idea what the authors are talking about. In the introduction, the authors must bring a paragraph talking about the taxa involved / targets of the study / found in the study sites

[19]. Their community assemblages can act as good indicators of the prevailing hydrological regime and water quality in aquatic systems. However, high densities of aquatic plants can lead to physical and chemical restrictions, causing fish avoidance behaviour, especially hypoxia at night during the hot season [13]. It is expected that these positive and negative effects on fish can explain their horizontal distribution and diel movements in lentic habitats near stands of macrophytes. The degree of response to these conditions is variable depending on fish assemblage composition, dominant life strategies and ontogenetic stage.

The role of macrophytes in the ecology of neotropicalichthyofauna is still not understood. There are some studies evaluating the effect of macrophytes coverage on fish assemblage structure [14 ; 21; 27]. In Côte d'Ivoire, no study addressing this topic has been conducted in marginal lagoons invaded by macrophytes. The present study aims to determine spatial and diel influence of macrophytes stands on fish assemblage and structure in Ono, Kodjoboue and Hebe lagoons of the South-eastern, Côte d'Ivoire.

2. Materials and methods

2.1. Study area

Ono lagoon (5°22'22"N and 3°33'53"W), Kodjoboue lagoon (5°14'11 "N and 3°35'9" W) and Hebe lagoon (5°12'14" N and 3°33'15" W) are three small lagoons of the Southeast of Ivory Coast (Figure 1). Their surfaces are respectively 400 ha, 423 ha and 244 ha. Because Ono lagoon is invaded by several habitat types such as emerged plants, free-floating macrophytes, floating leaf plants, submerged plants and white habitats, the exploitable surface is 162 ha. In the other lagoons, only the banks are occupied by macrophytes, with a pronounced degree of invasion in Kodjoboue lagoon. The main macrophytes are *Echinochloa pyramidalis*, *Eichhornia crassipes*, *Salvinia molesta*, *Pistia stratiotes* and *Hydrilla verticillata*. Hebe lagoon contains several species of mangroves such as

Comment [u4]: Present the ethical aspects of the research, authorization for the study. Fish collection methods assume that they were not returned to the lake after capture. In this case, scientific research should focus on reducing environmental impacts, not contributing to increasing impacts.

Rhizophoracemosa (Rhizophoraceae), *Avicenniagerminans* (Avicenniaceae) and *Conocarpus erectus*. These lagoons are irrigated by one or two small rivers and are connected in the downstream to Comoé River. This region has an equatorial climate, including two rainy seasons (April-July and October-November) and two dry seasons (December-March and August-September). The permanent linkage with the Comoé river produces typical freshwater characteristics of these lagoons.

Comment [u5]: Every taxon must be accompanied by the author descriptor.

Furthermore, species information must be in the introduction.




Figure 1: Map showing the sampling areas.

2.2. Environmental parameters sampling and analysis methods

The abiotic parameters namely temperature, transparency, pH, total dissolved solids (TDS), conductivity and dissolved oxygen (DO) were recorded *in situ*. Water samples were taken stored in polyethylene bottles (500 mL) and kept at a temperature below 4°C for further determination of ammonium-nitrogen (NH_4^+ ; mg/L), nitrate (NO_3^- ; mg/L), nitrite (NO_2^- ;

Comment [u6]: The physicochemical parameters should constitute an additional objective of the study. To review

mg/L) and phosphate (PO_4^{3-} ; mg/L). The samples were filtered through Whatman GF/C fibreglass filters and concentrations were determined using a spectrophotometer Model HACH DR 6000.

2.3. Fish sampling

In order to completely understand the effect of macrophytes on fish assemblages and determine their distribution characteristics, three lagoons were chosen following the occupation degree of macrophytes (Ono lagoon = 70%, Kodjoboue lagoon = 20% and Hebe lagoon = 2%). Fish were monthly sampled from September 2014 to August 2016 using traps, gill nets, harpoons and hawks. Specimens were systematically identified using the identification key of [22] and [16]. A sample of unknown fish were fixed in 10% formalin, placed in plastic bags and later transferred in 70% alcohol for further identification by specialists at the Oceanographic Research Center of Abidjan. Standard length of individuals was taken to the nearest 1 mm and body weight to the nearest 0.01 g using a Sartorius A200 S-F1 electronic balance.

2.4. Data analysis

Fish composition, rarefied richness (S) and numerical abundance (N) were determined during the month in each lagoon. Species diversity was measured considering the Shannon-Wiener diversity index (H') and Evenness index (E). In order to determine the permanence of the species in the evaluated sectors, the ecological index of consistency was used and each species was classified as constant ($C > 50$), incidental ($25\% \leq C \leq 50\%$), or accidental ($C < 25\%$) [4].

2.5. Statistical Analysis

Comment [u7]: Use a rarefaction analysis to assess and compare diversity/richness

The Shapiro-Wilk normality test for homoscedasticity were applied to the data, to determine whether the assumptions of the parametric and nonparametric analyses for the environmental variables (temperature, pH, DO, TDS, transparency, NH_4^+ , NO_3^- and NO_2^-), abundance, species richness, diversity and evenness were satisfied. To compare the effects of macrophytes on fish assemblage characteristics within and between sampling sites when the data were normally distributed, an analysis of variance (ANOVA) was used.

3. Results

3.1. Environmental variables

The environmental variables of Ono, Kodjoboue and Hebe lagoons were presented in Table 1. Significant variation in water parameters was observed among the sampling lagoons (ANOVA test, $p < 0.05$), except for nitrite and ammonium-nitrogen. However, no significant different (ANOVA test, $p > 0.05$) was observed between Kodjoboue and Hebe lagoons for all parameters. For the parameters such as temperature, pH, DO, conductivity and TDS, the values were significantly higher in Hebe lagoon whereas the values were lower for the temperature, pH and DO in Ono lagoon and for conductivity and TDS in Kodjoboue lagoon. On the other hand, the values of transparency, nitrate and phosphate were lower in Hebe lagoon and higher in Ono lagoon.

Table 1: Average values (mean \pm SD) of the physical and chemical parameters in Ono, Kodjoboue and Hebe lagoons between September 2015 and August 2016.

Parameters	Ono lagoon	Kodjoboue lagoon	Hebe lagoon
Temperature ($^{\circ}\text{C}$)	27.17 ± 1.56^a	29.73 ± 1.57^b	29.93 ± 1.38^b
Dissolved oxygen (mg/L)	2.29 ± 0.84^a	5.81 ± 0.97^b	6.22 ± 0.53^b
pH	6.32 ± 0.49^a	6.29 ± 0.58^a	6.67 ± 0.45^b
Conductivity ($\mu\text{S}/\text{cm}$)	18.09 ± 5.92^a	13.97 ± 6.53^a	35.87 ± 22.45^b
TDS (mg/L)	9.06 ± 1.58^a	6.96 ± 3.20^a	17.87 ± 3.16^b
Transparency (m)	1.59 ± 0.29^b	1.21 ± 0.20^a	0.71 ± 0.30^a
Nitrate (mg/L)	3.09 ± 0.93^b	2.31 ± 0.67^a	1.77 ± 0.57^a

Comment [u8]: Describe how macrophytes were evaluated in the environment, how were they sampled? how were they identified? how many samples were taken? Make it clear which variables were measured.

Comment [u9]: The subject of study is the fish community and macrophytes. Therefore, results should start with these findings. The parameters do not even constitute a study objective, therefore, they should not be the first to be presented.

Nitrite (mg/L)	0.21 ± 0.40 ^a	0.24 ± 0.47 ^a	0.14 ± 0.33 ^a
Ammonium-nitrogen (mg/L)	0.08 ± 0.04 ^a	0.06 ± 0.04 ^a	0.07 ± 0.04 ^a
Phosphate (mg/L)	0.48 ± 0.24 ^b	0.27 ± 0.15 ^a	0.31 ± 0.11 ^a

3.2. Fish Assemblage Characteristics

The composition of fish population (species, genera, families and order) sampled during this study is shown in Table 2. Specific richness showed a spatial pattern where the greatest number of fish species were recorded in Ono lagoon (42 species) vs 35 species in Kodjobouelagoon and 37 species in Hebe lagoon. The comparative analysis showed that six orders (Perciformes, Siluriformes, Osteoglossiformes, Characiformes, Cypriniformes, Clupeiformes) were common to all lagoons. The most diverse Order was that of Perciformes in Ono lagoon (10 families and 20 species), Kodjobouelagoon (3 families and 11 species) and Hebe lagoon (3 families and 12 species). This was followed by Siluriformes and Osteoglossiformes with 4 and 3 families, respectively, accounting for 5 and 4 species in Ono lagoon and 7 and 6 species in the other lagoons. The remaining orders presented one species by taxa and accounted for very low proportions of catches. Specially, all lagoons were dominated by Cichlids (10 species, 23.81 % in Ono lagoon, 9 species, 25.71% in Kodjobouelagoon and 12 species, 27.78% in Hebe lagoon). This family was followed by Alestidae (4 species, 9.52%), Mormyridae and Cyprinidae (3 species each, 7.14%) in Ono lagoon and by Clariidae, Alestidae and Mormyridae (4 species each) in Kodjobouelagoon and Hebe lagoons.

The spatial consistency analysis indicated that 11 species (26.19%) and ~~7~~ seven species (16.67%) were considered constant and incidental respectively in Ono lagoon. The remaining 24 species (57.14%) on the list were considered accidental. In Kodjobouelagoon, 8 species (22.86%) were defined as constant, 10 species (28.57%) as incidental and a further 17 species (48.57%) as accidental (Table 2). In Hebe lagoon, 10 species (27.03%) were classified as constant, 8 species (21.62%) as incidental and 19 species (51.35%) as accidental.

Comment [u10]: First result

Comment [u11]: Numbers less than 10, describe in full. see all cases

Table 2: Taxonomic list of the fish fauna with percentage and occurrence class in the Ono, Kodjoboue and Hebe lagoons between September 2015 and August 2016. Oc. = occurrence, % Oc. = percentage of occurrence, Oc. class = occurrence class, + = sporadic, ++ = frequent, +++ = very frequent, - = absent, Inc. = incidental, Acc. = accidental, C = constant.

Orders / Families	Species	Ono lagoon			Kodjoboue lagoon			Hebe lagoon		
		Oc.	%Oc.	Oc. class	Oc.	%Oc.	Oc. class	Oc.	%Oc.	Oc. class
SILURIFORMES										
CLARIIDAE	<i>Clariasebriensis</i>	+	20.55	Acc	+	13.80	Acc	+	28.33	Inc
	<i>Heterobranchuslongifilis</i>	-	-	-	+	22.46	Acc	+	26.66	Inc
	<i>Heterobranchusgariepinus</i>	-	-	-	+	18.66	Acc	+	25.67	Inc
	<i>Heterobranchusisopterus</i>	-	-	-	+	12.50	Acc	+	25.87	Inc
CLAROTEIDAE	<i>Chrysichthysmaurus</i>	+	15.65	Acc	-	-	-	-	-	-
	<i>Chrysichthysnigrodigitatus</i>	+++	100	C	++	68.86	C	+	74.25	C
					+					
SCHILBEIDAE	<i>Schilbemandibularis</i>	+	24.33	Acc	+	13.33	Acc	+	18.66	Acc
MALAPTERURIDAE	<i>Malapteruruselectricus</i>	+	06.45	Acc	+	10.33	Acc	+	11.67	Acc
MUGILIFORMES										
MUGILIDAE	<i>Liza falcipinnis*</i>	+	16.66	Acc	-	-	-	-	-	-
CHARACIFORMES										
HEPSETIDAE	<i>Hepsetusodoe</i>	+	18.86	Acc	++	15.45	Acc	+	20.66	Acc
ALESTIDAE	<i>Brycinusimberi</i>	+	15.33	Acc	++	26.25	Inc	+	18.67	Acc
	<i>Brycinusmacrolepidotus</i>	+	14.66	Acc	++	28.45	Inc	+	23.33	Acc
	<i>Brycinus nurse</i>	+	14.85	Acc	+	16.66	Acc	+	20.54	Acc
	<i>Brycinuslongipinnis</i>	+	18.28	Acc	+	13.66	Acc	+	24.66	Acc
DISTICHODONTIDAE	<i>Distichodusrostratus</i>	+	10.33	Acc	-	-	-	-	-	-
CYPRINIFORMES										
CYPRINIDAE	<i>Enteromiusablables</i>	+	19.25	Acc	+	12.25	Acc	+	16.67	Acc
	<i>Enteromiustrispilos</i>	+	23.33	Acc	-	-	-	-	-	Acc
	<i>Enteromiusburneensis</i>	-	-	-	++	25.15	Inc	+	18.66	Acc
	<i>Labeocoubie</i>	++	26.77	Inc	+	15.66	Acc	+	19.66	Acc
CLUPEIFORMES										
CLUPEIDAE	<i>Pellonulaleonensis</i>	++	25.33	Acc	+	20.66	Acc	+	15.67	Acc
ELOPTIFORMES										
ELOPTIDAE	<i>Eloplacerta</i>	++	25.33	Acc						
PERCIFORMES										

Comment [u12]: All names (taxa) must be presented accompanied by the author and year of the taxon. See all cases in the table and throughout the text

CHANNIDAE	<i>Parachannaobscura</i>	+++	81.50	C	++	45.26	Inc	++	51.67	C
GOBIIDAE	<i>Sicyopteruslagocephalus</i>	+	16.66	Acc	-	-		-	-	
POLYNEMIDAE	<i>Galeoidesdecadactylus</i>	+	15.67	Acc	-	-		-	-	
ANABANTIDAE	<i>Ctenopomapetherici</i>	++	26.66	Inc	+	19.67	Acc	+	23.33	Acc
HAEMULIDAE	<i>Pomadasyjsubelini</i>	+	11.89	Acc	-	-		-	-	
SPHYRAENIDAE	<i>Sphyraenaafra*</i>	+	06.67	Acc	-	-		-	-	
POLIMIDAE	<i>Polydactylusquadrifilis</i>	+	09.33	Acc	-	-		-	-	
MONODACTYLIDAE	<i>Monodactylussebae</i>	+	07.66	Acc	-	-		-	-	
ELEOTRIDAE	<i>Eleotrissenegalensis</i>	++	41.67	Inc	-	-		-	-	
	<i>Eleotrisvittata</i>	++	25.67	Inc	-	-		-	-	
CICHLIDAE	<i>Hemichromisbimaculatus</i>	+++	68.50	C	++	41.25	Inc	++	45.66	C
	<i>Hemichromisfasciatus</i>	+++	74.56	C	++	48.89	Inc	++	55.3	C
	<i>Oreochromisniloticus**</i>	++	45.67	Inc	+	18.33	Acc	+	22.56	Acc
	<i>Sarotherodonmelanotheron*</i>	+++	100	C	+++	100	C	+	100	C
	<i>Tilapia busumana</i>	++	30.28	Inc	-	-		-	-	
	<i>Coptodonguineensis*</i>	+++	98.26	C	+++	100	C	+++	100	C
	<i>Coptodonzillii</i>	+	27.33	Acc	+	17.66	Acc	++	36.86	Inc
	<i>Tylochromisjentinki*</i>	+++	86.66	C	+++	61.67	C	++	57.66	C
	<i>Pelmatolapiamarie</i>	+	15.50	Acc	+	12.15	Acc	+	18.15	Acc
	<i>Chromidotilapiaguntheri</i>	-	-	-	+	19.20	Acc	+	19.87	Acc
	Hybrid <i>Coptodon</i>	+++	100	C	+++	100	C	+++	100	C
	<i>Thysochromisansorgii</i>	-	-	-	-	-	-	+	10.12	Acc
OSTEOGLOSSIFORMES										
NOTOPTERIDAE	<i>Papyrocranusufer</i>	+++	86.87	C	+++	64.66	C	+++	77.55	C
MORMYRIDAE	<i>Marcuseniusfurcidens</i>	+++	50.50	C	+++	54.66	C	++	51.21	C
	<i>Marcuseniussheri</i>	++	48.67	Inc	++	45.55	Inc	++	48.67	Inc
	<i>Mormyropsanguilloides</i>	++	58.33	C	++	51.66	C	++	68.33	C
	<i>Marcuseniussenegalensis</i>	-	-		++	41.33	Inc	++	47.50	Inc
	<i>Mormyrusrume</i>	-	-		++	38.66	Inc	++	41.67	Inc
OSTEOGLOSSIDAE	<i>Heterotisniloticus**</i>	+	20.33	Acc	++	36.67	Inc	+	23.33	Acc
LEPDOSIRENIFORMES										
PROTOPTERIDAE	<i>Protopterusannectens</i>	-	-		-	-		+	4.33	Acc
9 orders / 25 families	51 species	42			35				37	

3.3. Diversity indices

Comment [u13]: Add a richness estimator, eg. Chao

The Shannon Specific Diversity Index (H') and Equitability Index (E) showed a low spatial variability (ANOVA test, $p > 0.05$) (Table 3). However, the recorded values were slightly higher in the Ono lagoon (4.5 bits/individual and 0.84 respectively) than that of Kodjoboue lagoon (4.45 bits/individual and 0.87 respectively) and Hebe lagoon (4.34 bits / individuals and 0.83, respectively).

Table 3: Diversity indices of Shannon and Equitability calculated for fish population of Ono, Kodjoboue and Hebe lagoons from September 2015 to August 2016.

Indices	Ono lagoon	Kodjoboue lagoon	Hebe lagoon
Shannon-Wiener index (H')	4.50 ^a	4.45 ^a	4.34 ^a
Equitability (E)	0.84 ^a	0.87 ^a	0.83 ^a

3.4. Fish abundance

In this study, the abundance of fish populations varied significantly (ANOVA test, $p < 0.05$) according to lagoons from September 2015 to August 2016 (Table 4). A total of 4138, 2630 and 2084 specimens were sampled respectively in Ono, Kodjoboue and Hebe lagoons. In general, Perciformes (48.11%), Osteoglossiformes (23.54%), Siluriformes (10.49%), Cypriniformes (7.66%) and Characiformes (7.62%) constituted the main orders in Ono lagoon. In Kodjoboue and Hebe lagoons, Perciformes was also the dominant order, accounting for respectively 58.95% and 56.65% of fish abundance, followed by Siluriformes (19.98% and 16.53% respectively), Osteoglossiformes (9.32% and 16.07% respectively) and Characiformes (9.97% and 6.25% respectively). These orders represented of about 97.42%, 98.18% and 99.00% of fish collected respectively in Ono, Kodjoboue and Hebe lagoons.

Table 4: Abundances of the main fish Orders caught in Ono, Kodjoboue and Hebe lagoons between September 2015 and August 2016.

Orders	Ono lagoon	Kodjoboue lagoon	Hebe lagoon
Perciformes	48.11	58.95	56.65
Siluriformes	10.49	19.98	16.53
Osteoglossiformes	23.54	9.32	16.07
Characiformes	7.62	6.97	6.25
Total	97.42	98.18	99.00

3.4. Factors influencing the distribution of species

The results of the Canonical Correspondence Analysis (CCA) showed that the correlation between environmental variables and fish was mainly explained by the first two axes represented (90,98%) of total variance (Figure 2). According to forward selection procedure and Monte Carlo permutation tests from CCA, several variables (macrophyte cover, temperature, DO, nitrate, phosphate, pH and ammonium-nitrogen) were significant in explaining patterns of occurrence and abundance of fish in the lagoons. The macrophytes, nitrate, ammonium-nitrogen and phosphate were positively correlated with the first axis whereas temperature and DO were negatively correlated with this axis. Concerning the second axis, it was positively correlated with pH. Ono lagoon was characterized by the abundance of macrophytes and high values of nitrate ammonia and phosphate and low values of temperature and dissolved oxygen. This lagoon was associated with the presence of numerous species belonging to Perciformes. On the other hand, Kodjoboue lagoon was characterized by high values of pH while Hebe lagoon was associated with low values of this parameter.

Comment [u14]: Canonical correspondence analysis is not mentioned in the method, nor is the software used. It also does not appear as an objective of the study. Authors should explain the use of this analysis, its purpose




Figure 2: Graphical representation of Canonical redundancy analysis based on the abundance of 50 fish species, environmental variables (Temp = water temperature, DO = dissolved oxygen, pH, NO₃⁻ = nitrate, PO₄³⁻ = phosphate and NH₄⁺ = ammonium-nitrogen), and lagoons (geometric figures) during the period-September 2015 to August 2016.

4. Discussion

Analysis of the physical and chemical parameters reveals that the parameters show significant variation between lagoon sites, except for nitrite and ammonium-nitrogen. Globally, the values of these parameters were higher in all lagoons due to aquatic macrophytes and numerous human activities, namely the intensification of poultry and pig

farm and the organic fertilizer from industrial plantations. However, the value of temperature, pH and dissolved oxygen were significantly lower in Ono lagoon than in the other lagoons. [9] reported that temperature plays an important role in the physical and chemical characteristics of lagoon environment, affecting the rate of CO₂ fixation by phytoplankton (primary productivity) and solubility of gases such as O₂, CO₂ and NH₄⁺ which on turn affect all aquatic organisms. The lowest values of oxygen levels may be due to the removal of free oxygen through respiration by, macrophytes, bacteria and animals as indicated by [25]. [3] noted that the dense mats of *Hydrillaverticillata* reduced water circulation and light penetration in water bodies and influenced dissolved oxygen concentrations. On the other hand, the value of nitrate, phosphate and ammonium-nitrogen were higher in Ono lagoon, indicating that this lagoon is most subject to anthropogenic pressures. In fact, its watershed is located in a zone which drain intensive agricultural activities using excessive application of chemical fertilizers and phytosanitary products. In addition, this lagoon is invaded by a multitude of macrophytes such as emergent plants, free-floating macrophytes, floating leafy plants, submerged plants and white habitats [5].

Comment [u15]: Add author

Fish assemblage structure seem to be influenced by both abiotic parameters and macrophytes cover. Fish assemblage responses were highlighted more in relation to the degree of macrophyte cover. The taxonomic richness, abundance, diversity and equitability decrease from areas most infested by macrophytes (Ono lagoon) to areas less infested (Hebe lagoon). Ono lagoon is heavily infested by several types of macrophytes, particularly *Eichhorniacrassipes*, *Hydrillavertillata* and *Echinochloapyramidalis* whereas in the other lagoons, only the banks are infested by *E. pyramidalis*, with a pronounced degree of invasion in Kodjoboue lagoon. Numerous studies showed that vegetated sites contain higher fish densities as compared to unvegetated areas [2; 12]. In fact, vegetation is used as breeding sites for some species of fish, nurseries and important refuges for their juvenile since they provide

Comment [u16]: Add author

minor shade, nesting and cover habitat for fishes [11]. Apparently, submerged vegetation (e.g. *H. vertillata*) is the key factor in the distribution and habitat use of adult fish. It is known that the sites of the plant communities in water column and their morphology attract and influence the production of epiphytic invertebrates which then serve as prey for a variety of fishes especially Cyprinidae, Percidae and Cyprinodontidae [18 ; 12]. Although aquatic plants are a vital contributor in maintaining stability within water bodies, excessive plant growth reduces growth and condition of fish due to reduced foraging efficiency [7; 12]. For [8], the limited role of physicochemistry influencing fish distribution may be a common phenomenon, especially when macrophytes do not develop excessive coverage. In addition, all the orders encountered present almost the same distribution pattern in the three lagoons. However, Perciformes, Elopiformes, Mugiliformes were frequently recorded in the Ono Lagoon having areas with high macrophyte complexity.

CCA results indicated that temperature, dissolved oxygen, nitrate, phosphate, pH and ammonium-nitrogen and macrophyte were key environmental variables and played important roles in determining species composition of fish assemblages. According to [10] and [1], oxygen concentration, predation and food availability are all influenced by macrophytes. Numerous authors predicted that higher structural complexity results in an increase in resource availability and variability, allowing the coexistence of a higher number of species in the same area [24 ; 26 ; 17].

Conclusion

6. References

1. Agostinho A.A., Thomaz S.M., Gomes L.C. & Baltar S., 2007. Influence of the macrophyte *Eichhornia azurea* on fish assemblage of the upper Parana River floodplain (Brazil). *Aquatic Ecology*, 41(4): 611-619.

Comment [u17]: What is the conclusion? A conclusion is missing. What do the results/findings indicate? for what? Conservation? conservation policies?

Comment [u18]: References must be updated. Only two of the last five years. Most must be recent, current.

2. Borawa J.C., Kerby J.H., Huish M.T. & Mullis A.W., 1979. Currituck Sound fish populations before and after infestation by Eurasian water-milfoil. *Proceedings of the Annual Conference South-eastern Association of Fish and Wildlife Agencies*, 32: 520-528.
3. Colon-Gaud J.C. 2003. Macroinvertebrate abundance and distribution of *Hydrilla* and *Ceratophyllum* habitats in the Atchafalaya River Basin, Louisiana. Masters Thesis, Louisiana State University, Baton Rouge, LA. 60p.
4. Dajoz R., 2002. Tratado de ecología. 2nd ed. Madrid: MundiPrensa.
5. Diarra B., Konan K.J., Yapo L.M. & Kouassi K.P., 2018. Aquatic macroinvertebrates associated with free-floating macrophytes in a marginal lentic ecosystem (Ono Lagoon, Côte d' Ivoire). *Journal of Entomology and Zoology Studies*, 6 (3): 1432-1441.
6. Dibble E.D., Killgore K.J. & Harrel S.L., 1996. Assessment of Fish-Plant Interactions. In: Miranda L. E. & DeVries D. R. (eds.). *Multidimensional Approaches to Reservoir Fisheries Management*, Am. Fish. Soc. Symposium 16, Bethesda, Maryland.
7. Dibble E.D., Killgore K.J. & Harrel S.L., 1997. Assessment of Fish-Plant Interactions. Aquatic Plant Control Miscellaneous Paper A-97-6. *American Fisheries Society Symposium*, 16: 357-372.
8. Dibble E.D. & Pelicice F.M., 2010. Influence of aquatic plant-specific habitat on an assemblage of small neotropical floodplain fishes. *Ecology of Freshwater Fish*, 19: 381-389.
9. Gawad S.S.A. & Abdel-Aal E.I., 2018. Impact of flood cycle on phytoplankton and the macro invertebrates associated with *Myriophyllum spicatum* in lake Nasser khors (Egypt). *Journal of Biological Sciences*, 18: 51-67.
10. Miranda L.E., Driscoll M.P. & Allen M.S., 2000. Transient physicochemical microhabitats facilitate fish survival in inhospitable aquatic plant stands. *Freshwater Biology*, 44 (4): 617-628.
11. Krischik V.A., Newman R. & Kyhl J., 1997. *Managing Aquatic Plants in Minnesota Lakes*. University of Minnesota. St Paul, MN, USA.
12. Meng L., Gray C., Taplin B. & Kupcha K., 2000. Using winter flounder growth rates to assess habitat quality in Rhode Island's coastal lagoons. *Marine Ecology Progress Series*, 201: 287-299.
13. Miranda L.E., Driscoll M.P. & Allen M.S., 2000. Transient physicochemical microhabitat facilitated fish survival in inhospitable aquatic plant stand. *Freshwater Biology*, 44: 617-628.
14. Meschiatti J., Arcifa M.S. & Fenerich-Verani N., 2000. Fish communities associated with macrophytes in Brazilian floodplain lakes. *Environmental Biology of Fishes*, 58: 133-143.
15. Neiff J.J., Neiff A.P. & Verón M.B.C., 2009. The role of vegetated areas on fish assemblage of the Paraná River floodplain: effects of different hydrological conditions. *Neotropical Ichthyology*, 7(1): 39-48.

16. Paugy D., Lévêque C. & Teugel G.G., 2003. Faune des poissons d'eaux douces et saumâtres de l'Afrique de l'ouest. Tome I & II. Edition IRD-MNHN-MRAC, Paris-Tervuren, 457 + 815 p.
17. Pierre J.I.S. & Kovalenko K.E., 2014. Effect of habitat complexity attributes on species richness. *Ecosphere*, 5(2):1-10.
18. Pípalová I., 2006. A Review of Grass Carp Use for Aquatic Weed Control and its Impact on Water Bodies. *Journal of Aquatic Plant Management*, 44: 1-12.
19. Sass G.G., Gille C.M., Hinke J.T. & Kitchell J.F., 2006. Whole-lake influences of littoral structural complexity and prey body morphology on fish predator-prey interactions. *Ecology of Freshwater Fish*, 15: 301-308.
20. Slade J.G., Dibble E.D. & Smiley P.C., 2005. Relationships between littoral zone macrophytes and the fish community in four urban Minnesota lakes. *Journal of Freshwater Ecology*, 20:635-640.
21. Suarez Y.R., Petrere Jr. M. & Catella A.C., 2001. Factors determining the structure of fish communities in Pantanal lagoons (MS, Brazil). *Fisheries Management and Ecology*, 8:173-186.
22. Teugels G.G., Lévêque C., Paugy D. & Traoré K., 1988. Etat des connaissances sur la faune ichthyologique des bassins côtiers de Côte d'Ivoire et de l'Ouest du Ghana. *Revue d'Hydrobiologie Tropicale*, 21(3): 221-237.
23. Thomaz S.M. & Cunha E.R., 2010. The role of macrophytes in habitat structuring in aquatic ecosystems: methods of measurement, causes and consequences on animal assemblages' composition and biodiversity. *Acta Limnologica Brasiliensia*, 22(2): 218-236.
24. Thomaz S.M., Dibble E.D., Evangelista L.R., Híguti J. & Bini L.M., 2008. Influence of aquatic macrophyte habitat complexity on invertebrate abundance and richness in tropical lagoons. *Freshwater Biology*, 53:358-367.
25. Tohouri P., Miessan Adja G.M., Soro G., Ake E.G., Konan I.N. & Biémi J., 2017. Physicochemical quality in rainy season water surface area Bonoua (Southeast of Ivory Coast). *International Journal of Innovation and Applied Studies*, 20(1):28-41.
26. Tokeshi M. & Arakaki S., 2012. Habitat complexity in aquatic systems: fractals and beyond. *Hydrobiologia*, 685:27-47.
27. Vono V. & Barbosa F.A.R., 2001. Habitats and littoral zone fish community structure of two natural lakes in southeast Brazil. *Environmental Biology of Fishes*, 61:371-379.