

## **Original Research Article**

### **Genetic variances, heritability and traits association of early maturing maize hybrids under induced drought at seedling and flowering stages**

#### ABSTRACT

**Aims:** Information on traits association and inheritance are crucial to designing appropriate breeding strategies for improving maize production and productivity in drought-prone ecologies. The objectives of this study were to investigate inter-trait relationships among maize hybrids and estimate genetic variances and heritability of drought tolerance parameters under seedling and flowering drought conditions.

**Methodology:** Sixty-six single cross hybrids generated using diallel mating design plus nine hybrid checks were evaluated using a 5 x 15 alpha lattice and randomized complete block designs in three replicates on the field and in the screenhouse respectively, during 2015 cropping season. Data were collected on grain yield, ears per plant, anthesis-silking interval, seedling aspect, chlorophyll content and leaf area and these data were subjected to analysis of variance as well as correlation and path coefficient analyses.

**Results:** Significant mean squares ( $P = .05$ ) were observed for all measured traits except leaf area and shoot fresh weight. Narrow-sense heritability estimate for grain yield was moderate (33.4%) on the field and low (0 – 25%) for all the seedling traits. The low narrow-sense heritability estimates observed for most seedling traits implied that the scope for improvement of these traits in the genotypes is limited. Seedling traits under drought stress were not directly correlated with grain yield on the field except number of dead leaves relative to the total number of leaves. Results of the path analysis revealed that number of leaves, number of dead leaves and chlorophyll content under the screenhouse conditions had significant direct effects on grain yield on the field.

**Conclusion:** Number of leaves, and chlorophyll content under drought at seedling stage could therefore be used as indicator traits for grain yield improvement in maize exposed to drought stress at flowering stage.

**Keywords:** Drought, heritability, genetic variance, path analysis, maize hybrids, screenhouse conditions.

36 **Introduction**

37 Maize is an important staple food, animal feed and industrial crop in sub-Saharan Africa  
38 (SSA). The savannas of the sub region offer ideal environments for maize production because  
39 they are characterized by high solar radiation, low night temperatures, low incidence of pests,  
40 and diseases. Over 85% of the population of people in the rural areas in Africa grows maize  
41 due to its suitability in diverse farming systems and its ability for increased yield under  
42 improved management practices as compared to other cereal crops [1]. Auta et al. [2]  
43 recommended that the development and accelerated deployment of maize hybrids would  
44 increase maize yields in the major maize producing countries of West and Central Africa  
45 (WCA). Studies have demonstrated that hybrids can increase farmers' maize yields by more  
46 than 40% in favourable growing environments and by more than 30% even under stressful  
47 conditions [3].

48 Drought is a major constraint in boosting maize production in Africa especially in the SSA,  
49 coupled with other abiotic stresses it has resulted in maize displacement by high value crops  
50 to marginal areas [4]. Drought exerts its effects on yield through its effects on physiological  
51 processes, and through losses in plant stand when such stress occurs during emergence and at  
52 the seedling stage. The extent of grain yield reduction in maize due to drought stress depends  
53 on the stage of crop development at the time of the stress and its severity. However, the  
54 flowering period in maize is the most sensitive to drought. The most economically feasible  
55 and sustainable way to boost maize production and productivity in SSA is to develop drought  
56 tolerant varieties for the farmers. Earlier studies conducted in the rainforest agro-climatic  
57 zone showed that maize planted early significantly out-yielded those planted later in the  
58 season primarily because grain filling coincided with the period of relatively high incident  
59 solar radiation [5]. Maize plants with excellent adaptive response to drought at seedling stage  
60 will help in combating the effect of early season drought in the rainforest agro-ecology of  
61 southwestern Nigeria. Therefore, the evaluation of maize crop at the seedling stage is an  
62 important aspect of crop breeding program with the objective to evolve drought tolerant  
63 varieties. Developing specific maize genotypes which can tolerate drought at seedling stage,  
64 especially in southwestern Nigeria where the rainfall pattern is bimodal, will offer the farmers  
65 in this region the opportunity of planting maize earlier in the year (that is, late February to  
66 early April), immediately after the first few rains and by extension West Africa. Therefore,  
67 development of high yielding cultivars that combine tolerance to drought at both seedling and  
68 flowering stages could be a coping strategy to combat detrimental effects of climate change  
69 on maize production and productivity in the sub-region.

**Comment [O1]:** WCA is an unnecessary abbreviation because used only once in the text

70 Grain yield is a complex trait and it is collectively influenced by various component traits,  
71 besides being polygenically inherited and highly influenced by environmental variation. The  
72 appropriate knowledge of interrelationships between grain yield and its contributing  
73 components can significantly improve the efficiency of breeding programs using appropriate  
74 selection indices [6]. In making selection for improved genotypes in maize, yield is the  
75 primary trait but selection under drought based on grain yield alone is inefficient due to low  
76 heritability of grain yield and the complexity of genotype-environment interactions [7].  
77 Genetic variance and heritability of maize grain yield are reduced under drought whereas  
78 secondary traits have relatively high genetic variance and heritability [8,9]. However, the  
79 relative usefulness of secondary traits as indirect selection criteria for grain yield is often  
80 inconclusive for all experiments because of the nature of the genetic materials and different  
81 conditions of experiments [10]. In the past years, the use of secondary traits with grain yield  
82 in making selection has increased selection breeding efficiency in maize grown under stress  
83 conditions by 20 to 50% [11, 12]. Badu-Apraku et al. [13] reported that the most reliable  
84 traits for selection for improved grain yield under drought stress in the early maturing  
85 germplasm were ear aspect, ears per plant, anthesis-silking interval, and plant aspect. There is  
86 dearth of information on reliable seedling drought traits for predicting improved grain yield  
87 under flowering drought conditions.

88 Correlation analysis helps to measure the level of relationships among traits and to establish  
89 the level at which these traits are mutually different [14]. Path coefficient analysis helps to  
90 know the nature, extent and direction of selection; it is the most valuable tool commonly used  
91 to establish the exact relationships in terms of cause and effect, identifying the direct, indirect  
92 and total (direct plus indirect) causal effects, as well as to remove any spurious effect that  
93 may be present [15]. Studies had reported that tolerance to drought stress at seedling stage  
94 and at flowering stage has no relationship, using correlation analysis [16]. However,  
95 correlation analysis, being a bivariate analysis only detect linear relationships between two  
96 variables while it assumes no significant influence of interrelationship of other variables.  
97 Therefore, the use of multivariate techniques such as path coefficient analysis, which  
98 considers the effect of several independent traits on a target (dependent) trait by partitioning  
99 the total correlation into direct and indirect effects (effects exerted through other independent  
100 variables) is necessary. In such analysis, traits with significant direct effect on a target trait  
101 are identified while those with indirect effect through other traits are also detected.  
102 Information obtained from such analysis is more important than mere correlation and would

103 not only help in identifying important secondary traits that could be used as indirect selection  
104 criteria for improving maize genotypes for a desired trait, but will provide adequate  
105 information useful to formulate base index which will in turn improve the efficiency of  
106 selection for the target trait.

107 The primary objectives of this present study were to (i) estimate genetic variance and  
108 heritability of traits for drought tolerance at seedling and flowering stages of maize hybrids;  
109 and (ii) investigate the inter-trait relationships under seedling and flowering drought stress  
110 conditions, using simple correlations and path co-efficient analyses.

111

## 112 **Materials and Methods**

### 113 **Generation of crosses**

114 Twelve inbred lines were planted in breeding nursery in a single-row plot, 5 m long at a  
115 spacing of 0.75 m x 0.25 m. Standard agronomic practices were employed to ensure good  
116 crop stand. At flowering, the twelve inbred parents were crossed in a diallel fashion to  
117 generate 66 single-cross hybrids. The different crosses were harvested separately, processed  
118 and packaged into trials for evaluation.

119

### 120 **Field performance evaluation**

121 Drought stress at flowering stage was achieved in the field. The experiment was conducted at  
122 the Crop Science Unit of the Teaching and Research Farm, Obafemi Awolowo University, Ile  
123 Ife (7°28'N 4°33'E, 244 m, 1200 mm rainfall) during the 2015 cropping season. The planting  
124 was done on the 21<sup>st</sup> September, such that flowering could coincide with a period of no  
125 rainfall (drought). A trial composed of the sixty-six (66) single cross hybrids plus nine checks  
126 was laid out using a 5 x 15 alpha lattice design with three replications. Each entry was  
127 planted into a single-row plot, 5 m long with 0.75 m spacing between rows and 0.50 m  
128 spacing between plants within a row. Three seeds were sown per hole and later thinned to  
129 two plants per hill to attain a population density of 53,333 plants ha<sup>-1</sup>. A compound fertilizer  
130 of NPK 15:15:15 was applied at the rates of 60 kg\_N, 60 kg\_P and 60 kg K<sub>2</sub>O ha<sup>-1</sup> at two  
131 weeks after planting. An additional 60 kg\_N ha<sup>-1</sup> was also applied as side dressing at four  
132 weeks after planting (WAP) using urea (46% N). On each plot on the field, data were  
133 collected on days to anthesis, days to silking, number of ears per plant (EPP), plant and ear  
134 heights. Plant aspect was rated on a scale of 1 to 9, where 1 = excellent overall phenotypic

**Comment [O2]:** WAP is an unnecessary abbreviation because used only once in the text

135 appeal and 9 = poor overall phenotypic appeal. Ear aspect was scored on a scale of 1 to 9,  
136 where 1 = clean, uniform and large ears and 9 = rotten, variable and small ears. Stay-green  
137 characteristic (leaf death score) was scored per plot at 70 days after planting, on a scale of 1  
138 to 9, where 1 = almost all leaves green and 9 = virtually all leaves dead. Harvested ears from  
139 each plot were shelled to determine the percentage grain moisture. Grain yield in kg ha<sup>-1</sup> was  
140 adjusted to 15% moisture content and computed from the shelled grain weight.

141

#### 142 **Screenhouse evaluation**

143 Drought stress at seedling stage was imposed in the screenhouse. The sixty-six single-cross  
144 hybrids plus nine hybrid checks were planted under drought stress imposed at the seedling  
145 stage at the screen house, Department of Crop Production and Protection, Obafemi Awolowo  
146 University, Ile Ife in 2015. The trial was laid out in a randomized incomplete block design  
147 with three replications. Six seeds of each inbred were sown per pot. The methodology  
148 proposed by Akinwale [et al.](#) [17] for screening maize genotypes for tolerance to drought at  
149 seedling stage was adopted, water was applied to each pot at the rate of 0.6 litres daily for 7  
150 days, after which watering stopped. Data were collected at two-day intervals from 3 days  
151 after watering had stopped (DAWS) till 9 DAWS on the following: plant height, number of  
152 leaves, number of dead leaves, leaf length and breadth, leaf area, and number of dead leaves  
153 relative to the total number of leaves (RDL) in percentage. In addition, seedling aspect was  
154 scored on a scale of 1 to 9, where 1 = absence of visible symptoms of stress: vigorous plants,  
155 no wilting, no dead leaves, no chlorosis, no height reduction and unrolled turgid leaves and 9  
156 = total collapse or 100% death of seedlings, dried leaves and stem, as described by Akinwale  
157 [et al.](#) [17].

158

#### 159 **Data analysis**

160 Data collected were subjected to analysis of variance using PROC general linear model  
161 (GLM) procedure of Statistical Analysis Software (SAS), version 9.2 [18]. In the analysis,  
162 genotype was considered as a random factor and to achieve that, random statement option in  
163 GLM was used. Genetic variances were estimated using Proc Varcomp procedure of SAS.  
164 Estimate of narrow-sense heritability was performed as the proportion of additive variance  
165 over phenotypic variance expressed in percentage and this was done on plot mean basis for  
166 all traits under the two study conditions. In addition, path coefficient analysis was carried out  
167 to partition total correlation among traits into direct and indirect effects considering grain

168 yield as the dependent variable. Pearson correlation (phenotypic correlation) was first  
169 calculated and then the correlation coefficients were partitioned into direct and indirect  
170 effects through path coefficient analysis using PATHSAS program developed by Cramer et  
171 al. [19].

172

## 173 **Results and Discussion**

174 Results of analysis of variance (ANOVA) showed significant ( $P = .05$ ) differences among the  
175 genotypes for all seedling traits evaluated in the screenhouse except leaf area and fresh shoot  
176 weight (Table 1). Similarly, significant ( $P = .05$ ) effects were observed for genotype for all  
177 measured traits under stress conditions on the field (Table 2). The significant mean squares  
178 observed among the genotypes under the different stress conditions indicated presence of  
179 wide genetic variability within the genetic materials, which implies that genetic progress  
180 could be achieved from selection for improvement for drought tolerance at both growth  
181 stages. The coefficients of variation (CVs) associated with the stressed environments are  
182 usually higher and the coefficients of determination ( $R^2$ ) lower than those associated with the  
183 non-stressed environments. Thus, CVs of less than 30% was obtained only for seedling  
184 aspect score, chlorophyll content and plant height under seedling drought conditions and for  
185 all traits measured under flowering drought conditions except ear aspect. In addition, the  
186 proportion of variation explained by the ANOVA model ( $R^2$ ) is greater than 50% for  
187 chlorophyll content, dry shoot weight and number of dead leaves and greater than 60% for all  
188 the flowering drought traits, indicating the reliability of the model.

189 The additive variance measures the variation due to the average effects of alleles (additive  
190 effects) and the variation in the effects that are transmitted from one generation to another  
191 while the dominance variance is the variance due to interaction of average effects of alleles  
192 (dominance effects) [20]. The dominance variance is a function of allele frequencies and the  
193 level of dominance. In our present study, the additive variance was greater than non-additive  
194 (dominance) variance for all traits except number of ears per plant under the flowering  
195 drought conditions (Table 3). In contrast, the dominance variance was greater than the  
196 additive variance for all traits under the seedling drought conditions, which implied that  
197 tolerance to drought involves different genes or combination of genes depending on the stage  
198 at which the drought set in in the crop cycle. Estimates of additive genetic variance were  
199 significantly different from zero for all traits under field drought conditions and for all traits  
200 under the screenhouse conditions except seedling aspect, dry shoot weight and fresh shoot

Formatted: Not Strikethrough

201 weight. Dominance variances were significantly different from zero for all measured traits  
202 except seedling fresh shoot weight. Due to negative estimates, additive variance of dry shoot  
203 weight and dominance variance of fresh shoot weight were equated to zero. Thus, their  
204 dominance to additive genetic variance ratio could not be estimated. The dominance to  
205 additive genetic variance ratios were greater than 0.5 for four out of the eight traits under  
206 drought at the flowering stage and for seven out of the nine traits under seedling drought  
207 conditions.

208

UNDER PEER REVIEW

209 Table 1: Mean squares of seedling aspects and other seedling traits of 66 hybrids plus 9 checks evaluated under induced drought at seedling stage at the  
 210 screen house, 2015

SV	Df	Seedling aspect score	Chlorophyll content ( $\mu\text{mol L}^{-1}$ )	Plant height (cm)	Leaf area, ( $\text{cm}^2$ )	Number of dead leaves	Number of leaves	RDL	Fresh shoot weight (g)	Dry shoot weight (g)
Rep	2	12.20**	25.39	102.09**	90.15	5.32	39.66	854.30	9.07**	0.16*
Entry	74	1.83*	55.62**	29.04*	94.20	10.00**	55.56**	717.58**	0.26	0.05**
Error	148	1.27	16.39	19.34	79.73	3.17	35.91	416.36	0.26	0.03
R <sup>2</sup>		0.46	0.63	0.45	0.37	0.62	0.44	0.47	0.49	0.53
CV		20.65	11.95	18.61	40.84	37.11	30.95	41.90	50.95	41.68

\* significant at 0.05 probability level, \*\* significant at 0.01 probability level

RDL = number of dead leaves relative to the total number of leaves.

213 Table 2: Mean squares for grain yield and other agronomic traits of 66 hybrids plus 9 hybrid checks evaluated under drought at flowering stage, at the  
 214 Teaching and Research Farm Obafemi Awolowo University, Ile Ife, 2015

Source of variation	Df	Grain yield	Ears per plant	Plant aspect	Ear Aspect	Anthesis - silking interval	Plant height	Ear height	Stay green characteristics
Block (Replicate)	12	652613.3**	0.16**	6.91**	5.64**	9.55**	1223.52**	388.68**	3.99**
Replicate	2	977096.1**	0.47**	9.76**	21.63**	9.75**	2628.29**	882.79**	13.26**
Genotype	74	318140.3**	0.112**	2.24**	2.73**	4.69**	455.57**	162.38**	2.17**

Formatted Table

Formatted Table

Error	136	120633.0	0.06	0.77	1.59	1.65	124.48	50.26	0.93	215
R <sup>2</sup>		0.68	0.62	0.75	0.63	0.71	0.79	0.75	0.68	
CV		27.26	33.22	15.35	16.7	31.69	11.35	16.35	16.3	216


217 \* significant at 0.05 probability level, \*\* significant at 0.01 probability level  
218

219

220

221

UNDER PEER REVIEW



222 Table 3: Additive ( $\hat{\sigma}_A^2$ ) and non-additive ( $\hat{\sigma}_{NA}^2$ ) variances, narrow-sense heritability estimates ( $h^2$ ),  
 223 for grain yield and other agronomic traits of hybrids generated from diallel crosses evaluated under  
 224 terminal drought in the field and under imposed drought in the screen house

Traits	$\hat{\sigma}_A^2 \pm SE$	$\hat{\sigma}_{NA}^2 \pm SE$	$\hat{\sigma}_{NA}^2 / \hat{\sigma}_A^2$	$h^2$ (%)
Flowering drought environment				
Grain yield	40465.76±3512.730	36639.40±2026.930	0.91	33.4
Ears per plant	0.02±0.001	0.02±0.001	1.00	7.3
Plant aspect	0.35±0.030	0.22±0.020	0.63	35.7
Ear aspect	0.53±0.040	0.06±0.020	0.11	46.0
Anthesis-silking interval	0.90±0.070	0.37±0.030	0.41	45.0
Plant height	99.55±7.380	30.60±2.590	0.31	49.9
Ear height	17.70±1.590	13.58±1.010	0.77	31.1
Stay green characteristic	0.436±0.030	0.20±0.020	0.45	44.0
Seedling drought environment				
Seedling aspect	0.002±0.010	0.140±0.015	61.670	0.39
Chlorophyll content	3.154±0.390	15.270±0.360	4.840	17.0
Plant height	2.000±0.240	2.080±0.230	1.040	18.7
Leaf area	1.520±0.540	4.970±0.910	3.270	4.3
Number of dead leaves	0.412±0.070	2.080±0.060	5.060	11.6
Number of leaves	6.870±0.690	12.200±0.500	1.770	25.0
RDL	5.320±1.190	30.060±1.630	5.650	7.6
Fresh weight	0.001±0.001	0±0.003	0	0.38
Dry weight	0±0.0002	0.011±0.001	0	0.0

225  
 226  
 227  
 228  
 229  
 230  
 231  
 232  
 233  
 234

235

236 Narrow-sense heritability estimate for grain yield was moderate (33.4%). The values of  
237 narrow-sense heritability for the traits ranged from 7.3% for ears per plant to 50% for plant  
238 height under drought at flowering and 0.38% for fresh weight to 25% for number of leaves  
239 under seedling drought conditions. Heritability estimates provide information on how  
240 probable a trait could be transmitted from parents to their offspring [20]. Low narrow-sense  
241 heritability estimate observed for most measured traits, especially under drought at seedling  
242 stage is an indication that the scope for improvement of these traits in the parents is limited.  
243 However, it might be desirable to increase the number of replications and locations to  
244 increase the accuracy of estimated entry means, and thus the heritability.

245 Adequate knowledge on the inter-relationship among traits is important in designing effective  
246 selection programs for crop improvement. Results showed that seedling traits were not  
247 directly correlated with grain yield except number of dead leaves relative to the total number  
248 of leaves which displayed significant negative relationship (Table 4). Ears per plant showed  
249 significant negative relationship with number of dead leaves and number of dead leaves  
250 relative to the total number of leaves. This implied that a hybrid with high number of dead  
251 leaves under drought at seedling stage will possibly give lower number of ears harvested per  
252 plot under drought conditions at flowering. Plant aspect had significant negative relationship  
253 with leaf area, number of leaves, shoot fresh weight and shoot dry weight. This suggests that  
254 maize hybrids under drought stress at seedling stage with high number of leaves, leaf area,  
255 shoot fresh weight and shoot dry weight would give rise to plants with excellent physical  
256 appearance under the same condition at flowering stage. In addition, stay green characteristic  
257 showed significant positive relationship with number of dead leaves and number of dead  
258 leaves relative to the total number of leaves. Though some of the seedling drought traits were  
259 found to be significantly correlated with traits taken on the field under flowering drought  
260 conditions, the strength of the relationships was generally low. Therefore, the results of this  
261 study showed that seedling drought tolerance traits cannot be used as the main criteria for  
262 predicting grain yield in the hybrids under flowering drought conditions. This corroborates  
263 the findings from the study of Meeks et al. [16] who reported that the seedling drought  
264 conditions were independent of drought responses at flowering on the field, possibly due to  
265 the type of screening environments used and the high genetic diversity that segregated for  
266 traits conditioning drought tolerance. Even though low values of correlation coefficients ( $r$ )  
267 were obtained,  $r$  measures only linear relationship, which path analysis helps to decompose

268 among the studied characters, thereby enhancing better interpretation of relationships as well  
269 as pattern of the effects of one trait on the other.

270 An important objective of this study was to identify seedling drought traits with greatest  
271 influence on grain yield for use in future breeding programs. Thus, the correlation  
272 coefficients of the seedling drought traits with grain yield were further partitioned into direct  
273 and indirect effects through path analysis (Table 5). The low residual effect (0.25) indicates  
274 that the seedling traits altogether contributed substantially to grain yield. Among the seedling  
275 traits, path analysis identified number of leaves (0.294), and number of dead leaves (-0.242),  
276 chlorophyll content (0.052) as traits with significant direct contributions to grain yield,  
277 whereas number of leaves had the highest direct effect. This result was not surprising since  
278 leaf synthesizes the photosynthates, which are stored in plant tissues and culminate in yield  
279 for the crop. Number of leaves was related to the number of photosynthetic components such  
280 as chloroplasts and therefore an increase in the number of leaves improves photosynthetic  
281 capacity. Path coefficient analysis showed that number of seedling leaves was the most  
282 important trait for grain yield improvement through its direct and indirect effects on grain  
283 yield. Number of leaves, relative number of dead leaves to the total number of leaves are  
284 contributing indirectly to the number of dead leaves. For number of leaves, number of dead  
285 leaves recorded the highest indirect effect. Under seedling drought conditions, in our study,  
286 number of leaves, chlorophyll content and leaf area were identified as the most important  
287 traits contributing to the variation in grain yield, suggesting that they are reliable secondary  
288 traits under drought stress conditions. In addition, these traits can be used as criteria for  
289 selecting for drought-tolerant genotypes under seedling stress conditions in early maturing  
290 maize hybrids. Under the field conditions, the path analysis identified ear aspect (-0.48), ears  
291 per plant (0.331), ear height (0.283), plant height (-0.255) and plant aspect (-0.219) as traits  
292 with the highest direct effect on grain yield (Table not shown). These results are similar with  
293 those of Badu-Apraku et al. [9], who identified plant and ear aspects and plant and ear heights  
294 as the most reliable traits for the simultaneous selection in the extra-early inbreds for  
295 improved yield under low-N and drought stress environments.

296 Table 4: Pearson correlation of traits evaluated under drought induced at the seedling at the screenhouse, Faculty of Agriculture and under drought at  
 297 flowering stage at the Teaching and Research Farm, Obafemi Awolowo University, Ile Ife, 2015.

		Traits under drought at flowering stage							
		Anthesis- silking interval	Ears per plant	Plant height	Ear height,	Plant aspect	Stay green characteristics	Ear aspect	Grain yield,
Traits under drought at seedling	Number of dead leaves	0.15*	-0.15*	0.07	0.03	0.01	0.30**	0.17**	-0.07
	Leaf area	0.02	0.08	0.06	0.02	-0.14*	-0.08	-0.02	-0.02
	RDL	0.14*	-0.18**	-0.03	-0.06	0.11	0.28**	0.24**	-0.14*
	Number of leaves	0.06	-0.11	0.11	0.10	-0.18**	0.09	0.05	0.12
	Seedling aspect	0.02	-0.13	-0.03	-0.05	0.12	0.13	0.07	-0.02
	Chlorophyll content	0.06	0.07	-0.09	-0.03	0.08	0.05	-0.03	0.09
	Shoot fresh weight	-0.03	0.03	0.10	0.08	-0.20**	-0.12	-0.09	0.04
	Shoot dry weight	-0.09	0.09	0.17**	0.16**	-0.22**	-0.10	-0.11	0.06

\* significant at 0.05 probability level, \*\* significant at 0.01 probability level, \*\*\* Significant at 0.05, 0.01 probability levels, respectively

298  
 299  
 300  
 301  
 302  
 303

304 Table 5: Estimates of direct (diagonal values in bold) and indirect (off-diagonal values) effects based on path analysis of yield attributing characters on grain  
 305 yield, evaluated under drought induced at the seedling stage in the screenhouse, and under drought that coincided with flowering stage.

	Seedling height	Number of dead leaves	Leaf area	RDL	Number of leaves	Seedling aspect	Chlorophyll content	Shoot fresh weight	Shoot dry weight
Seedling height	<b>-0.051</b>	-0.055	0.003	0.001	0.108	-0.002	-0.005	-0.006	-0.018
Number of dead leaves	-0.011	<b>-0.242</b>	0.001	0.149	0.169	0.003	0.001	0.006	-0.002
Leaf area	-0.021	-0.023	<b>0.008</b>	-0.007	0.069	-0.001	-0.009	-0.005	-0.020
RDL	0.000	-0.183	-0.003	<b>0.196</b>	0.035	0.003	-0.001	0.005	0.005
Number of leaves	-0.019	-0.139	0.002	0.023	<b>0.294</b>	0.001	-0.001	-0.001	-0.013
Seedling aspect	0.012	-0.109	-0.001	0.086	0.055	<b>0.007</b>	0.006	0.013	0.019
Chlorophyll content	0.005	-0.003	-0.001	0.001	-0.007	0.001	<b>0.052</b>	0.002	0.008
Shoot fresh weight	-0.013	0.068	0.002	-0.047	0.008	-0.004	-0.004	<b>-0.022</b>	-0.034
Shoot dry weight	-0.013	-0.007	0.002	-0.014	0.052	-0.002	-0.006	-0.011	<b>0.073</b>

306 Residual effect = 0.2517

307 **Conclusion**

308 There was narrow genetic base for seedling drought tolerance traits in the early-maturing  
309 maize germplasm studied and this might limit the scope of improvement of the traits under  
310 drought stress. In addition, low additive variances and heritability estimates were obtained  
311 for most traits. Number of leaves and number of dead leaves under seedling drought stress  
312 had the highest direct effects on grain yield under field conditions, indicating that these  
313 seedling traits are reliable predictors of grain yield of early maize hybrids.

314

315 **References**

- 316 1. Badu-Apraku B, Fakorede MAB, Oyekunle M, Yallou GC, Obeng-Antwi K, Haruna A.  
317 Gains in grain yield of early maize cultivars developed during three breeding eras under  
318 multiple environments. *Crop Sci.* 2015;55:527–539. doi:10.2135/cropsci2013.11.0783
- 319 2. Auta SJ, Akpoko JG, Mamman H, Hassan HA. Production trends of small-scale maize  
320 farmers under the SASSKAWA Global 2000 Programme in Kaduna State of Nigeria,  
321 2001:412-418.
- 322 3. Byerlee D, Jewell D.. The technological foundation of the revolution, pp. 127-143;  
323 1997.
- 324 4. Banziger M, Cooper M.. Breeding for low-input conditions and consequences for  
325 participatory plant breeding - examples from tropical maize and wheat. *Euphytica*  
326 2001;122:503-509. <https://doi.org/10.1023/A:1017510928038>
- 327 5. Fakorede MAB, Opeke B. . Weather factors affecting the response of maize to planting  
328 dates in a tropical rainforest location. *Experimental Agriculture* 1985;21:31-40.  
329 <https://doi.org/10.1017/S0014479700012229>
- 330 6. Mohammadia SA, Prasanna BM, Singh NN. . Sequential path model for determining  
331 interrelationship among grain yield and related characters in maize. *Crop Sci.*  
332 2003;43:1690-1697. ISSN:1435-0653.
- 333 7. Araus JL, Serret MD, Edmeades GO.. Phenotyping maize for adaptation to drought.  
334 *Frontiers in Physiology* 2012;3:305. doi:10.3389/fphys.2012.00305.
- 335 8. Ashraf M. Inducing drought tolerance in plants: Recent advances. *Biotechnol. Adv.*  
336 2010;28:169-183. DOI: 10.1016/j.biotechadv.2009.11.005.
- 337 9. Badu-Apraku B. Akinwale RO, Franco J, Oyekunle M. Assessment of reliability of  
338 secondary traits in selecting for improved grain yield in drought and low-nitrogen  
339 environments. *Crop Sci.* 2012;52:2050-2062.

- 340 10. Dao A, Sanou J, Traore EVS, Gracen V, Danquah EY. Selection of drought tolerant  
341 maize hybrids using path coefficient analysis and selection index. *Pakistan Journal of*  
342 *Biological Sciences* 2017. DOI:10.3923/pjbs.2017.132.139
- 343 11. Edmeades GO, Bolanos J, Chapman SC. Value of secondary traits in selecting for  
344 drought tolerance in tropical maize. In: Edmeades G. O., Banziger M., Mickelson H. R.,  
345 and Pena-Valdivia C. B., editors, *Developing drought and low N tolerant maize.*  
346 *Proceedings of a Symposium, El Batan, Mexico. D.F. 25-29 Mar. 1996. CIMMYT, El*  
347 *Batan, Mexico, D.F. p. 222-234; 1997.*
- 348 12. Chapman SC, Edmeades GO. Selection improves drought tolerance in tropical maize  
349 populations: II. Direct and correlated responses among secondary traits. *Crop Sci.*  
350 1999;39:1315-1324. DOI: 10.2135/cropsci1999.3951315x
- 351 13. Badu-Apraku B. Akinwale RO, Ajala SO, Menkir A, Fakorede MAB, Oyekunle M.  
352 Relationships among traits of tropical early maize cultivars in contrasting environments.  
353 *Agron. J.* 2011;103(3):717-729.
- 354 14. Bocanski J, Sreckov Z, Nastasic A.. Genetic and phenotypic relationship between grain  
355 yield and components of grain yield of maize (*Zea mays* L.). *Genetika* 2009;41(2):145-  
356 154. DOI:10.2298/gensr0902145b
- 357 15. Hefny M. Genetic parameters and path analysis of yield and its components in corn  
358 inbred lines (*Zea mays* L.) at different sowing dates. *Asian J. Crop Sci.* 2011; 3(3):106-  
359 117. DOI: 10.3923/ajcs.2011.106.117
- 360 16. Meeks M, Murray SC, Steve H, Dirk H. Measuring maize seedling drought response in  
361 search of tolerant germplasm. *Agronomy* 2013;3:135-147. DOI:  
362 10.3390/agronomy3010135.
- 363 17. Akinwale RO, Awosanmi FE, Ogunniyi OO, Fadoju AO. Determinants of drought  
364 tolerance at seedling stage in early and extra-early maturing maize hybrids. *Maydica*  
365 2016;62(1):9.
- 366 18. SAS Institute. *SAS/STAT User's Guide version 9.1, (TS1M0)*, SAS Institute  
367 Incorporated, Cary, North Caroline, USA; 2002.
- 368 19. Cramer CS, Wehner TC, Donaghy SB. PATHSAS: A SAS computer program for path  
369 coefficient analysis of quantitative data. *Journal of heredity* 1999;90: 260-262.  
370 DOI:10.1093/jhered/90.1.260
- 371 20. Hallauer A, Carena M, Miranda F. *Quantitative genetics in Maize breeding.* Springer. p  
372 57-59; 2010.