

## Original Research Article

### Morphological screening and genetic variability analysis in rice (*Oryza sativa* L.) genotypes for salinity tolerance

#### Abstract:

This ~~experimente study~~ was conducted to estimate the genetic variability and correlation between yield and ~~other~~ yield related characters of 22 rice genotypes at Genetics and Plant Breeding Farm of Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, ~~material consisted of 22 rice genotypes~~. This experiment was ~~conducted-laid~~ in Randomized Block Design (RBD) during *Kharif* season ~~of 2022~~ under saline condition. ~~In analysis of variance mean sum of square due to treatments were highly significant and hence there was sufficient variability among rice genotypes~~ Analysis of variance shows significant differences among treatments which shows that there was significant variation among 22 rice genotypes under saline condition. GCV (Genotypic coefficient of variation) and PCV (Phenotypic coefficient of variation) estimates were higher for traits such as panicle bearing tillers/plant, harvest index and biological yield indicating selection of rice genotypes for yield and component traits under saline condition. High heritability coupled with higher genetic advance was observed for traits like grains/panicle, days to 50% flowering, spikelets number and plant height. Grain yield/plant was strongly related to harvest index, panicle bearing tiller /plant and biological yield per plant. Path coefficient analysis showed that harvest index followed by biological yield /plant and panicle bearing tillers/plant has positive direct effect on grain yield/plant.

**Keywords:** ~~Genotypic and phenotypic coefficient of variation~~ GCV, PCV, Correlation, Heritability, Genetic advance, Path analysis.

## INTRODUCTION

Rice (*Oryza sativa*) is ~~comes under from family~~ Poaceae ~~family~~ (Gramineae). The haploid chromosome number of rice is  $n=12$  ( $2n=24$ ). ~~Either the species is diploid or tetraploid. Both *O. sativa* L. and *O. glaberrima* L. are diploid.~~ It is *Kharif*, annual and tropical crop. Rice after wheat is 2<sup>nd</sup> most important crop covering almost 90% of area across Asia alone (Ghosh *et al.*, 2016). United States Department of Agriculture estimates world rice production 2022-23 of 503.27 million metric tonnes. India approximately had 46 million hectares of land under cultivation of rice. Total production of rice during 2022-23 in India ~~was~~ 104.99 million metric tonnes (Ministry of farmer and agriculture welfare).

In India, saline soil covers roughly 9.38 million ha, in which 5.5 million ha are coastal saline and 3.88 million ha are other saline soils. Saline soils ~~area~~ are ~~found from~~ Gujarat ~~in (~~ West) to Jammu & Kashmir (Ladakh region) ~~in the (~~North) and from the Andaman & Nicobar Islands ~~in the (~~East) to Kanyakumari (~~in the~~ South). Saline soil ~~poses causes~~ significant danger on the ability to raise food production to keep up with rising demand (Dagar, 2005). Rice is extremely susceptible to salinity, especially when it is ~~still at~~ seedling ~~stage~~. The main ~~factor~~reason in a variety of physiological impairments and plant-inhibiting process is a high concentration of harmful ions, especially  $\text{Na}^+$ , which is brought on by salinity. From a physiological perspective, it interferes with potassium absorption, which is crucial for maintaining membrane potential, enzyme activity and cell turgor (Huong *et al.*, 2020). ~~Correct the statement {Due to correlation, path coefficient analysis determines the direct and indirect effects used by various traits on grain yield. Through critical examination of the genetic variation influencing yield related quantitative parameters, crop productivity under salt stressed soil can be improved. The heritability is the percentage of the total that is passed down from parent to offspring. Genetic progress and heritability work together to predict genetic gain (Singh and Verma, 2018).}~~

## MATERIALS AND METHODS

### Plant materials and Experimental design

The present investigation ~~were carried out consisted of 22 rice genotypes. These were grown~~ during *Kharif* season ~~of 2022-22~~ in Randomized Block Design (RBD) at Genetics and Plant Breeding Farm of Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.) ~~during 2022-23-224229. The All 22~~ genotypes were sown in nursery

bed on 26 June 2022 and 35 days old [healthy](#) seedlings were transplanted on 30 July 2022 with a spacing of 20 ~~×~~ 15 cm with one seedling per hill.

### **Data collection:**

Observations on quantitative traits like days to 50% flowering, plant height(cm), panicle length(cm), panicle bearing tillers per plant, spikelets per panicle, grains per panicle, spikelet fertility(%), test weight(g), biological yield /plant(g), harvest index and grain yield per plant(g) were recorded on five randomly selected plants from each replication excluding border rows while days to 50% flowering and grain yield was calculated on plot basis. Agronomic practices were carried out as per recommendation to raise a good and healthy crop.

### **Statistical analysis:**

The analysis of variance [for the design of experiment](#) was carried out according to the procedure outlined by Panse and Sukhatme (1967), coefficient of variability was done as per Burton and de Vane (1953), correlation was calculated as per method suggested as per Searle (1961), heritability and genetic advance was calculated as per methods of Hanson *et al.*, (1956) and Johnson *et al.*, (1955) respectively and path analysis was computed according to Dewey and Lu (1959).

## **RESULT AND DISCUSSION**

For all 11 characters, an analysis of variance (ANOVA) was performed [using 22 rice genotypes](#) under saline condition. Analysis of variance shows significant differences among treatments which shows that there was significant variation among 22 rice genotypes as presented in table 1.

Estimates of grand mean, phenotypic coefficient of variability (PCV), genotypic coefficient of variability (GCV), heritability in broad sense ( $h^2b$ ) and genetic advance in percent of mean for 11 characters in 22 rice genotypes under saline condition are presented in table 2. It shows that in general phenotypic coefficient of variation (PCV) higher than genotypic coefficient of variation (GCV) this shows the effect of environment on the character.

High magnitude of PCV was observed for panicle bearing tillers/plant, harvest index and biological yield. This indicates that these characters can be manipulated for high yielding varieties through hybridization and selection in subsequent generations in rice improvement.



**Table 2: Estimates of grand mean, (PCV), (GCV), heritability in broad sense ( $h^2_b$ ) and genetic advance in percent of mean for 11 characters in 22 rice genotypes under saline condition**

Characters	Grand mean	Coefficient of variation		Heritability in broad sense (%)	Genetic advance in percent of mean
		PCV	GCV		
Days to 50% flowering	92.53	10.78	10.27	90.9	20.17
Plant height (cm)	100.04	11.52	10.49	82.9	19.66
Panicle bearing tillers/plant	9.71	15.57	11.75	56.9	18.26
Panicle length(cm)	21.54	10.9	9.49	75.7	17.01
Spikelets number	124.32	14.4	13.42	86.9	25.78
Grains / panicle	97.37	14.48	13.86	91.6	27.33
Spikelet fertility (%)	79.52	6.35	5.3	69.6	9.12
Test weight(g)	24.21	9.94	7.42	55.7	11.41
Biological Yield(g)	40.73	14.5	11.71	65.3	19.49
Harvest Index(g)	35.69	15.24	11.61	58	18.23
Grain yield(g)	14.33	11.92	10.15	72.5	17.80

**Table 3: Phenotypic correlation under saline condition**

Characters	Days to 50% flowering	Plant height (cm)	Panicle bearing tillers/plant	Panicle length	Spikelet No.	Grain per panicle	Spikelet Fertility %	Test weight (g)	Biological Yield/plant (g)	Harvest Index (%)	Grain yield / plant (g)
Days to 50% flowering	<b>1.000</b>	0.197	0.048	-0.179	0.119	0.148	-0.204	-0.068	0.069	-0.285*	-0.276*
Plant height (cm)		<b>1.000</b>	-0.101	0.273*	-0.040	-0.017	-0.224	-0.034	-0.058	0.017	-0.052
Panicle bearing tillers/plant			<b>1.000</b>	0.030	-0.143	0.206	0.175	0.048	0.287*	-0.018	0.369**
Panicle length				<b>1.000</b>	-0.176	-0.080	0.052	0.334**	0.236	0.011	0.274*
Spikelet No.					<b>1.000</b>	0.712**	-0.253*	-0.150	0.218	-0.011	0.261*
Grain per panicle						<b>1.000</b>	0.064	-0.224	0.222	0.022	0.293*
Spikelet Fertility %							<b>1.000</b>	0.184	-0.054	0.099	0.024
Test weight (g)								<b>1.000</b>	0.201	0.053	0.281*
Biological Yield/plant (g)									<b>1.000</b>	-0.655**	0.322**
Harvest Index (%)										<b>1.000</b>	0.489**

\*,\*\*Significant at 5% and 1% probability levels, respectively.

**Table 4: Direct and indirect effect for different characters on grain yield per plant under saline condition at phenotypic level in rice genotypes**

Characters	Days to 50% flowering	Plant height (cm)	Panicle bearing tillers/plant	Panicle length	Spikelet No.	Grain per panicle	Spikelet Fertility %	Test weight (g)	Biological Yield/plant (g)	Harvest Index (%)	Grain yield / plant (g)
Days to 50% flowering	<b>-0.027</b>	-0.001	0.006	-0.005	0.012	-0.007	0.008	-0.001	0.071	-0.332	-0.276*
Plant height (cm)	-0.005	<b>-0.006</b>	-0.013	0.007	-0.004	0.001	0.009	0	-0.059	0.02	-0.052
Panicle bearing tillers/plant	-0.001	0.001	<b>0.126</b>	0.001	-0.015	-0.01	-0.007	0	0.295	-0.021	0.369**
Panicle length	0.005	-0.002	0.004	<b>0.026</b>	-0.018	0.004	-0.002	0.003	0.242	0.013	0.274*
Spikelet No.	-0.003	0	-0.018	-0.005	<b>0.101</b>	-0.035	0.01	-0.001	0.224	-0.012	0.261*
Grain per panicle	-0.004	0	0.026	-0.002	0.072	<b>-0.049</b>	-0.003	-0.002	0.228	0.026	0.293*
Spikelet Fertility %	0.006	0.001	0.022	0.001	-0.026	-0.003	<b>-0.039</b>	0.001	-0.055	0.115	0.024
Test weight (g)	0.002	0	0.006	0.009	-0.015	0.011	-0.007	<b>0.008</b>	0.206	0.062	0.281*
Biological Yield/plant (g)	-0.002	0	0.036	0.006	0.022	-0.011	0.002	0.002	<b>1.028</b>	-0.762	0.322**
Harvest Index (%)	0.008	0	-0.002	0	-0.001	-0.001	-0.004	0	-0.673	<b>1.163</b>	0.489**

**R Square = 0.9742 Residual effect = 0.1606**

**Bold values show direct and normal values show indirect effect.**

## CONCLUSION:

The analysis of variance shows there is presence of sufficient variability among rice genotypes under saline condition. Values of phenotypic coefficient of variation were higher than genotypic coefficient of variation for all the eleven characters under saline condition. Higher magnitudes of phenotypic coefficient of variation were found for panicle bearing tillers/plant, harvest index, biological yield/plant, grain/panicle and spikelet number. High estimates of heritability were observed for grain per panicle, days to 50% flowering, spikelet number, plant height. Whereas medium estimates of heritability were observed for grain yield per plant, spikelet fertility, biological yield per plant, harvest index, panicle bearing tillers per plant and test weight. Under saline condition grains/panicle showed highest value of genetic advance and spikelet fertility showed lowest value of genetic advance. Grain yield showed highly significant and positive phenotypic correlation with harvest index, panicle bearing and biological yield/plant. Harvest index showed maximum positive direct effect on grain yield followed by biological yield/plant and panicle bearing tillers/plant.

## References:

~~Adhikari, B. N., Joshi, B. P., Shrestha, J., & Bhatta, N. R. (2018). Genetic variability, heritability, genetic advance and correlation among yield and yield components of rice (*Oryza sativa* L.). *J. of Agri. and Natural Res.*, **1(1)**:149-160. Not mentioned any where in main manuscript~~

Beena, R., Veena, V., and Jaslam, M. P. K. (2021) Germplasm innovation for high-temperature tolerance from traditional rice accessions of Kerala using genetic variability, genetic advance, path coefficient analysis and principal component analysis. *J. Crop Sci. Biotechnol.* **24**, 555–566.

~~Correct this Bhargava, K., Shivani, D., Phushpavalli, S.N.C.V.L., Sundaram, R.M., Beulah, P. and Senguttuvel, P.(2021). *Electr. J. of Plant Breed.* **12(2)**: 549-555.~~

Chamar, J. P., Joshi, R. P., Katkani, D., Sahu, D. and Patel, V. (2021) Genetic variability, heritability and correlation coefficient analysis in rice. *The PharmaInnov. J.* **10 (5)**:763-769.

Formatted: Font color: Text 1

- Dhurai, S. Y., Bhati, P. K. and Saroj, S. K. (2014) Studies on genetic variability for yield and quality characters in rice (*Oryza sativa* L.) under integrated fertilizer management. *The Bioscan* **9(2)**: 845-848.
- Kulsum, U., Sarker, U. and Rasul, M.G. (2022). Genetic variability, heritability and interrelationship in salt-tolerant lines of T. Aman rice. *Genetika*, **54(2)**:761-776.
- Kumar, G., Dwivedi, D.K., Maurya, H. and Yadav, V.K. (2018). Correlation and path coefficient analysis of thirteen rice genotypes for grain yield and other yield attributing traits. *J. Pharmacogn Phytochem* **7(3)**:3506-3510.
- Lingaiah, N.Satish , B., Venkanna, V., Rukmini, K. and Hari, Y.(2020). Genetic Variability and Correlation Studies in Yield Traits of Elite Rice (*Oryza sativa* L.) Genotypes. *Ind. J.Pure App. Biosci.* **8(6)**: 359-363.
- Mishra, V.K., Dwivedi, D.K., Sunil, K. and Verma, O.P. (2012). Screening for salinity tolerance at vegetative and reproductive stages in rice (*Oryza sativa* L.). *Environment and Ecology*, **30(2)**:355-360.
- Nihad, S.A.I., Manidas, A.C., Hasan, K., Hasan, M.A., Honey, O. and Latif, M.A.(2021) Genetic variability, heritability, genetic advance and phylogenetic relationship between rice tungro virus resistant and susceptible genotypes revealed by morphological traits and SSR markers. *Curr. Plant Biol.***25**: 2214-6628.
- Rahman, M. A., Haque, M., Sikdar, B., Islam, M. A., &Matin, M. N. (2014). Correlation Analysis of Flag Leaf with Yield in Several Rice Cultivars. *J. of Life and Earth Sci.* **8**:49–54.
- Singh, S.K., Singh, C.M. and Lal, G.M. (2011) Assessment of genetic variability for yield and its component characters in rice (*Oryza sativa* L.) *Res. in Plant Biol.***1(4)**:73-76.