

Original Research Article

Morphological screening and genetic variability analysis in rice (*Oryza sativa* L.) genotypes for salinity tolerance

Abstract:

The study was conducted to estimate the genetic variability and correlation between yield and other yield related characters of rice genotypes at Genetics and Plant Breeding Farm of Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya, material consisted of 22 rice genotypes. The experiment was conducted in Randomized Block Design (RBD) during *Kharif* season of 2022 under saline condition. In analysis of variance mean sum of square due to treatments were highly significant and hence there was sufficient variability among rice genotypes under saline condition. GCV and PCV estimates were higher for traits such as panicle bearing tillers/plant, harvest index and biological yield indicating selection of rice genotypes for yield and component traits under saline condition. High heritability coupled with higher genetic advance was observed for traits like grains/panicle, days to 50% flowering, spikelets number and plant height. Grain yield/plant was strongly related to harvest index, panicle bearing tiller /plant and biological yield per plant. Path coefficient analysis showed that harvest index followed by biological yield /plant and panicle bearing tillers/plant has positive direct effect on grain yield/plant.

Keywords: Genotypic and phenotypic coefficient of variation, Correlation, Heritability, Genetic advance, Path analysis.

INTRODUCTION

Rice (*Oryza sativa*) is from family Poaceae (Gramineae). The haploid chromosome number of rice is $n=12$ ($2n=24$). Either the species is diploid or tetraploid. Both *O. sativa* L. and *O. glaberrima* L. are diploid. It is *Kharif*, annual and tropical crop. Rice after wheat is 2nd most important crop covering almost 90% of area across Asia alone (Ghosh *et al.*, 2016). United

Comment [IA1]: Introduction should also contain a factual background, clearly defined problem, proposed solution, a brief literature survey and the scope and justification of the work done. Reference style for this journal is number in bracket []

Comment [IA2]: [1]

States Department of Agriculture estimates world rice production 2022-23 of 503.27 million metric tonnes. India approximately had 46 million hectares of land under cultivation of rice. Total production of rice during 2022-23 in India is 104.99 million metric tonnes (Ministry of farmer and agriculture welfare).

Comment [IA3]: [2], etc

In India, saline soil covers roughly 9.38 million ha, in which 5.5 million ha are coastal saline and 3.88 million ha are other saline soils. Saline soils are Gujarat in West to Jammu & Kashmir (Ladakh region) in the North and from the Andaman & Nicobar Islands in the East to Kanyakumari in the South. Saline soil poses significant danger on the ability to raise food production to keep up with rising demand (Dagar, 2005). Rice is extremely susceptible to salinity, especially when it is still a seedling. The main factor in a variety of physiological impairments and plant-inhibiting process is a high concentration of harmful ions, especially Na^+ , which is brought on by salinity. From a physiological perspective, it interferes with potassium absorption, which is crucial for maintaining membrane potential, enzyme activity and cell turgor (Huong *et al.*, 2020). Due to correlation, path coefficient analysis determines the direct and indirect effects used by various traits on grain yield. Through critical examination of the genetic variation influencing yield-related quantitative parameters, crop productivity under salt-stressed soil can be improved. The heritability is the percentage of the total that is passed down from parent to offspring. Genetic progress and heritability work together to predict genetic gain (Singh and Verma, 2018).

MATERIALS AND METHODS

Plant materials and Experimental design

Comment [IA4]: Growth medium and saline condition (sodium concentration) should also be described.

The present investigation consisted of 22 rice genotypes. These were grown during *Kharif* season of 2022-22 in Randomized Block Design (RBD) at Genetics and Plant Breeding Farm of Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.)-224229. The genotypes were sown in nursery bed on 26 June 2022 and 35 days old seedlings were transplanted on 30 July 2022 with a spacing of 20 X 15 cm with one seedling per hill.

Data collection:

Comment [IA5]: A method for estimating phenotypic and genotype variation should also be provided.

Observations on quantitative traits like days to 50% flowering, plant height(cm), panicle length(cm), panicle bearing tillers per plant, spikelets per panicle, grains per panicle, spikelet

fertility(%), test weight(g), biological yield /plant(g), harvest index and grain yield per plant(g) were recorded on five randomly selected plants from each replication excluding border rows while days to 50% flowering and grain yield was calculated on plot basis. Agronomic practices were carried out as per recommendation to raise a good and healthy crop.

Statistical analysis:

The analysis of variance for the design of experiment was carried out according to the procedure outlined by Panse and Sukhatme (1967), coefficient of variability was done as per Burton and de Vane (1953), correlation was calculated as per method suggested as per Searle (1961), heritability and genetic advance was calculated as per methods of Hanson *et al.*, (1956) and Johnson *et al.*, (1955) respectively and path analysis was computed according to Dewey and Lu (1959).

RESULT AND DISCUSSION

For all 11 characters, an analysis of variance (ANOVA) was performed using 22 rice genotypes under saline condition. Analysis of variance shows significant differences among treatments which shows that there was significant variation among 22 rice genotypes as presented in table 1.

Estimates of grand mean, phenotypic coefficient of variability(PCV), genotypic coefficient of variability (GCV), heritability in broad sense (h^2b) and genetic advance in percent of mean for 11 characters in 22 rice genotypes under saline condition are presented in table 2. It shows that in general phenotypic coefficient of variation (PCV) higher than genotypic coefficient of variation(GCV) this shows the effect of environment on the character.

High magnitude of PCV was observed for panicle bearing tillers/plant, harvest index and biological yield. This indicates that these characters can be manipulated for high yielding varieties through hybridization and selection in subsequent generations in rice improvement.

High estimates of heritability were observed for grains per panicle, days to 50% flowering, spikelet number and plant height (Table 2) under saline condition. Grains/panicle, spikelets number, days to 50% flowering, plant height, biological yield/plant, spikelet fertility % , harvest index, panicle length, test weight and grain yield/plant showed high values of genetic advance in percent of mean. Fiyazet *al.*, (2011) and Rahman *et al.*, (2013) showed similar result.

The simple linear correlation coefficient between all the pairs of studied traits are given in table 3. Grain yield showed highly significant and positive correlation with harvest index, panicle bearing tillers/plant and biological yield/plant and significant and negative correlation with days to 50% flowering. Under saline condition harvest index showed maximum positive direct effect followed by biological yield/plant and panicle bearing tillers/plant on grain yield/plant as presented in table 4. This suggests that these traits can be utilized for developing high yielding varieties. Mishra *et al.*, (2012) and Kulsum *et al.*, (2022) showed similar result.

Table 1: Analysis of variance for different traits

Comment [IA6]: P value for the significance differences should also be provided in the table.

Comment [IA7]: The variation for this study is 22 genotype

Characters	Sources of variation		
	Replication	Treatment	Error
d.f.	2	21	42
Days to 50% flowering	8.92	280.05**	9.08
Plant height (cm)	61.6	352.97**	22.77
Panicle bearing tillers/ plant	3.05	4.89**	0.99
Panicle length(cm)	1.61	13.87**	1.34
Number of spikelets	3.18	877.08**	41.88
Grains / panicle	42.16	563.16**	16.73
Spikelet fertility (%)	5.16	61.08**	7.75
Test weight	1.67	12.24**	2.56
Biological Yield	10.45	80.35**	12.1
Harvest Index	6.81	63.93**	12.41
Grain yield	2.39	7.15**	0.8

*, **Significant at 5% and 1% probability levels, respectively.

Table 2: Estimates of grand mean, (PCV), (GCV), heritability in broad sense (h^2_b) and genetic advance in percent of mean for 11 characters in 22 rice genotypes under saline condition

Characters	Grand mean	Coefficient of variation		Heritability in broad sense (%)	Genetic advance in percent of mean
		PCV	GCV		
Days to 50% flowering	92.53	10.78	10.27	90.9	20.17
Plant height (cm)	100.04	11.52	10.49	82.9	19.66
Panicle bearing tillers/plant	9.71	15.57	11.75	56.9	18.26
Panicle length(cm)	21.54	10.9	9.49	75.7	17.01
Spikelets number	124.32	14.4	13.42	86.9	25.78
Grains / panicle	97.37	14.48	13.86	91.6	27.33
Spikelet fertility (%)	79.52	6.35	5.3	69.6	9.12
Test weight(g)	24.21	9.94	7.42	55.7	11.41
Biological Yield(g)	40.73	14.5	11.71	65.3	19.49
Harvest Index(g)	35.69	15.24	11.61	58	18.23
Grain yield(g)	14.33	11.92	10.15	72.5	17.80

Table 3: Phenotypic correlation under saline condition

Characters	Days to 50% flowering	Plant height (cm)	Panicle bearing tillers/plant	Panicle length	Spikelet No.	Grain per panicle	Spikelet Fertility %	Test weight (g)	Biological Yield/plant (g)	Harvest Index (%)	Grain yield / plant (g)
Days to 50% flowering	1.000	0.197	0.048	-0.179	0.119	0.148	-0.204	-0.068	0.069	-0.285*	-0.276*
Plant height (cm)		1.000	-0.101	0.273*	-0.040	-0.017	-0.224	-0.034	-0.058	0.017	-0.052
Panicle bearing tillers/plant			1.000	0.030	-0.143	0.206	0.175	0.048	0.287*	-0.018	0.369**
Panicle length				1.000	-0.176	-0.080	0.052	0.334**	0.236	0.011	0.274*
Spikelet No.					1.000	0.712**	-0.253*	-0.150	0.218	-0.011	0.261*
Grain per panicle						1.000	0.064	-0.224	0.222	0.022	0.293*
Spikelet Fertility %							1.000	0.184	-0.054	0.099	0.024
Test weight (g)								1.000	0.201	0.053	0.281*
Biological Yield/plant (g)									1.000	-0.655**	0.322**
Harvest Index (%)										1.000	0.489**

*, **Significant at 5% and 1% probability levels, respectively.

Table 4: Direct and indirect effect for different characters on grain yield per plant under saline condition at phenotypic level in rice genotypes

Characters	Days to 50% flowering	Plant height (cm)	Panicle bearing tillers/plant	Panicle length	Spikelet No.	Grain per panicle	Spikelet Fertility %	Test weight (g)	Biological Yield/plant (g)	Harvest Index (%)	Grain yield / plant (g)
Days to 50% flowering	-0.027	-0.001	0.006	-0.005	0.012	-0.007	0.008	-0.001	0.071	-0.332	-0.276*
Plant height (cm)	-0.005	-0.006	-0.013	0.007	-0.004	0.001	0.009	0	-0.059	0.02	-0.052
Panicle bearing tillers/plant	-0.001	0.001	0.126	0.001	-0.015	-0.01	-0.007	0	0.295	-0.021	0.369**
Panicle length	0.005	-0.002	0.004	0.026	-0.018	0.004	-0.002	0.003	0.242	0.013	0.274*
Spikelet No.	-0.003	0	-0.018	-0.005	0.101	-0.035	0.01	-0.001	0.224	-0.012	0.261*
Grain per panicle	-0.004	0	0.026	-0.002	0.072	-0.049	-0.003	-0.002	0.228	0.026	0.293*
Spikelet Fertility %	0.006	0.001	0.022	0.001	-0.026	-0.003	-0.039	0.001	-0.055	0.115	0.024
Test weight (g)	0.002	0	0.006	0.009	-0.015	0.011	-0.007	0.008	0.206	0.062	0.281*
Biological Yield/plant (g)	-0.002	0	0.036	0.006	0.022	-0.011	0.002	0.002	1.028	-0.762	0.322**
Harvest Index (%)	0.008	0	-0.002	0	-0.001	-0.001	-0.004	0	-0.673	1.163	0.489**

R Square = 0.9742 Residual effect = 0.1606

Bold values show direct and normal values show indirect effect.

CONCLUSION:

The analysis of variance shows there is presence of sufficient variability among rice genotypes under saline condition. Values of phenotypic coefficient of variation were higher than genotypic coefficient of variation for all the eleven characters under saline condition. Higher magnitudes of phenotypic coefficient of variation were found for panicle bearing tillers/plant, harvest index, biological yield/plant, grain/panicle and spikelet number. High estimates of heritability were observed for grain per panicle, days to 50% flowering, spikelet number, plant height. Whereas medium estimates of heritability were observed for grain yield per plant, spikelet fertility, biological yield per plant, harvest index, panicle bearing tillers per plant and test weight. Under saline condition grains/panicle showed highest value of genetic advance and spikelet fertility showed lowest value of genetic advance. Grain yield showed highly significant and positive phenotypic correlation with harvest index, panicle bearing and biological yield/plant. Harvest index showed maximum positive direct effect on grain yield followed by biological yield/plant and panicle bearing tillers/plant.

References:

- Adhikari, B. N., Joshi, B. P., Shrestha, J., & Bhatta, N. R. (2018). Genetic variability, heritability, genetic advance and correlation among yield and yield components of rice (*Oryza sativa* L.). *J. of Agri. and Natural Res.*, **1(1)**:149–160.
- Beena, R., Veena, V., Jaslam, M.P.K.(2021) Germplasm innovation for high-temperature tolerance from traditional rice accessions of Kerala using genetic variability, genetic advance, path coefficient analysis and principal component analysis. *J. Crop Sci. Biotechnol.* **24**, 555–566.
- Bhargava, K., Shivani, D., Phushpavalli, S.N.C.V.L., Sundaram, R.M., Beulah, P. and Senguttuvel, P.(2021). *Electr. J. of Plant Breed.***12(2)**: 549-555.
- Chamar, J.P., Joshi, R.P., Katkani, D., Sahu, D. and Patel, V.(2021) Genetic variability, heritability and correlation coefficient analysis in rice. *The PharmaInnov. J.* **10 (5)**:763-769.

Comment [IA8]: This should briefly state the major findings of the study and do not repeat the results. For example, genotype...is the highest tolerance genotype to the saline condition.

Comment [IA9]: References should be listed in the order that they appear in the text.

- Dhurai, S. Y., Bhati, P. K. and Saroj, S. K. (2014) Studies on genetic variability for yield and quality characters in rice (*Oryza sativa* L.) under integrated fertilizer management. *The Bioscan* **9(2)**: 845-848.
- Kulsum, U., Sarker, U. and Rasul, M.G. (2022). Genetic variability, heritability and interrelationship in salt-tolerant lines of T. Aman rice. *Genetika*, **54(2)**:761-776.
- Kumar, G., Dwivedi, D.K., Maurya, H. and Yadav, V.K. (2018). Correlation and path coefficient analysis of thirteen rice genotypes for grain yield and other yield attributing traits. *J. PharmacognPhytochem***7(3)**:3506-3510.
- Lingaiah, N.Satish , B., Venkanna, V., Rukmini, K. and Hari, Y.(2020).Genetic Variability and Correlation Studies in Yield Traits of Elite Rice (*Oryza sativa* L.) Genotypes. *Ind. J.Pure App. Biosci.* **8(6)**: 359-363.
- Mishra, V.K., Dwivedi, D.K., Sunil, K. and Verma, O.P. (2012). Screening for salinity tolerance at vegetative and reproductive stages in rice (*Oryza sativa* L.). *Environment and Ecology*, **30(2)**:355-360.
- Nihad, S.A.I., Manidas, A.C., Hasan, K., Hasan, M.A., Honey, O. and Latif, M.A.(2021) Genetic variability, heritability, genetic advance and phylogenetic relationship between rice tungro virus resistant and susceptible genotypes revealed by morphological traits and SSR markers. *Curr. Plant Biol.***25**: 2214-6628.
- Rahman, M. A., Haque, M., Sikdar, B., Islam, M. A., &Matin, M. N. (2014). Correlation Analysis of Flag Leaf with Yield in Several Rice Cultivars. *J. of Life and Earth Sci.* **8**:49–54.
- Singh, S.K., Singh, C.M. and Lal, G.M. (2011) Assessment of genetic variability for yield and its component characters in rice (*Oryza sativa* L.) *Res. in Plant Biol.***1(4)**:73-76.