

## **An Indicator Framework for Assessing Socio-Ecological Resilience of Candaba Wetlands, Philippines**

### **Abstract**

Candaba Wetlands is important production landscape and declared as a priority wetland in the Philippines. The socio-ecological system of Candaba Wetlands is characterized by the synergy of agriculture and ecology in providing various ecosystem services and improving the well-being of the residents. However, the wetland is not secured from recurrent disturbances. This paper presents an indicator framework utilized in the analysis of the socio-ecological resilience of Candaba Wetlands. A review of literature related to resilience and Socio-Ecological Production Landscape and Seascapes (SEPLS) served as the backbone in integrating various perspectives. SEPLS resilience highlights the interactions of the ecological and social system. Thus, the indicators identified in this paper is grounded on four areas of SEPLS: (a) ecosystem protection and maintenance of biodiversity; (b) agricultural biodiversity; (c) knowledge, learning and innovation, and (d) social equity and infrastructure. The indicators in each area were further classified according to ecosystem, agricultural and social dimensions of Candaba Wetlands. The framework also offers a way to communicate with the stakeholders in maintaining and enhancing the resilience of the wetland.

*Keywords: resilience, socio-ecological system, Candaba Wetlands, wetland, indicators*

## 1. Introduction

The concept of resilience has become a focal point for many researchers studying the disturbance and change of socio-ecological systems (SES) (Callo-Concha & Ewert, 2014; Ifejika Speranza et al., 2014a; Quandt, 2018). SES are complex networks shaped by the intricate interactions between social systems and ecosystems (Chaffin & Scown, 2018; Cumming & Peterson, 2017; Dennis, 2015; Ives et al., 2007; Knapp et al., 2010; Peterson et al., 2003). These interactions have profound effects on biodiversity and ecological processes within SES (Jeynes-Smith et al., 2023). Moreover, temporal changes in SES can significantly impact the ecosystem services of wetlands (Liu et al., 2007)

From a socio-ecological system perspective, the resilience characteristic of a system allows it to absorb disturbances and reorganize while undergoing change, thereby retaining its function, structure, identity, and (Adger, 2006; Folke, 2006; Gunderson & Holling, 2002; Holling, 1973; Panpakdee & Limnirankul, 2018; Smit & Wandel, 2006; Urruty et al., 2016; Walker et al., 2006). However, due to the complexity and dynamics of interrelations in SES, measuring its resilience poses a significant challenge (Lade et al., n.d.; Tamberg et al., 2020).

To address this, the Japan Satoyama-Satoumi Association (2010) integrated resilience thinking into socio-ecological production landscapes and seascapes (SEPLS). SEPLS refers to the mosaic production landscapes formed by human-nature interaction that sustain human well-being while maintaining ecosystem services. It links the biological, physical, and social components, which are manifested in traditional knowledge and the sustainable use of biodiversity (Boafo et al., 2016; Boafo & Ichikawa, 2016; Fiksel, 2007; Folke et al., 2010; Takeuchi et al., 2016)

In the Philippines, the Candaba Wetlands have been exposed to various recurrent disturbances such as agricultural activities and flooding. Despite these disturbances, this wetland remains a key biodiversity area in the Philippines and is globally important in the East Asian Australasian Flyway of waterfowls. The residents have adapted to the changing landscape through shifting livelihoods, demonstrating the resilience of the wetland (Meyer, n.d.).

The question then arises, "What makes Candaba Wetlands resilient?" Considering the concepts of resilience and SEPLS, we identified indicators suitable to the context of Candaba Wetlands that could be used in measuring or assessing its socio-ecological resilience and to gain a better understanding of the adaptive mechanisms.

## 2. Resilience Defined

Resilience, as a concept, has been a subject of extensive debate and exploration in ecological studies. Holling (1973) introduced and defined resilience as a measure of a system's persistence and its ability to absorb changes in state variables, driving variables, and parameters while maintaining the same relationships with these variables. He contrasted resilience with stability, defining the latter as the system's ability to return to an equilibrium state after a temporary disturbance, resulting in the degree of fluctuation.

The resilience and stability perspectives on system behavior have distinct implications for resource management. While resilience emphasizes the "domains of attraction and the need for persistence," stability focuses on maintaining equilibrium and harvesting nature's excess production with minimal fluctuation (Iglesias & Whitlock, 2020). Some scholars view resilience as the time required for a system to return to an equilibrium or steady state following a perturbation. In this perspective, resilience is measured by how far the system has deviated from equilibrium and the speed at which it returns to its steady state (Schlüter & Herrfahrdt-Pähle, 2011).

Gunderson (2000) posited that resilience is an emergent property of ecosystems, linked to self-organized behavior over time. He associated self-organization with the interaction between structure and process, leading to system development. This perspective aligns with Holling's adaptive cycle's exploitation phase, where a system can absorb a range of disturbances, indicating high ecological resilience. However, when a system reaches its conservative growth limits, it becomes primed for rapid structural changes.

Walker et al. (2004) offered a comprehensive definition of resilience as "the capacity of a system to absorb disturbance and reorganize while undergoing change to retain substantially the same function, structure, identity, and feedbacks." This definition underscores that an ecosystem's ability to reorganize and renew itself after perturbation depends on states and dynamics at various scales. Resilience is dynamic, changing over time, with human actions potentially eroding resilience. Disturbances that were previously absorbed can lead to a regime shift. Thus, resilience encompasses both recovery and change elements, emphasizing adaptive capacity while considering the sensitivity of human and ecological systems (Ibarraran et al., 2008).

Mumby et al. (2014) distinguished between two forms of resilience discussed in literature: ecological and engineering resilience. Ecological resilience considers ecosystems that can evolve towards one or more community types, even without acute disturbance events. While it is ideal for ecosystems at risk of losing their recovery ability and moving to an undesirable state, it can be challenging to quantify (Lu, 2010). In contrast, engineering resilience applies to systems that consistently recover towards the same long-term state. It is beneficial where ecosystem recovery is common but falls short in systems with frequent disturbances since it focuses on an equilibrium state (Schlee et al., 2012).

### **3. The Socio-Ecological Production Landscapes and Seascapes Concept**

#### **3.1. Origin of the SEPLS Concept**

Satoyama, a Japanese term, refers to a mosaic of different ecosystems such as secondary forests, farmlands, irrigation ponds, grasslands, and human settlements (Ichikawa, 2013; JSSA, 2010). This type of landscape provides diverse land uses that offer benefits for people's lives and agricultural production. Beyond provisioning services, the satoyama landscape also provides regulating services such as flood prevention, watershed conservation, landslide and erosion prevention functions, and pollination (Takeuchi et al., 2016). The various environmental conditions of the mosaic of ecosystems harbor more species across the entire landscape. The concept has been extended to satoumi, which encompasses marine and coastal ecosystems. Satoumi refers to the spatial structure of coastal areas and the use and management of fisheries resources within these areas.

These two concepts depict the relationship of the mutually beneficial interaction between humans and the environment, allowing for the sustainable use of resources. This satoyama-satoumi concept, internationally known as traditional socio-ecological production landscapes and seascapes (SEPLS), is a historical model for environmental stewardship and resource management that contributes to human well-being (Hashimoto & Nakamura, 2014; JSSA, 2010).

The Japan Satoyama-Satoumi Assessment (JSSA) focuses on the link between humans and different landscape ecosystems in Japan and how they interrelate with each other. It follows the framework of sub-global assessments developed by the Millennium Ecosystem Assessment. JSSA adopts the ecosystem services conceptual framework developed by MA due to the following reasons: (a) centrality of human well-being in considerations of ecosystem services, (b) recognition of the interdependency, synergy, and trade-offs between ecosystem services and human well-being, and (c)

acknowledgment of different temporal and spatial scales that impact interdependency (Takahashi et al., 2022).

SEPLS represent harmonious human-nature interactions resulting in positive outcomes for both biodiversity and human well-being, thus implying synergies among multiple nature's contributions to people (NCP) are possible. In case studies of ten projects selected from biodiversity hotspots under the GEF-Satoyama Project, it was found that synergies in NCP exist within SEPLS and management interventions that enhanced these synergies were identified. Among the management options that enhanced NCP in SEPLS were food-centered approaches entailing organic agriculture, eco-labelling, branding and improved agricultural practices. Habitat-centered approaches included participatory biodiversity monitoring, ecosystem restoration, co-management and conservation agreements with landowners. Synergies in NCP were generated by integrating these interventions with enabling governance structures and through community empowerment (Takahashi et al., 2022)

### 3.2. Concepts and Characteristics of SEPLS

Socio-Ecological Production Landscapes and Seascapes (SEPLS) are dynamic mosaics of habitats and land and sea areas shaped by the interactions between people and the environment over time. These interactions not only maintain biodiversity but also provide a bundle of ecosystem services that contribute to human well-being ((Ciftcioglu, 2016; Gu & Subramanian, 2014)).

SEPLS are “*found in many parts of the world, of different socio-economic context, where humans are directly linked with nature through various forms of resource use, including primary industries, subsistence activities, tourism, and cultural uses*” (Gu & Subramanian, 2014). In SEPLS, biodiversity and ecosystems support humans by delivering ecosystems services such as food, fuels, shelter, cultural values, while in return, humans support biodiversity and ecosystems through managing the natural resources for its sustainable use (Subramanian et al., 2017).

The SEPLS concept underscores the importance of traditional knowledge, which is instrumental in building sustainable management strategies for the further development of the landscape. The socio-ecological characteristics of SEPLS vary depending on their location, as their sustainable use and management are influenced by the local inhabitants' traditional norms, beliefs, practices, and customs (Boafo et al., 2016; Kadoya & Washitani, 2011).

SEPLS is also characterized with enhanced biodiversity due to mosaic ecosystems that can support organisms. The SEPLS concept aligns with the core objective of the Intergovernmental Science-Policy Platform for Biodiversity and Ecosystem Services (IPBES), which aims to understand ecosystem production, function, and services to balance conservation and development goals, integrate scientific findings into policies, and improve human well-being (Indrawan et al., 2014).

From a sustainability perspective, SEPLS shares the basic premise of socio-ecological systems and resilience thinking, which posits that social and ecological systems are interconnected, and adaptive management is necessary to address ecosystem changes (Gu & Subramanian, 2014). SEPLS not only provide ecosystem services for local communities but also for larger populations beyond their borders. The production and harvesting processes within SEPLS are influenced by external demands, pressures, and policy decisions at national and international levels (Gu & Subramanian, 2014; Hashimoto & Nakamura, 2014). SEPLS are dynamic in nature and are sustained through adaptive co-management regimes, with interlinkages amongst natural and cultural processes that operate on different time frames (Gu & Subramanian, 2014).

UNDER PEER REVIEW

### 3.3. Changes in SEPLS

SEPLS across the globe have been experiencing a rapid decline due to a multitude of anthropogenic and natural drivers, jeopardizing their sustainability. These influential factors are largely dependent on the socio-economic context of each region (Gu & Subramanian, 2014; Ichikawa, 2013; JSSA, 2010).

The changes in SEPLS are primarily shaped by the interactions between people and nature. Several factors have been identified as drivers of these changes. Indirect factors include economics, culture and religion, science and technology, population, and public policy. Direct factors encompass changes in land use, underuse, overhunting/overharvesting, climate change, the introduction of non-native species, and pollution (JSSA, 2010).

These drivers can be further categorized into socio-political transformation, legal rights and autonomy, economic globalization, and demographic and socio-cultural changes (Gu & Subramanian, 2014). The manifestation of these drivers varies across different regions, as illustrated by the following case studies:

In Africa, SEPLS face threats due to interrelated natural and anthropogenic factors, including climate change, unsustainable resource extraction, population pressure, and governance and institutional issues. Changing rainfall and temperature patterns have resulted in water scarcity and severe impacts on rain-fed agricultural production, affecting food security. The increasing global demand for mineral resources and fossil fuels has led to the conversion of large tracts of rural landscapes into mining concessions, resulting in adverse impacts on the landscape (Boafo et al., 2016; Mapani & Kribek, 2012)

In Japan, particularly in the Noto Peninsula, the sustainability of SEPLS is threatened by rapid depopulation and an aging local community. These factors have led to the gradual collapse of the balance between humans and nature, causing the abandonment of farmland and forests, and increasing the incidence of human-wildlife conflict (Hashimoto & Nakamura, 2014)

In Cambodia, SEPLS are becoming uniform due to land use change. The 1990s saw a focus on timber production from high-value forests managed by private companies under large-scale forest concessions. This led to high levels of illegal logging and degradation of forest resources, exacerbated by conflicts over rights with local communities who traditionally used to collect natural resources from the concession forests (Ichikawa, 2013)

In the Lefke Region of North Cyprus, the mosaic of forest, maquis, agriculture, coast, and marine ecosystems plays a significant role in the production of ecosystem services. However, this mosaic is threatened by land use change due to the impacts of urbanization and land abandonment. Other factors such as low level of management, migration, lack of job opportunities, and an aging society have led to the transformation of the SEPLS of the Lefke Region (Ciftcioglu, 2016, 2017)

These case studies illustrate the complex interplay of factors contributing to changes in SEPLS. Policies on mining concession in Africa and forestry concession in Cambodia, for instance, have triggered the conversion of rural landscapes. Although these policies aim to improve livelihood and food security and reduce poverty, they have negative impacts on maintaining the SEPLS structures and dynamics (Indrawan et al., 2014) Changes in demography also affect land use change (Gu & Subramanian, 2014). The increasing population has impacted the unsustainable use of natural resources in Africa and the conversion of shifting agriculture to cash crops in Cambodia. Conversely, rapid depopulation and aging of the community have led to the abandonment of SEPLS in rural areas in the Lefke Region of North Cyprus and the Noto Peninsula of Japan.

## 4. The Candaba Wetlands, Pampanga, Philippines

### 4.1. Geographical Overview

The Candaba Wetlands, also known as the Candaba Swamp or Candaba Marsh, is a socio-ecological system that encompasses a diverse range of wetlands within the municipality. This area is recognized as a Key Biodiversity Area and is one of the three sites included in the Northwest Manila Bay regional grouping as reported in the Philippine National Report on Wetlands (2015). Geographically, the wetland is situated approximately 50 km north-north-west of Metro Manila, near the towns of Candaba in Pampanga and San Miguel and San Ildefonso in Bulacan, all in Central Luzon (DENR-PAWB, 2005; Juarez-Lucas et al., 2018; Melendres, 2014)

The Candaba Wetlands spans an area of 32,000 hectares, which reduces to 10,000 hectares during dry seasons. This seasonal wetland comprises a complex of freshwater ponds, wetlands, and marshes, surrounded by seasonally flooded grassland, arable land, and palm savanna on a vast alluvial flood plain. The majority of the wetland is privately owned, which presents unique challenges for conservation efforts (Canning et al., 2023)

### 4.2. Ecological Dimension

#### 4.2.1. Natural Flood Retention

The Candaba Wetlands serve as a natural flood retention basin, holding the wet season overflow from the Maasim, San Miguel, Garlang, Bulu, and Penaranda Rivers, and draining into the Pampanga River (DENR-PAWB, 2005; Juarez-lucas et al., 2016; Juarez-Lucas et al., 2018; B. Shrestha et al., 2014; B. B. Shrestha et al., 2014). It functions as a retarding basin, absorbing most flood flows from the eastern sections of the basin and overflows from the Pampanga River via the Cabiao floodway (Garcia, 2010; B. B. Shrestha et al., 2014). This flood control function is similar to the green infrastructure practices observed in Japan, where flood-control basins are used to reduce flood risk while also providing habitat for biodiversity (Ishiyama et al., 2022)

#### 4.2.2. Habitat to Biodiversity

The Candaba Wetlands bolster a seasonal fishery, aligning with the onset of monsoon floodwaters. Despite the predominant agricultural use of the flood plain, the native vegetation is characterized by water lilies, lotus, water hyacinths, and aquatic reeds. Additionally, there are sporadic patches of *Nypa fruticans* and mangroves in adjacent areas (DENR-PAWB, 2005). In a study by Garcia (2010), 18 aquatic species, predominantly *Cyprinids*, were identified in specific regions of the wetlands. The wetland is also a crucial habitat for migratory waterfowls, particularly during October and November. During these months, the wetland becomes a temporary home for 5,000 to 10,000 migratory birds, including egrets, wild ducks, herons, and kingfishers. This influx coincides with the icy conditions in the Northern Hemisphere. Additionally, the wetland provides sanctuary for diverse wildlife, including small mammals, reptiles, and amphibians. The ecological richness of the Candaba Wetlands can be paralleled with the high Andean wetlands of Peru, which are celebrated for their ecosystem services and contribution to local economies (León et al., 2021).

#### 4.2.3. Historical and Cultural Importance

The Candaba Wetlands, with its rich biodiversity and ecological significance, also holds historical and cultural importance. In 1951, the expansive depth of the wetlands and the nearby wilderness of Mount Arayat provided refuge for the HUKBALAHAP, a communist movement in the Philippines (Fifield, 2014). Furthermore, an archaeological site was uncovered in Donya Simang, situated at the heart of the Candaba Wetlands. This discovery was made by Melendres (2014) and his colleagues during a bird-watching expedition. The site's exposure was a result of the construction of an elevated road leading to a bird camp. Among the artifacts found were indigenous undecorated earthenware, believed to date back to before 1000 AD. Additionally, oriental trade ceramics from China, dating from 1300 – 1600 AD, brown glazed stoneware jars from the 13th to 14th century, and blue and white porcelains from the Hongzi period in the late 15th to 16th century were also discovered.

#### 4.2.4. Tourism

The Candaba Wetlands is a favored destination for both local and international bird watchers and naturalists. Its proximity to Metro Manila and its rich biodiversity make it an ideal spot for nature-oriented outdoor recreation and conservation education. The wetland is renowned for its duck products, particularly salted eggs and the exotic delicacy "balut" or boiled unhatched duck eggs. Recognizing the wetland's dual role as a fishing ground and a stopover for migratory birds, the Local Government Unit of Candaba initiated the "Ibun-Ebun (Bird-Egg) Festival" (Melendres, 2014). Such initiatives highlight the potential of the wetlands for ecotourism. In fact, protected areas, due to their recognized natural, ecological, and cultural values, are valuable resources for the rural economy and play a crucial role in the tourism industry (Grgić et al., 2021). Adopting a sustainable tourism model for the Candaba Wetlands can not only elevate its stature but also champion its preservation (Amara, 2017)

### 4.3. Agriculture and Fisheries Dimension

#### 4.3.1. Agriculture and Wild Fisheries

At the municipal level, agriculture and fisheries are the primary sources of income for 70% and 13% of the total households, respectively. During the dry season, from late November to April, the area is converted into rice fields and watermelon plantations. During the rainy season, the impoundment is used as a fishpond and supports a seasonal fishery (Garcia, 2010; Juarez-lucas et al., 2016). Most of the privately-owned flood plain has been converted to agricultural and residential land except for the core area of approximately 500 ha. Since floods are recurrent in the area, livestock and poultry production in Candaba Wetlands are slowly increasing. The locals raise carabaos or water buffaloes, pigs, chickens, ducks, goats, sheep, and cows.

#### 4.3.2. Cropping Season

Cropping season varies within the three subdivisions of the municipality. In Riverside areas, farmers only experience one cropping season which falls January to April or May due to imminent flooding during months of June to November. The farmers from Tagalog and Kapampangan regions enjoy two cropping seasons. The first cropping starts from January to April or May and the second cropping falls from October to December.

#### 4.3.3. Farmers and Fisherfolks

The 2018 data from the Municipal Agriculture Office listed 7,519 farmers were recorded during dry season and 5,492 during the wet season. Barangay Mandili had the highest number of farmers in both seasons. A total of 2,434 fisherfolks were also recorded in Candaba and mostly from Barangays San Agustin, Mandili and Bambang.

The sustainability of these agricultural and fisheries practices is closely linked to the health of the Candaba Wetlands ecosystem. As Ramachandra (2022) notes, the integrity of the catchment area, which includes vegetation, plays a crucial role in maintaining soil moisture and water availability, both of which are essential for farming and fisheries. In areas with more than 60% vegetation of native species, streams are perennial, and soil moisture is higher, facilitating the farming of commercial crops with higher economic returns to the farmers. In contrast, streams are intermittent or seasonal in catchments dominated by monoculture plantations or with vegetation cover lower than 30%. This highlights the need to maintain ecosystem integrity to sustain water availability and, by extension, agricultural and fisheries productivity in the Candaba Wetlands.

#### 4.4. Social Dimension

##### 4.4.1. Demography

The Municipality of Candaba had a total population of 102,400 and a total of 20,211 households with a mean household size of five members as of the 2010 Philippine Census. The majority, 67%, of the population resided in rural areas (Juarez-lucas et al., 2016; Juarez-Lucas et al., 2018) This rural lifestyle, often characterized by close-knit communities and reliance on natural resources, can contribute to a unique social dynamic that is integral to the identity of the Candaba Wetlands (Rao. M. & Pasala, 2019).

##### 4.4.2. Economic Activities

The Municipal Treasury Office recorded a total of 613 establishments registered in Candaba in 2018, with 25% being new. These establishments were predominantly located in Barangay Bahay Pare and the Poblacion area, which are the most populous and commercial areas, respectively. The service sector, mainly engaged in retail trade and general services, comprised a significant portion of the economy. The presence of 13 associations and 15 cooperatives also indicated a strong sense of community and mutual support among the residents (Antipova & Momeni, 2021).

##### 4.4.3. Health

Health services in Candaba were provided by one hospital/infirmarary, three rural health units, 31 barangay health stations, and seven private clinics. Acute respiratory infection was the leading cause of morbidity, followed by hypertension, diabetes mellitus, and urinary tract infection. The availability and accessibility of these health services are crucial for the well-being of the residents, particularly the elderly, who require more frequent health care services (Mohammadi et al., 2021)

The social dimension of the Candaba Wetlands is a complex interplay of demographic, economic, and health factors. Understanding these factors is crucial in planning and implementing sustainable development initiatives in the area (Estebansari et al., 2020)

## 5. Conceptualizing the socio-ecological resilience of Candaba Wetlands

The Socio-Ecological Production Landscapes and Seascapes (SEPLS) framework offers a unique approach to understanding socio-ecological systems, treating both social and ecological dimensions with equal depth (Binder et al., 2013; Ciftcioglu, 2017)). Recognizing the intricate and dynamic nature of SEPLS, indicators have been developed to capture the essential elements that underpin a resilient landscape. While a set of indicators exists, they are not rigid measurements but rather a guide to understand and bolster SEPLS resilience (UNU-IAS, 2013; UNU-IAS et al., 2009; UNU-IAS & IGES, 2015). These indicators can be adapted and localized based on the specific SEPLS in question.

SEPLS resilience encompasses four primary areas: (1) ecosystem protection and maintenance of biodiversity; (2) agricultural biodiversity; (3) knowledge, learning and innovation; (4) social equity and infrastructure. But these areas can be further grouped according to its respective dimension.

### a. Ecosystem Dimension

Ecosystem protection and maintenance of biodiversity could enhance the resilience of an ecosystem. The diversity and interactions of species in Candaba Wetlands reflects the ecosystem health (Fischman, 2004). Biodiversity, encompassing variation from within species to across landscapes, is crucial in the resilience of underpinning ecosystem functions and the services (Convention on Biological Diversity, 1992; Levin, 2013; Oliver et al., 2015). By protecting the ecosystem and maintaining the biodiversity, the community settling in the Candaba Wetlands could gain from the ecosystem services which are either sustained or degraded by the human practices and the institution that regulate the use of natural resources (Yao et al., 2015). Spatial heterogeneity and multifunctionality of the landscape create a mosaic of habitats, thus, increasing the biodiversity (Benton et al., 2003; Oliver et al., 2015; Sutti, 2016). Designating protected areas is also critical in conserving biodiversity as this measure safeguard not only the species, but also the habitat (BMB, 2015; CBD, 2008). The rate of recovery from the extreme environmental and climate change-related stresses and shocks is also vital as this will determine the ability of the Candaba Wetlands to bounce back to its original state (Adger, 2005; UNU-IAS, 2013).

### b. Agricultural Dimension

The agricultural biodiversity in Candaba Wetlands creates the nexus of its ecosystem health and the community (UNU-IAS, 2013). This is linked by the array of functions such as maintenance of agroecosystem functions, sustainable production of food and other agricultural products, and biological support to production (Convention on Biological Diversity, 2013; Cromwell et al., 1999; FAO, 2019; GIZ, 2014). Thus, it is essential to maintain, document and conserve the wetland's agricultural biodiversity. To conserve the Candaba Wetlands, a range of different measures such as on-farm management, restricting farming intensity, prevention of farmland abandonment, establishment of gene bank, botanical garden, agricultural research stations and tissue culture collections, and following eco-agriculture practices (Fransiska et al., 2015; GIZ, 2014; Kleijn et al., 2011) could be conducted. The agricultural biodiversity could also be improved by involving and recognizing the role of the farmers in conservation (Abay et al., 2009), thereby, also have a positive impact in the adaptive capacity of Candaba Wetlands (Dennis, 2015). Horizontal networks such as intercommunity resource trade could also enhance the resilience to perturbation, particularly climate variability.

### c. Social Dimension

The social dimension of Candaba Wetlands encompasses knowledge, learning and innovation; and social equity and infrastructure. The resilience of a community is strengthened by learning from different knowledge systems such as traditional and local adaptation strategies, customs, agricultural traditions, local languages, cultural values, and social institutions (UNU-IAS et al., 2009). Though the local knowledge is considered outdated, local knowledge is built upon the practical knowledge of the past experience and delivered from skilled generations (Panpakdee & Limmirankul, 2018). These knowledge systems that shaped the Candaba Wetlands is strongly linked the community and ecosystem. Maintenance of the local knowledge is significantly anchored by the ability of the community to document local knowledge (Ifejika Speranza et al., 2014b; Panpakdee & Limmirankul, 2018), access and exchange of local knowledge (Ifejika Speranza et al., 2014b; Panpakdee & Limmirankul, 2018; Subramanian et al., 2017), transmit local knowledge to younger generations (Elgar, 2013; Panpakdee & Limmirankul, 2018; UNU-IAS, 2013), uphold cultural traditions related to biodiversity (Ducusin et al., 2019; UNU-IAS, 2013), interact with the landscape, use of terminology or indigenous languages (UNESCO, 2003), and maintain diversity of local food system (Bioversity International, 2019) and local medical system (Alves & Alves, 2011; Nascimento et al., 2016).

The resilience of Candaba Wetlands is also reliant on the availability of and access to functioning social infrastructures, such as communication, health, and education. The resilience of the SEPLS could also improve with effective local resource governance with efficient institutional devolution and institutional arrangements (Senga, 2001; World Bank, 2003) that distinguishes property rights and autonomy in relation to land and resource management (Young, 2011). Fostering gender equality could also build and enhance resilience in the community as the women's leadership and decision-making are empowered and recognized (Chanamoto & Hall, 2015; Collord, 2018; Jenson, 2015). Resilience in the Candaba Wetlands is also dependent on the adaptive mechanism to disasters (UNISDR, 2015) as flooding is recurrent in the area. As a hotspot for migratory birds, ecotourism also has a contribution to resilience of the wetland (Hall, 2013) as this nature-based type of tourism was considered a solution to the disconnect between sustainability and development (Kiper, 2013).

In conclusion, the social dimension of the Candaba Wetlands is a complex interplay of demographic, economic, and health factors. Understanding these factors is crucial in planning and implementing sustainable development initiatives in the area (Estebesari et al., 2020). Accordingly, a set of suitable resilience assessment indicators for the ecological, social and agricultural systems of the Candaba Wetlands was developed by reviewing relevant references. The suitable indicators are detailed in table below:

Table 1. Resilience Indicators for Candaba Wetlands

Area	Indicator	Variable	References
<b>Ecosystem Dimension</b>			
Ecosystem Protection and Maintenance of Biodiversity	Flora and fauna	Degree of decline in biodiversity	(Fischman, 2004; UNU-IAS, 2013)
	Spatial heterogeneity and multifunctionality	Number of ecosystems and land uses	(Benton et al., 2003; Sutti, 2016; UNU-IAS, 2013)
	Areas protected for cultural importance	Number of designated protected area	(UNU-IAS, 2013)

<b>Area</b>	<b>Indicator</b>	<b>Variable</b>	<b>References</b>
	Rate of recovery	Length of recovery time	(Adger, 2005; UNU- IAS, 2013)
<b>Agricultural Dimension</b>			
Agricultural Biodiversity	Crops cultivated	Number of crops cultivated	(UNU-IAS, 2013)
	Animals domesticated	Number of animals domesticated	(Matsushita et al., 2016; UNU-IAS, 2013)
	Maintenance of agricultural biodiversity	Use of local varieties and breeds	(Convention on Biological Diversity, 2013; UNU-IAS, 2013)
	Documentation and conservation of agricultural biodiversity	Degree of documentation and conservation of agricultural biodiversity	(Fransiska et al., 2015; GIZ, 2014; Kleijn et al., 2011; UNU-IAS, 2013)
	Innovation in agricultural biodiversity management	Degree of community reception to changes	(Abay et al., 2009; UNU-IAS, 2013)
	Access and exchange of agricultural biodiversity	Presence of networks of exchange and access of agricultural biodiversity	(UNU-IAS et al., 2009)
<b>Social Dimension</b>			
Knowledge, learning and innovation	Transmission of local knowledge	Degree of intergenerational transmission of knowledge	(Campilan, 2002; Elgar, 2013; Panpakdee and Limmirankul, 2018; UNU-IAS, 2013)
	Number of generations interacting with the landscape	Number of generations interacting with the landscape	(Elgar, 2013; UNU- IAS, 2013)
	Cultural activities related to biodiversity	Presence of cultural activities and community involvement	(Ducusin et al., 2019; UNU-IAS, 2013)
	Practices of documentation and exchange of knowledge	Presence of institutions and systems for documentation and exchange of knowledge	(Ifejika Speranza et al., 2014; Panpakdee and Limmirankul, 2018; UNU-IAS, 2013)
	Use of local terminologies	Degree of use of local terminologies	(UNESCO, 2003; UNU-IAS, 2013)
	Practice of traditional healing methods	Degree of practice of traditional healing methods	(Nascimento et al., 2016)
	Maintenance of local food diversity and system	Degree of local food diversity and system	(UNU-IAS, 2013; Wahlqvist, 2003)
	Social Equity and Infrastructure	Dependence on natural resources	Degree of dependence on natural resources
Local resource governance		Presence of institutions and programs governing the area	(Senga, 2001; UNU- IAS, 2013)

Area	Indicator	Variable	References
	Gender equality	Degree of women's participation in community activities	(Chanamoto and Hall, 2015; Jenson, 2015; UNU-IAS, 2013)
	Clear ownership and autonomy in relation to land resource management	Degree of land tenure and autonomy	(DFID, 1999; UNU-IAS, 2013)
	Social infrastructures	Presence and access to social infrastructures	(DFID, 1999; UNU-IAS, 2013)
	Health care	Access to health care	(UNU-IAS, 2013)
	Livelihood diversity	Presence of on-farm and off-farm livelihood	(DFID, 1999; Panpakdee and Limmirankul, 2018; UNU-IAS, 2013)
	Access to credit and loans	Presence and access to credit and loans	(DFID, 1999; UNU-IAS, 2013)
	Local disaster management	Presence of local disaster management plans and equipment	(Cutter et al., 2010; UNISDR, 2015)
	Ecotourism	Presence of ecotourism policies and activities	(Gallaher, 2007; Kiper, 2013)

## 6. Conclusion

This paper delves into the socio-ecological resilience of the Candaba Wetlands, anchoring its analysis on the Socio-Ecological Production Landscape and Seascape (SEPLS) framework. The SEPLS model, renowned for its participatory nature, offers a comprehensive lens to decipher the intricate interplay between the wetland's ecological and social systems. By adopting the SEPLS approach for the Candaba Wetlands, we aim to harness its inherent participatory methodology to glean insights into the dynamics of these intertwined systems.

One of the salient features of the SEPLS model is its recommendation to employ the Likert scale for assessment. This scale offers a straightforward method to quantitatively encapsulate individuals' perceptions and viewpoints (Ifejika Speranza et al., 2014; UNU-IAS, 2013). Beyond merely capturing impressions, this measurement scale can be instrumental in monitoring the roles and contributions of various stakeholders. Specifically, it can track their efforts in bolstering the resilience of the wetland's socio-ecological system, especially during times of disturbances or perturbations (Jain, 2012).

Furthermore, the SEPLS model underscores the pivotal role of knowledge systems. In the context of the Candaba Wetlands, the anthropogenic activities and their subsequent impacts are intrinsically linked to the knowledge and learning paradigms of the local inhabitants. This recognition of the knowledge system's significance accentuates the need for continuous learning and adaptation, especially in the face of evolving challenges.

The SEPLS approach fosters collaboration. By leveraging this model, key stakeholders from the Candaba Wetlands can convene, fostering a collaborative spirit. Such collaboration can pave the way for

the joint formulation of policies and tools that resonate with the multifaceted dimensions of the wetland, ensuring a holistic approach to its conservation and sustainable development.

UNDER PEER REVIEW

## 7. References

- Abay, F., Gandarillas, E., Shrestha, P., Waters-Bayer, A., & Wongtschowski, M. (2009). Recognising and enhancing local innovation in managing agricultural biodiversity. *Prolinnova Working Paper, September*, 1–13.
- Adger, W. N. (2005). Social-Ecological Resilience to Coastal Disasters. *Science*, *309*(5737), 1036–1039. <https://doi.org/10.1126/science.1112122>
- Adger, W. N. (2006). Vulnerability. *Global Environmental Change*, *16*(3), 268–281. <https://doi.org/10.1016/j.gloenvcha.2006.02.006>
- Alves, R. R. N., & Alves, H. N. (2011). The faunal drugstore: Animal-based remedies used in traditional medicines in Latin America. *Journal of Ethnobiology and Ethnomedicine*, *7*, 1–43. <https://doi.org/10.1186/1746-4269-7-9>
- Amara, D. F. (2017). Responsible marketing for tourism destinations: Saint Catherine Protectorate, South Sinai, Egypt. In *www.jbrmr.com A Journal of the Academy of Business and Retail Management* (Vol. 11). ABRM. [www.jbrmr.com](http://www.jbrmr.com)
- Antipova, A., & Momeni, E. (2021). Unemployment in Socially Disadvantaged Communities in Tennessee, US, During the COVID-19. *Frontiers in Sustainable Cities*, *3*. <https://doi.org/10.3389/frsc.2021.726489>
- Benton, T. G., Vickery, J. A., & Wilson, J. D. (2003). Farmland biodiversity: Is habitat heterogeneity the key? *Trends in Ecology and Evolution*, *18*(4), 182–188. [https://doi.org/10.1016/S0169-5347\(03\)00011-9](https://doi.org/10.1016/S0169-5347(03)00011-9)
- Binder, C. R., Pahl-wostl, C., Hinkel, J., & Bots, P. W. G. (2013). Comparison of Frameworks for Analyzing Social-ecological Systems. *Ecology and Society*, *18*(4). <https://doi.org/10.5751/ES-05551-180426>
- Biodiversity International. (2019). *Diet diversity for nutrition and health*. [https://www.biodiversityinternational.org/fileadmin/user\\_upload/research/research\\_portfolio/Diet\\_diversity/Biodiversity\\_International\\_Dietary\\_Diversity.pdf](https://www.biodiversityinternational.org/fileadmin/user_upload/research/research_portfolio/Diet_diversity/Biodiversity_International_Dietary_Diversity.pdf)
- BMB. (2015). *Guidebook to Protected Areas of the Philippines*. DENR.
- Boafo, Y. A., & Ichikawa, K. (2016). *Socio-ecological Production Landscapes and Seascapes in Africa* (Y. A. Boafo & I. Kaoru, Eds.). United Nations University Institute for the Advanced Study of Sustainability.
- Boafo, Y. A., Kaoru, I., Dunbar, W., & Manago, C. (2016). Understanding the current status, trends and efforts at revitalization of socio-ecological production landscapes and seascapes in Africa. In Y. A. Boafo & I. Kaoru (Eds.), *Socio-ecological Production Landscapes and Seascapes in Africa* (p. 106). UNU-IAS & IR3S/UTIAS 2016.
- Callo-Concha, D., & Ewert, F. (2014). Using the Concepts of Resilience, Vulnerability and Adaptability for the Assessment and Analysis of Agricultural Systems. *Change and Adaptation in Socio-Ecological Systems*, *1*(1), 1–11. <https://doi.org/10.2478/cass-2014-0001>

- Canning, A. D., Smart, J. C. R., Dyke, J., Curwen, G., Hasan, S., & Waltham, N. J. (2023). Constructed Wetlands Suitability for Sugarcane Profitability, Freshwater Biodiversity and Ecosystem Services. *Environmental Management*, 71(2), 304–320. <https://doi.org/10.1007/s00267-022-01734-4>
- CBD. (2008). Protected Areas in Today's World: Their Values and Benefits for the Welfare of the Planet. In *CBD Technical Series No. 36* (Issue 36). Secretariat of the Convention on Biological Diversity.
- Chaffin, B. C., & Scown, M. (2018). Social-ecological resilience and geomorphic systems. *Geomorphology*, 305, 221–230. <https://doi.org/10.1016/j.geomorph.2017.09.038>
- Chanamoto, N. J. C., & Hall, S. J. G. (2015). Gender equality, resilience to climate change, and the design of livestock projects for rural livelihoods. *Gender and Development*, 23(3), 515–530. <https://doi.org/10.1080/13552074.2015.1096041>
- Ciftcioglu, G. C. (2016). Revealing major terrestrial- and marine species-based provisioning ecosystem services provided by the socio- ecological production landscapes and seascapes of Lefke Region in North Cyprus. *Environment, Development and Sustainability*. <https://doi.org/10.1007/s10668-016-9877-5>
- Ciftcioglu, G. C. (2017). Assessment of the relationship between ecosystem services and human wellbeing in the social-ecological landscapes of Lefke Region in North Cyprus. *Landscape Ecology*, 32(4), 897–913. <https://doi.org/10.1007/s10980-017-0494-y>
- Collord, G. (2018). *Policy seminar : Building resilience by fostering gender equity at the household level* Event Page : <http://www.ifpri.org/blog/policy-seminar-building-resilience-fostering-gender-equity-household-level>
- Convention on Biological Diversity, 30 (1992).
- Convention on Biological Diversity. (2013). *What is Agricultural Biodiversity?* <http://www.cbd.int/agro/whatis.shtml>
- Cromwell, E., Cooper, D., & Mulvany, P. (1999). Agriculture, biodiversity and livelihoods: issues and entry points. *Natural Resources Perspectives*, 38, 1–34.
- Cumming, G. S., & Peterson, G. D. (2017). Unifying Research on Social–Ecological Resilience and Collapse. *Trends in Ecology and Evolution*, 32(9), 695–713. <https://doi.org/10.1016/j.tree.2017.06.014>
- Dennis, M. (2015). *The role of community-led innovation in the adaptive capacity of ecosystem services in an urban social-ecological system*. June.
- DENR-PAWB. (2005). *National Report on Wetlands - Philippines*. DENR.
- Ducusin, R. J. C., Espaldon, M. V. O., Rebanco, C. M., & De Guzman, L. E. P. (2019). Vulnerability assessment of climate change impacts on a Globally Important Agricultural Heritage System (GIAHS) in the Philippines: the case of Batad Rice Terraces, Banaue, Ifugao, Philippines. *Climatic Change*, 153, 395–421. <https://doi.org/10.1007/s10584-019-02397-7>
- Elgar, G. (2013). Transmission of Traditional Agricultural Knowledge : Intergenerational or International ? Examining Youth's Involvement in Agriculture. *Independent Study Project (ISP) Collection*.

- Estebansari, F., Jalili-Bahabadi, F., Rohani, C., Rahimi Khalifeh Kandi, Z., & Mostafaei, D. (2020). Socio-economic determinants affecting the lifestyle of pregnant women. *Iranian Journal of Health Education and Health Promotion*, 8(3), 261–271. <https://doi.org/10.29252/ijhehp.8.3.261>
- FAO. (2019). *Agricultural Biodiversity*. FAO.
- Fifield, R. H. (2014). *Institute of Pacific Relations The Hukbalahap Today* (Vol. 20, Issue 2). Institute of Pacific Relations.
- Fiksel, J. (2007). Sustainability and Resilience: Toward a Systems Approach. *IEEE Engineering Management Review*, 35(3), 5–5. <https://doi.org/10.1109/emr.2007.4296420>
- Fischman, R. L. (2004). The meanings of biological integrity, diversity, and environmental health. *Natural Resources Journal*, 44(4), 989–1026.
- Folke, C. (2006). Resilience: The emergence of a perspective for social-ecological systems analyses. *Global Environmental Change*, 16(3), 253–267. <https://doi.org/10.1016/j.gloenvcha.2006.04.002>
- Folke, C., Carpenter, S. R., Walker, B., Scheffer, M., Chapin, T., & Rockstrom, J. (2010). Resilience Thinking: Integrating Resilience, Adaptability and Transformability. *Ecology and Society*, 15(4), 20. <http://www.ccologyandsociety.org/vol15/iss4/art20/%0AInsight>
- Fransiska, G. A., Rahajeng, S., Dewi, K., & González, V. A. (2015). Conserving Traditional Seed Crops Diversity. *Global Sustainable Development Report*, 1–8.
- Garcia, L. M. B. (2010). Species composition and length-weight relationship of fishes in the Candaba wetland on Luzon Island, Philippines. *Journal of Applied Ichthyology*, 26(6), 946–948. <https://doi.org/10.1111/j.1439-0426.2010.01516.x>
- GIZ. (2014). Financing the conservation of agricultural diversity. *People, Food, and Biodiversity*.
- Grgić, I., Zrakić Sušac, M., Jež Rogelj, M., Kunštović, L., Perčin, A., Mikuš, O., Kovačiček, T., & Hadelan, L. (2021). Respondents' attitudes towards tourism in protected natural areas: the example of Lonjsko polje. *Journal of Central European Agriculture*, 22(4), 868–880. <https://doi.org/10.5513/JCEA01/22.4.3232>
- Gu, H., & Subramanian, S. M. (2014). Drivers of Change in Socio-Ecological Production Landscapes : Implications. *Ecology and Society*, 19(1). <https://doi.org/http://dx.doi.org/10.5751/ES-06283-190141>
- Gunderson, L. H. (2000). Ecological Resilience-In Theory and Application. *Annu. Rev. Ecol. Syst.*
- Gunderson, L. H., & Holling, C. S. (2002). *Panarchy* (L. H. Gunderson & C. S. Holling, Eds.). Island Press.
- Hall, M. C. (2013). Tourism and the implementation of the Convention on Biological Diversity. *International Governance on Environmental Issues*, 111–118. [https://doi.org/10.1007/978-94-015-8826-3\\_7](https://doi.org/10.1007/978-94-015-8826-3_7)
- Hashimoto, S., & Nakamura, S. (2014). Mapping and characterizing ecosystem services of social – ecological production landscapes : case study of Noto , Japan. *Sustainability Science*. <https://doi.org/10.1007/s11625-014-0285-1>

- Holling, C. S. (1973). Resilience and Stability of Ecological Systems. *Annual Review of Ecology and Systematics*, 4(1), 1–23. <https://doi.org/10.1146/annurev.es.04.110173.000245>
- Ibarraran, M., Malone, E. L., & Brenk, A. L. (2008). *Climate Change Vulnerability and Resilience : Current Status and Trends for Mexico ME Ibarrarán* (Issue December). Pacific Northwest National Laboratory.
- Ichikawa, K. (2013). Understanding Socio-Ecological Production Landscapes in the Context of Cambodia. *International Journal of Environmental and Rural Development*, 4(1), 57–62.
- Ifejika Speranza, C., Wiesmann, U., & Rist, S. (2014a). An indicator framework for assessing livelihood resilience in the context of social-ecological dynamics. *Global Environmental Change*, 28(1), 109–119. <https://doi.org/10.1016/j.gloenvcha.2014.06.005>
- Ifejika Speranza, C., Wiesmann, U., & Rist, S. (2014b). An indicator framework for assessing livelihood resilience in the context of social-ecological dynamics. *Global Environmental Change*, 28(1), 109–119. <https://doi.org/10.1016/j.gloenvcha.2014.06.005>
- Iglesias, V., & Whitlock, C. (2020). If the trees burn, is the forest lost? Past dynamics in temperate forests help inform management strategies. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375(1794). <https://doi.org/10.1098/rstb.2019.0115>
- Indrawan, M., Yabe, M., Nomura, H., & Harrison, R. (2014). Deconstructing satoyama - The socio-ecological landscape in Japan. *Ecological Engineering*, 64, 77–84. <https://doi.org/10.1016/j.ecoleng.2013.12.038>
- Ishiyama, N., Yamanaka, S., Ooue, K., Senzaki, M., Kitazawa, M., Morimoto, J., & Nakamura, F. (2022). Flood-Control Basins as Green Infrastructures: Flood-Risk Reduction, Biodiversity Conservation, and Sustainable Management in Japan. In *Green Infrastructure and Climate Change Adaptation* (pp. 189–207). [https://doi.org/10.1007/978-981-16-6791-6\\_12](https://doi.org/10.1007/978-981-16-6791-6_12)
- Ives, C., Taylor, M. P., & Davies, P. (2007). Ecological condition and biodiversity value of urban riparian and non-riparian bushland environments : Ku-ring-gai , Sydney. *Proceedings of the 5th Australian Stream Management Conference. Australian Rivers: Making a Difference.*, 163–168.
- Jenson, J. (2015). The Fading Goal of Gender Equality: Three Policy Directions that Underpin the Resilience of Gendered Socio-economic Inequalities. *Social Politics*, 22(4), 539–560. <https://doi.org/10.1093/sp/jxv035>
- Jeynes-Smith, C., Bode, M., & Araujo, R. P. (2023). *Identifying and Explaining the Resilience of Ecological Networks*. <http://arxiv.org/abs/2306.08795>
- JSSA. (2010). *Satoyama - Satoumi Ecosystems and Human Well-Being: Socio-ecological Production Landscapes of Japan - Summary for Decision Makers*. United Nations University.
- Juarez-lucas, A. M., Kibler, K. M., Ohara, M., & Sayama, T. (2016). Benefits of flood-prone land use and the role of coping. *Natural Hazards*, 1–6. <https://doi.org/10.1007/s11069-016-2551-2>

- Juarez-Lucas, A. M., Kibler, K. M., Sayama, T., & Ohara, M. (2018). Flood risk-benefit assessment to support management of flood-prone lands. *Journal of Flood Risk Management*.  
<https://doi.org/10.1111/jfr3.12476>
- Kadoya, T., & Washitani, I. (2011). The Satoyama Index : A biodiversity indicator for agricultural landscapes. *Agriculture, Ecosystems and Environment*, 140(1–2), 20–26.  
<https://doi.org/10.1016/j.agee.2010.11.007>
- Kiper, T. (2013). Role of Ecotourism in Sustainable Development. *Advances in Landscape Architecture*.  
<https://doi.org/10.5772/55749>
- Kleijn, D., Rundlöf, M., Scheper, J., Smith, H. G., & Tscharntke, T. (2011). Does conservation on farmland contribute to halting the biodiversity decline? *Trends in Ecology and Evolution*, 26(9), 474–481.  
<https://doi.org/10.1016/j.tree.2011.05.009>
- Knapp, A. K., Smith, M. D., Magnuson, J. J., Melack, J. M., Collins, S. L., Harlan, S. L., Orenstein, D. E., Grove, J. M., Robertson, G. P., Grimm, N. B., Childers, D. L., Whitmer, A. C., Gragson, T. L., Kofinas, G. P., Carpenter, S. R., Ogden, L. A., McDowell, W. H., Swinton, S. M., & Kaye, J. P. (2010). An integrated conceptual framework for long-term social–ecological research. *Frontiers in Ecology and the Environment*, 9(6), 351–357. <https://doi.org/10.1890/100068>
- Lade, S. J., Walker, B. H., & Haider, L. J. (n.d.). *Resilience as pathway diversity: Linking systems, individual and temporal perspectives on resilience*.
- Levin, S. (2013). *Encyclopedia of Biodiversity* (S. Levin, Ed.; Second). Elsevier Ltd.
- Liu, J., Moran, E., Carpenter, S. R., Ostrom, E., Deadman, P., Lubchenco, J., Folke, C., Pell, A. N., Redman, C. L., Kratz, T., Provencher, W., Taylor, W. W., Liu, J., Ouyang, Z., Alberti, M., Dietz, T., & Schneider, S. H. (2007). Complexity of Coupled Human and Natural Systems. *Science*, 317(5844), 1513–1516.  
<https://doi.org/10.1126/science.1144004>
- Lu, F. (2010). Patterns of Indigenous Resilience in the Amazon: A Case Study of Huaorani Hunting in Ecuador. In *Journal of Ecological Anthropology* (Vol. 14, Issue 1).  
<https://digitalcommons.usf.edu/jea/vol14/iss1/1>
- Mapani, B., & Kribek, B. (2012). Environmental and Health Impacts of Mining in Africa. In B. Mapani & B. Kribek (Eds.), *Annual Workshop IGCP/SIDA* (Issue 594, pp. 1–30). Czech Geological Survey.  
<https://doi.org/10.1016/B978-0-444-52858-2.00007-4>
- Melendres, R. G. (2014). The Utilization of Candaba Swamp from Prehistoric to Present Time : Evidences from Archaeology , History and Ethnography. *Bhatter College Journal of Multidisciplinary Studies*, 4, 81–93. [http://bcjms.bhattercollege.ac.in/V4/09\\_Candaba\\_Swamp.pdf](http://bcjms.bhattercollege.ac.in/V4/09_Candaba_Swamp.pdf)
- Meyer, K. J. (n.d.). *A Dynamical Systems Framework for Resilience in Ecology*.
- Mohammadi, F., Bastani, F., & Hosseini, R. (2021). Successful Aging Focused on the Dimensions of Perceived Social Support and Mental Vitality in the Elderly Referred to the Health Centers of Iran University of Medical Sciences (2020). *Iran Journal of Nursing*, 34(131), 55–69.  
<https://doi.org/10.29252/ijn.34.131.55>

- Mumby, P. J., Chollett, I., Bozec, Y., & Wolff, N. H. (2014). ScienceDirect Ecological resilience , robustness and vulnerability : how do these concepts benefit ecosystem management ? *Current Opinion in Environmental Sustainability*, 7, 22–27. <https://doi.org/10.1016/j.cosust.2013.11.021>
- Nascimento, A. L. B., Lozano, A., Melo, J. G., Alves, R. R. N., & Albuquerque, U. P. (2016). Functional aspects of the use of plants and animals in local medical systems and their implications for resilience. *Journal of Ethnopharmacology*, 194, 348–357. <https://doi.org/10.1016/j.jep.2016.08.017>
- Oliver, T. H., Heard, M. S., Isaac, N. J. B., Roy, D. B., Procter, D., Eigenbrod, F., Freckleton, R., Hector, A., Orme, C. D. L., Petchey, O. L., Proença, V., Raffaelli, D., Suttle, K. B., Mace, G. M., Martín-López, B., Woodcock, B. A., & Bullock, J. M. (2015). Biodiversity and Resilience of Ecosystem Functions. *Trends in Ecology & Evolution*, 30(11), 673–684. <https://doi.org/10.1016/j.tree.2015.08.009>
- Panpakdee, C., & Limnirankul, B. (2018). Indicators for assessing social-ecological resilience: A case study of organic rice production in northern Thailand. *Kasetsart Journal of Social Sciences*, 39(3), 414–421. <https://doi.org/10.1016/j.kjss.2017.07.003>
- Peterson, G. D., Cumming, G. S., & Carpenter, S. R. (2003). Scenario planning: A tool for conservation in an uncertain world. *Conservation Biology*, 17(2), 358–366. <https://doi.org/10.1046/j.1523-1739.2003.01491.x>
- Quandt, A. (2018). Measuring livelihood resilience: The Household Livelihood Resilience Approach (HLRA). *World Development*, 107, 253–263. <https://doi.org/10.1016/j.worlddev.2018.02.024>
- Ramachandra, T. V. (2022). Innovative ecological approaches to ensure clean and adequate water for all. *Journal of Environmental Biology*, 43. <https://doi.org/10.22438/Editorial>
- Rao, M., S., & Pasala, S. R. (2019). Inter-Regional Variations in the Process of Promoting Socio-Economic Empowerment of the Tribal Women through Self Help Groups (SHGs) Activities. *IRA-International Journal of Management & Social Sciences (ISSN 2455-2267)*, 15(2), 36. <https://doi.org/10.21013/jmss.v15.n2.p3>
- Schlee, M. B., Tamminga, K. R., & Tangari, V. R. (2012). A method for gauging landscape change as a prelude to urban watershed regeneration: The case of the Carioca river, Rio de Janeiro. *Sustainability*, 4(9), 2054–2098. <https://doi.org/10.3390/su4092054>
- Schlüter, M., & Herrfahrdt-Pähle, E. (2011). Exploring resilience and transformability of a river basin in the face of socioeconomic and ecological crisis: an example from the Amudarya river basin, Central Asia. *Ecology and Society*, 16(1), 32. <http://www.ecologyandsociety.org/vol16/iss1/art32/>
- Senga, R. G. (2001). Establishing Protected Areas in the Philippines: Emerging trends, challenges and prospects. *Protected Areas in East Asia*, 18(2), 9.
- Shrestha, B. B., Okazumi, T., Miyamoto, M., Nabesaka, S., Tanaka, S., & Sugiura, A. (2014). Fundamental Analysis for Flood Risk Management in the Selected River Basins of Southeast Asia. *Journal of Disaster Research*, 9(5), 858–869.

- Shrestha, B., Okazumi, T., Miyamoto, M., & Sawano, H. (2014). *Development of flood risk assessment method for data-poor river basins: A case study in the Pampanga River Basin, Philippines* (pp. 1–12).
- Smit, B., & Wandel, J. (2006). Adaptation, adaptive capacity and vulnerability. *Global Environmental Change*, 16(3), 282–292. <https://doi.org/10.1016/j.gloenvcha.2006.03.008>
- Subramanian, S. M., Chakraborty, S., & Ichikawa, K. (2017). *Mainstreaming Socio-Ecological Production Landscape Management Approaches*. 8. [http://collections.unu.edu/view/UNU:6026#.W0eqSYPFyG4.mendeley&title=Mainstreaming Socio-Ecological Production Landscape Management Approaches - UNU Collections](http://collections.unu.edu/view/UNU:6026#.W0eqSYPFyG4.mendeley&title=Mainstreaming+Socio-Ecological+Production+Landscape+Management+Approaches+-+UNU+Collections)
- Sutti, F. (2016). *Importance of Agricultural Systems as Multifunctional Landscapes*. University of Vermont.
- Takahashi, Y., Park, K. J., Natori, Y., Dublin, D., Dasgupta, R., & Miwa, K. (2022). Enhancing synergies in nature's contributions to people in socio-ecological production landscapes and seascapes: lessons learnt from ten site-based projects in biodiversity hotspots. *Sustainability Science*, 17(3), 823–836. <https://doi.org/10.1007/s11625-021-00927-w>
- Takeuchi, K., Ichikawa, K., & Elmqvist, T. (2016). Satoyama landscape as social – ecological system : historical changes and future perspective. *Current Opinion in Environmental Sustainability*, 19, 30–39. <https://doi.org/10.1016/j.cosust.2015.11.001>
- Tamberg, L. A., Heitzig, J., & Donges, J. F. (2020). *A modeler's guide to studying the resilience of social-technical-environmental systems*. <http://arxiv.org/abs/2007.05769>
- UNESCO. (2003). *Language Vitality and Endangerment* (Issue March). <https://doi.org/10.1024/1662-9647/a000144>
- UNISDR. (2015). Sendai Framework for Disaster Risk Reduction 2015 - 2030. *Proceedings of the 3rd United Nations World Conference on DRR, Sendai, Japan*, 32. <https://doi.org/A/CONF.224/CRP.1>
- UNU-IAS. (2013). *Indicators of Resilience in Socio-ecological Production Landscapes (SEPLs)*. United Nations University Institute for the Advanced Study of Sustainability.
- UNU-IAS, & IGES. (2015). *Generating collective knowledge on the conservation, management and sustainable use of socio-ecological production landscapes and seascapes - A summary of a review of 80 case studies under the International Partnership for the Satoyama Initiative (IPSI)*. United Nations University Institute for the Advanced Study of Sustainability.
- UNU-IAS, UNDP, & IGES. (2009). *Toolkit for the Indicators of Resilience in socioecological production landscapes and seascapes*. <https://doi.org/10.1596/978-0-8213-8865-5>
- Urruty, N., Tailliez-Lefebvre, D., & Huyghe, C. (2016). Stability, robustness, vulnerability and resilience of agricultural systems. A review. *Agronomy for Sustainable Development*, 36(1), 1–15. <https://doi.org/10.1007/s13593-015-0347-5>

Walker, B., Gunderson, L., Kinzig, A., Folke, C., Carpenter, S., & Schultz, L. (2006). A Handful of Heuristics and Some Propositions for Understanding Resilience in Social-Ecological Systems. *Ecology and Society*, 11(1).

Walker, B., Holling, C. S., Carpenter, S. R., & Kinzig, A. (2004). Resilience, Adaptability and Transformability in Social – ecological Systems. *Ecology and Society*, 9(2).

World Bank. (2003). *Governance of Natural Resources in the Philippines: Lessons from the Past, Directions for the Future* (Issue November).

Yao, J., Sánchez-Pérez, J. M., Sauvage, S., Teissier, S., Attard, E., Lauga, B., Duran, R., Julien, F., Bernard-Jannin, L., Ramburn, H., & Gerino, M. (2015). Biodiversity and ecosystem purification service in an alluvial wetland. *Ecological Engineering*, 103, 359–371.  
<https://doi.org/10.1016/j.ecoleng.2016.02.019>

Young, O. R. (2011). Land use, environmental change, and sustainable development: The role of institutional diagnostics. *International Journal of the Commons*, 5(1), 66–85.  
<https://doi.org/10.18352/ijc.244>

UNDER PEER REVIEW