

## Role of Phenylpropanoids and Flavonoids in Plant defense mechanism

### ABSTRACT:

Phenylpropanoids and flavonoids are specialized metabolites that play a crucial role in plant defenses against biotic and abiotic stresses. Secondary metabolites in plants, such as phenylpropanoids, flavonoids, lignins, monolignols, phenolic acids, stilbenes and coumarins, play a significant role in biotic and abiotic stress responses and interactions with their environment. Under stress-free conditions, flavonoids influence pollen tube growth, seed maturation, dormancy, and the longevity of plant reproductive organs and seeds. Under adverse abiotic conditions, flavonoids can mediate defense responses, such as the accumulation of flavonoids in leaves and glandular trichomes. Metabolomics has identified specialized PPP metabolites correlated with plant resistance to fungi and oomycetes. However, the correlation can be in opposite directions depending on the pathogen and a systemic approach is required to achieve increased resistance without decreasing resistance to another pathogen. The PAL gene is located upstream in the PPP pathway and phenylpropanoids and flavonoids have antifungal activity against *Fusarium oxysporum*, *Rhizoctonia solani* and *Alternaria alternata*. Phenylpropanoids have been studied in vitro to develop biological alternatives to synthetic phytoprotectants but the host's defense system is the most effective mode of action. Post-inoculation transcriptomic shifts reveal that some genes encoding enzymes in the PPP are rapidly activated, while others are induced later in the infection response. Insects are the most problematic macroscopic pests on cultivated plants and plants' natural defense mechanisms against herbivores include releasing volatile organic compounds, attracting enemies and producing phenylpropanoids and flavonoids. Increasing the content of flavonoids in infested plants can reduce the population of insects that cause damage to plants and spread viruses. Plant-environment interaction, mediated by secondary metabolites like phenylpropanoids and flavonoids, leads to diversity in gene structures. Systemic biology approach will be realized through sequencing, structural elucidation, and analytical tools. Model plants and crops can be used for exploration.

**Keywords:** Phenylpropanoids, Flavonoids, Plant defense, Metabolomics, Volatile organic compounds

### 1. INTRODUCTION


"In plants, secondary metabolites perform many biological functions and play an important role in biotic and abiotic stress responses and plant interactions with their environment. Some of the metabolites play a fundamental role in the attraction of pollinators mentioned in chemical ecology, while others help the organism cope with stressful stimuli. At the time of discovery, they were considered secondary metabolites, not essential to plant growth and development. However, their definition has changed significantly over the past two decades" [1,2,3,4,5]. Some authors refer to these metabolites as "specialized" metabolites, while others refer to them as "central" metabolites because they play a key role in plant plasticity and response to various environmental factors [6, 7, 8]. "In addition, high-throughput sequencing has allowed the publication of genomes of a number of species, revealing that genes involved in the biosynthesis pathways of these metabolites are present in a large array of genomes" [9]. "A number of plant specialized metabolites and the enzymes involved in their biosynthesis have been explored large-scale using metabolomics and functional genomics technologies" [10, 11]. An author in a review clearly mentioned these metabolites as "specialized metabolites" because they fully adhere to the concept that these metabolites are of great importance [1].

"PPP leads to the accumulation of many families of compounds, including phenylpropanoids, flavonoids, lignins, monolignols, phenolic acids, stilbenes, and coumarins" [12]. "Subfamilies of flavonoid molecules are categorised based on their structure, such as flavones, isoflavones, anthocyanidins, flavonols, flavanols, flavanones, aurones and chalcones. In each subfamily, there is a wide variety of molecules resulting from various conjugation processes, such as C- or O-methylation, sulfation, or glycosylation. The growth and maturation of pollen tubes, seed maturation, dormancy, and longevity of plant reproductive organs and seeds are affected by flavonoids under stress-free conditions" [13, 14]. "Additionally, they contribute to the attractiveness of plants to pollinators via the colour and scent of their flowers. Also, they facilitate plant-microorganism communication for the establishment of symbiosis, such as rhizobium interactions during nodulation" [14,15]. "When plants are under adverse abiotic conditions, they can be used to mediate defense responses. For example,

under water stress, plants have to deal with oxidative stress caused by reactive oxygen species (ROS). In order to limit the lipid peroxidation of cell membranes, high antioxidant activity is necessary” [16, 17]. The process can be achieved by up regulating the genes involved in the phenolic flavonoid biosynthesis described (Fig. 1).

“In *Chrysanthemum morifolium* L. cultivars under water stress, genes encoding enzymes such as Phenylalanine ammonia-lyase (PAL), Chalcone isomerase (CHI) and flavanone 3-hydroxylase (F3H) were upregulated, resulting in an increase in flavonoids” [18]. Salt and UV stress also resulted in a significant increase in the biosynthesis of flavonoid glycosides and caffeic acid derivatives [19]. “Several genes that code for the Myloblastosis (MYB) transcription factor, Chalcone synthase (CHS), and Chalcone isomerases have been mutated to alter *Arabidopsis thaliana* (L.) Heynh's freeze tolerance” [20]. *Phillyrea latifolia* L. accumulates strong amounts of flavonoids in its leaves and glandular trichomes when exposed to excessive light [25] leading authors to propose that they play a protective role in *P. latifolia*'s integrated mechanisms of acclimation.

“As a keynote, plant defense can be mediated indirectly by flavonoids and phenylpropanoid compounds acting as signalling molecules in biotic stress situations or directly by phytoanticipins (active compounds accumulating in plant tissues) and phytoalexins (newly synthesised active compounds that reduce pathogen detection)” [22, 23, 24]. It has been shown that genes involved in their biosynthesis increase resistance to biotic stress [14, 1]. “This has been described in the literature from two perspectives. The study of basal defense has addressed a variety of constitutive defense mechanisms, including physical barriers and chemical barriers” [26, 27, 28]. “Among these are phenylpropanoid derivatives for cell wall reinforcement. The abundance of phenylpropanoids in Maize (*Zea mays* L.) grain pericarps has been shown to reduce disease symptoms in genotypes resistant to *Fusarium graminearum*, *Fusarium verticillioides*, and Maize weevil (*Sitophilus zeamais* Motsch.)” [28, 29, 30]. “As an alternative, research focused on induced metabolites investigated how these metabolites may suppress or limit invader pathogenicity through their potential resistance toxicity. A growing interest in decoding the molecular dialogue between the host and the pathogen has stimulated the study of this direct effect of bioactive compounds against pathogens. However, these mechanisms are still unknown. Contrary to this, plant-derived compounds-especially flavonoids-have been extensively studied in drug research” [31, 32]. As a result of searching for natural anti-inflammatory compounds, some flavones have been found to have intracellular targets [33].



**Figure 1.** Main steps have the phenylpropanoid and flavonoid pathways. Enzymes mentioned in this paper are shown in green and yellow for central phenylpropanoid pathways and flavonoid biosynthesis, respectively. Complete arrows refer to one step in the biosynthetic pathway, whereas dashed arrows represent undetailed pathways leading to one molecule or molecule subfamilies [25].

## 2. Secondary metabolites defence mechanisms against different pathogens

### 2.1 Fungal pathogens

“Metabolomics has greatly contributed to the identification of specialized PPP metabolites correlated with plant resistance to fungi and oomycetes” [34, 35, 36, 37]. “O-glycosylated flavonoids are the most frequently reported host-pathogen interactions. In most mechanistic studies published so far, phenolic derivative accumulation in plant tissues associated with fungus resistance has been emphasised in terms of quantitative or spatiotemporal aspects”. [59]

“In order to illustrate quantitative aspects, Carrot leaves (*Daucus carota* L.) will be used. A comparison with susceptible genotypes revealed significantly higher levels of feruloylquinic acid, apigenin (4'-O- and 7'-O-glycosides), luteolin, and chrysoeriol in genotypes more resistant to *Alternaria dauci*” [35]. There are similar differences in PPP metabolites between resistant and susceptible genotypes in numerous other ecosystems, supporting plant breeders. In spite of the same combination of factors, disease resistance and metabolite contents can be correlated in opposite directions, depending on the pathogen. A specific example was observed in Potato tubers enriched in rutin and nicotiflorin that were resistant to *Pectobacterium atrosepticum*, but susceptible to *Phytophthora infestans* [38], a biotrophic fungal pathogen. Thus, a systemic approach is required to achieve increased resistance to a specific pathogen without decreasing resistance to another.

“An example of a spatial aspect is the Maize-*Fusarium graminearum* pathosystem and the Maize-*Fusarium verticillioides* pathosystem, where ferulic acid levels in grain tissues were found to correlate with a lesser extent of disease in resistant genotypes” [29, 30]. “To prevent pathogen invasion, this strategy of localising the defense metabolite seems very strategic. For example, in Cotton (*Gossypium hirsutum* L. and *Gossypium barbadense* L.), catechin and gallic acid were dominant near the *Verticillium dahliae* infection site in the vessels, creating a toxic environment that confined the pathogen to the vessel lumens” [39]. “By preventing the spread of vascular disease through the formation of tyloses, this local accumulation contributed to preventing its systemic spread. It has also been observed in Grapevines (*Vitis vinifera* L.) defense against *Phaeoacremonium chlamydospora* and *Phaeoacremonium* species” [40].

“As an example of temporality, the Date palm tree (*Phoenix dactylifera* L.) is resistant to *Fusarium oxysporum* fungal diseases. This resistance has been attributed to a quantitative differential in the concentration of 5-O-caffeoyl-shikimic acid at physiological stage 3 (ripening of dates). Often, the difference in phenotypic characteristics between resistant and susceptible cultivars of fungal diseases is explained by the early or constitutive availability of the compound of interest in the tissues, which results in an efficient defense response, as demonstrated by preformed chlorogenic acid in Tobacco plants resistant to *Cercospora nicotianae*” [25]. “It has also been found that 194 metabolites have been identified in the Barley (*Gibberella zeae*) pathosystem and that they accumulate significantly more in a resistant genotype than in a susceptible genotype” [34]. “The active compounds included kaempferol-3-O-rhamnopyranoside, naringenin-7-O-glucopyranoside, kaempferol-3-O-glucopyranoside, and kaempferol-3-O-glucopyranoside-7-O-rhamnopyranoside-7-O-sophoroside. It is becoming increasingly apparent that constitutive accumulation of flavonoids occurs in various plant-fungus pathosystems as a result of disease resistance” [34, 42, 43]. “Besides an early or constitutive synthesis of the defense compounds, the maintenance of their concentration over time, *i.e.*, when disease pressures are high. As a result, plant resistance to fungal diseases is more complicated than a simple quantitative difference between resistant and susceptible genotypes during infection. Also, the durability of the system is determined by the stability of the metabolite contents in the tissues over time. It is thus necessary to maintain an efficient metabolic ratio according to disease and plant development cycles” [25].

An integrated approach of reverse genetics and biochemical analysis of the resulting proteins has helped explain how phenylpropanoid derivatives mediate fungal disease resistance. First of all, since the PAL gene is located upstream in the PPP pathway, modifications would prevent the plant from synthesising many compounds driven by downstream genes. Tobacco PAL mutants (PAL-suppressed YE-6-16 transformants) exhibited rapid growth of lesions after infection by *Cercospora nicotianae*, while PAL gene overexpression reduced symptoms [25, 44]. The inactivation of the gene encoding a CHS in *Arabidopsis thaliana* decreased anthocyanin content and *Verticillium dahliae* resistance. However, overexpression of genes encoding CHS, CHI, and dihydroflavonol reductase (DFR) in flax (*Linum usitatissimum* L.) has been linked to increased resistance to *Fusarium* species. Poplar (*Populus tomentosa* Carr.) resistance to *Dothiorella gregaria* was clearly impacted by overexpression and mutation of an R2R3 MYB transcription factor, respectively [45]. Additional evidence shows that chemical inhibition of a CHS enzyme and downregulation of its corresponding gene suppressed *Podosphaera xanthii* induced resistance in cucumber (*Cucumis sativus* L.).

“The antifungal activity of phenylpropanoids and flavonoids is supported by metabolomic and functional genomic data. Flavonoids extracted from the needles of *Picea neoveitchii* Mast. used at 1 mg/mL, exhibited very interesting antifungal activities: kaempferol-7-O- (2"-E-p-coumaroyl) - $\alpha$ -L-arabinofuranoside exhibited strong activity against *Fusarium oxysporum* with a relative inhibitory percentage of 108.1%, while 5,7,4'-trihydroxy-3,8,-di- methoxy-6-C-methylflavone, 5,8,4'-trihydroxy-3,7-dimethoxy-6-C-methylflavone, 7-methoxy-6-C-methylkaempferol and kaempferol-7-O-(2"-E-p-coumaroyl)- $\alpha$ -L-arabino- furanoside were active against *Rhizoctonia solani*, with 49.5%, 53.3%, 95.3% and 49.5% relative inhibitory percentages, respectively” [46]. “Compared to Carbendazim, a synthetic chemical fungicide used to treat these two pathogens, these compounds proved to be as active as Carbendazim. In addition, other flavonoids, such as eriodictyol, homoeriodictyol, dihydroquercetin, and luteolin, were isolated from *Ficus sarmentosa*. It was found that Henryi (King) Corner was highly effective against pathogenic fungi, including *Fusarium graminearum* and *Septoria zeicola*. Luteolin showed the strongest antagonistic activity among these flavonoids, with half-maximal inhibitory concentrations (IC<sub>50</sub>) of 56.38 and 81.48 mg·L<sup>-1</sup> for each fungus, respectively” [47]. “In an in vivo assay, Cherry tomatoes sprayed with Laurel (*Laurus nobilis* L.) oil containing 44% eugenol and 30% cinnamaldehyde were less infected by *Alternaria alternata* after 5 days of storage. In particular, decayed tomatoes treated with laurel oil were reduced by 86.4% [48]. A few studies have assessed the integrity of plant fungal pathogens using phenylpropanoid derivatives. One of them is the study shown above on essential oil from laurel leaves: fungicidal activity was demonstrated by invaginations and folds in the fungus cell walls”. [59]

## 2.2 Bacterial pathogens

There have been many studies investigating how PPP affects bacterial resistance. It has been shown that infection increases the content of flavonoids and other phenylpropanoid derivatives, especially coumaric acid, in Tobacco (*Nicotiana tabacum* L.)-*Pseudomonas syringae* pathosystems [27]. Orange leaves (*Citrus sinensis* L.) infected with *Candidatus liberibacter asiaticus* produced significantly more flavonoid glycosides and hydroxycinnamic acids [49]. A necrotrophic bacterial pathogen, *Pectobacterium atrosepticum*, was shown to be resistant to rutin (quercetin-3-O-rutinoside) and nicotiflorin (kaempferol-3-O-rutinoside) in Potato tubers [38]. A higher concentration of *pelargonidin-3-O-rutinoside-5-O-glucopyranoside* and *peonidin-3-O-rutinoside*, both acylated with p-coumaric acid, was found in the *E. carotovora* of transgenic potato tubers [50].

In the context of plant protection, phenylpropanoids have been heavily studied in vitro to develop biological alternatives to synthetic phytoprotectants. Nevertheless, [51] claims that efficient protection is not the result of significant toxicity but rather of the host's defense system. Therefore, it is important to explore both the direct and indirect modes of action of these compounds for effective pest management. Consequently, it is imperative to explore their biosynthesis and targets. Post-inoculation transcriptomic shifts revealed that some genes encoding enzymes in the PPP are rapidly activated. However, others are induced much later in the infection response [52, 27]. Among the earliest genes activated were CHS, F3H and isoflavone synthase I (IFS I) in soybeans infected by *Pseudomonas syringae* [52]. A number of PAL genes were found in tobacco infected by the same bacterium, including cinnamic acid 4-hydroxylase (C4H), 4-coumarate: CoA-ligase (4CL), Orthomethyltransferases (OMT) and ferulate-5-hydroxylase (F5H). Although this discovery may pave the way for the development of new plant varieties resistant to bacteria, little is known about how these

enzymes work biologically. PPP compounds have been reported to disrupt vital cellular functions and jeopardise bacterial survival without detailing the mechanisms. In *Clavibacter michiganensis*, fragarin from strawberry leaves disrupted cell membrane integrity [1].

The electron transport pathway and cell survival, in particular, can be significantly affected by compounds with better penetrability. A case study on *Micrococcus luteus* demonstrated that two retrochalcones isolated from Chinese licorice (*Glycyrrhiza inflata* L.) roots compromised the enzymatic activity of the NADH-cytochrome C reductase with licochalcones A and C. A compound's antibacterial activity may also be attributed to preventing cell proliferation through direct interference with cell division. Chlorogenic acid inhibits cell division by inhibiting the polymerization of the cytoskeletal protein [25], which is essential for cytokinesis. It has been suggested that chlorogenic acid forms hydrogen bonds and hydrophobic interactions with various residues of this protein, altering its conformation and inhibiting GTPase activity [1].

### 2.3 Viruses

In response to viral agents, plants produced salicylic acid (SA) and other defense metabolites from the PPP to initiate systemic acquired resistance (SAR). Viruses such as Sugarcane mosaic and Dwarf mosaic virus have been shown to cause Dwarf mosaic diseases in Maize [53]. Tobacco mosaic virus (TMV) has been investigated in the exploration of the antiviral activity of candidate specialized metabolites owing to its host diversity, which includes several economically important plant species, including Tobacco and Tomato (*Solanum lycopersicum* L.). Although 5-O-caffeoylquinic acid and quercetin were abundant at the TMV infection site in Tobacco leaves, kaempferol was predominant in a distant part of the plant with SAR [1]. In most publications, quercetin and kaempferol are cited as triggers of the plant's defense response rather than direct actions on viral particles. Such as *Datura stramonium* L.-TMV and *Chenopodium amaranticolor* -TMV pathosystems [54]. Several studies suggest that metabolite levels are associated with plant resistance to viruses, but they do not assess whether they are potentially harmful to viruses.

### 2.4 Insects

The insects are the most problematic macroscopic pests on cultivated plants, not just because of the direct damage caused by their herbivorous larvae but also because of the indirect damage caused by their ability to spread microbial pathogens. A number of studies have been conducted to identify plants' natural defense mechanisms against herbivores. Defensive compounds are produced constitutively or in response to plant damage and affect herbivore feeding, growth, and survival. Aside from releasing volatile organic compounds, plants also attract herbivores' natural enemies. Phenylpropanoids and flavonoids are defensive compounds against insects. It has been shown that chlorogenic acid and feruloylquinic acid distinguish between resistant and susceptible chrysanthemum genotypes and Thrips (*Frankliniella occidentalis*). Both molecules have higher amounts in thrips-resistant genotypes. The wild-cultivated crosses of Groundnut plants (*Arachis hypogaea* L. X *Arachis kempff-mercadoi* Krapov.) contain similar amounts of quercetin, chlorogenic acid and rutin [25].

In carrot leaves, the flavone, luteolin and the phenylpropanoid, sinapic acid significantly distinguished thrips-resistant cultivars from susceptible ones. The ratio of specific metabolites played a preponderant role in plant resistance to insects, as in Carrot, which showed resistance to the fly *Psylla rosae* [1]. *F. occidentalis* thrips were shown to be significantly harmed by chlorogenic acid when they were fed artificial diets containing 5% chlorogenic acid [25]. Castor bean leaf extracts (*Ricinus communis* L.) contain quercetin, which can kill adults and also prevent oviposition and ovipos from developing. One of the most common Bean weevils, *Callosobruchus chinensis* L., is known as a pest to many stored legumes [55]. While some molecules do not directly have insecticidal activity, their physical and chemical properties can enhance other compounds solubility, improving their penetration and effectiveness. This kind of synergy was illustrated with non-PPP metabolites in an in vitro analysis that revealed an up to 19-fold increase in penetration of camphor in a binary mixture with 1,8-cineole through the larval integument of the Cabbage looper (*Trichoplusia ni*) compared to Camphor alone. As shown by the LD50 for 1,8-cineole: camphor of 60:40 (186.9 g/insect) [56, 57], such a synergy was found in PPP metabolites. One metabolite may act in opposing ways against different targets, as we have discussed previously for fungal targets. For instance, chlorogenic acid

may function as an auxiliary to control thrips but on the other hand, promote oviposition by the Black swallowtail (*Papilio polyxenes* Fabr.). It is therefore necessary to adopt an integrative approach to protect against one target without increasing the severity caused by another [25].

Insects can cause direct damage to plants, as well as be vectors of economically threatening diseases, including Pierce's disease caused by *Xylella fastidiosa* [58], Grapevine yellow disease caused by phytoplasmas [59] or other major crop viruses. A major concern in crop management is preventing host contact with disease-carrying vectors in order to limit epidemics. As a result, Su and collaborators [60] linked metabolic changes in Tomato leaves with vector behaviour to address this dimension. The whiteflies (*Bemisia tabaci*) recognised plants that had previously been attacked by conspecifics due to decreased terpenoid and flavonoid contents. By treating Tomato plants infested by *B. tabaci* with naringenin they increased their content of rutin, kaempferol-rhamnopyranoside, quercetin-trisaccharide, 3-O-methylmyricetin and anthocyanin to the same level as those measured in non-infested plants. Later studies showed that the preference of *B. tabaci* for oviposition on previously infested plants was reversed. As a consequence of a reduced *B. tabaci* population, both the pest and the vectored virus (e.g., the Tomato yellow leaf curl virus (TYLCV)) that cause damage to the plant can be decreased. Moreover, whiteflies fed less on flavonoid-rich Tomato leaves, and TYLCV spread was reduced [61]. Their findings suggest that flavonoids are more likely to inhibit host–vector interaction than to have antiviral activity, since only disease expression was delayed. Flavonoids were also accumulated in grapevines after *phytoplasma* infection. Flavonoids were thought to repel insects [62].

### 3. Conclusion

A complex plant-environment interaction mediated in part by secondary metabolites, including phenylpropanoids and flavonoids, results in diversity within the structural and regulatory genes of several superfamilies involved in secondary metabolism. Ultimately, the desired systems biology approach will be realised through rapid progress in sequencing, structural elucidation, and analytical tools. Several model plants and crops, like Arabidopsis, alfalfa, poplar, and rice, could be used at the beginning. A comprehensive understanding of the pathway leading to phenylpropanoids and flavonoids formation and function should continue to be explored.

### 4. References

1. Mierziak, J.; Kostyn, K.; Kulma, A. Flavonoids as Important Molecules of Plant Interactions with the Environment. *Molecules* 2014, *19*, 16240–16265.
2. Falcone Ferreyra, M.L.; Rius, S.P.; Casati, P. Flavonoids: Biosynthesis, Biological Functions, and Biotechnological Applications. *Front. Plant Sci.* 2012, *3*, 222.
3. Bily, A.C.; Reid, L.M.; Taylor, J.H.; Johnston, D.; Malouin, C.; Burt, A.J.; Bakan, B.; Regnault-Roger, C.; Pauls, K.P.; Arnason, J.T.; et al. Dehydrodimers of Ferulic Acid in Maize Grain Pericarp and Aleurone: Resistance Factors to *Fusarium graminearum*. *Phytopathology* 2003, *93*, 712–719.
4. Samanta, A.; Das, G.; Das, S. Roles of Flavonoids in Plants. *Carbon* 2011, *100*, 12–35.
5. Tak, J.H.; Isman, M.B. Penetration-Enhancement Underlies Synergy of Plant Essential Oil Terpenoids as Insecticides in the Cab-bage Looper, *Trichoplusia Ni*. *Sci. Rep.* 2017, *7*, 42432.
6. Sampietro, D.A.; Fauguel, C.M.; Vattuone, M.A.; Presello, D.A.; Catalán, C.A.N. Phenylpropanoids from Maize Pericarp: Resistance Factors to Kernel Infection and Fumonisin Accumulation by *Fusarium Verticillioides*. *Eur. J. Plant Pathol.* 2013, *135*, 105–113.
7. Vogt, T. Phenylpropanoid Biosynthesis. *Mol. Plant* 2010, *3*, 2–20.
8. Schulz, E.; Tohge, T.; Zuther, E.; Fernie, A.R.; Hincha, D.K. Flavonoids Are Determinants of Freezing Tolerance and Cold Acclimation in Arabidopsis Thaliana. *Sci. Rep.* 2016, *6*, 34027.
9. Kratočić, E.; Rusak, G.; Bezić, N.; Krajčić, M. Inhibition of Tobacco Mosaic Virus Infection by Quercetin and Vitexin. *Acta Virol.* 2008, *52*, 119–124.
10. Crasci, L.; Basile, L.; Panico, A.; Puglia, C.; Bonina, F.P.; Basile, P.M.; Rizza, L.; Guccione, S. Correlating in Vitro Target-Oriented Screening and Docking: Inhibition of Matrix Metalloproteinases Activities by Flavonoids. *Planta Med.* 2017, *83*, 901–911.
11. Wang, X. gui; Wei, X. yi; Tian, Y. qing; Shen, L. tao; Xu, H. hong Antifungal Flavonoids from *Ficus sarmentosa* var. *Henryi* (King) Corner. *Agric. Sci. China* 2010, *9*, 690–694.
12. Xu, S.; Yan, F.; Ni, Z.; Chen, Q.; Zhang, H.; Zheng, X. In Vitro and in Vivo Control of Alternaria Alternata in Cherry Tomato by Essential Oil from *Laurus Nobilis* of Chinese Origin. *J. Sci. Food Agric.* 2014, *94*,
13. Stevenson, P.C.; Nicolson, S.W.; Wright, G.A. Plant Secondary Metabolites in Nectar: Impacts on Pollinators and Ecological Functions. *Funct. Ecol.* 2017, *31*, 65–75.

14. Feeny, P.; Sachdev, K.; Rosenberry, L.; Carter, M. Luteolin 7-O-(6"-O-Malonyl)- $\beta$ -d-Glucopyranoside and Trans-Chlorogenic Acid: Oviposition Stimulants for the Black Swallowtail Butterfly. *Phytochemistry* 1988, 27, 3439–3448.
15. Pichersky, E.; Lewinsohn, E. Convergent Evolution in Plant Specialized Metabolism. *Annu. Rev. Plant Biol.* 2011, 62, 549–566.
16. Hodaie, M.; Rahimmalek, M.; Arzani, A.; Talebi, M. The Effect of Water Stress on Phytochemical Accumulation, Bioactive Compounds and Expression of Key Genes Involved in Flavonoid Biosynthesis in *Chrysanthemum morifolium* L. *Ind. Crops Prod.* 2018, 120, 295–304.
17. Shadle, G.L.; Wesley, S.V.; Korth, K.L.; Chen, F.; Lamb, C.; Dixon, R.A. Phenylpropanoid Compounds and Disease Resistance in Transgenic Tobacco with Altered Expression of L-Phenylalanine Ammonia-Lyase. *Phytochemistry* 2003, 64, 153–161.
18. Kim, J.; Buel, C.R. A Revolution in Plant Metabolism: Genome-Enabled Pathway Discovery. *Plant Physiol.* 2015, 169, 1532–1539.
19. Agati, G.; Biricolti, S.; Guidi, L.; Ferrini, F.; Fini, A.; Tattini, M. The Biosynthesis of Flavonoids Is Enhanced Similarly by UV Radiation and Root Zone Salinity in *L. vulgare* Leaves. *J. Plant Physiol.* 2011, 168, 204–212.
20. Su, Q.; Chen, G.; Mescher, M.C.; Peng, Z.; Xie, W.; Wang, S.; Wu, Q.; Liu, J.; Li, C.; Wang, W.; et al. Whitefly Aggregation on Tomato Is Mediated by Feeding-Induced Changes in Plant Metabolites That Influence the Behaviour and Performance of Con-specifics. *Funct. Ecol.* 2018, 32, 1180–1193.
21. Pott, D.M.; Osorio, S.; Vallarino, J.G. From Central to Specialized Metabolism: An Overview of Some Secondary Compounds Derived from the Primary Metabolism for Their Role in Conferring Nutritional and Organoleptic Characteristics to Fruit. *Front. Plant Sci.* 2019, 10, 835.
22. Rackova, L.; Firakova, S.; Kostalova, D.; Stefek, M.; Sturdik, E.; Majekova, M. Oxidation of Liposomal Membrane Suppressed by Flavonoids: Quantitative Structure-Activity Relationship. *Bioorganic Med. Chem.* 2005, 13, 6477–6484.
23. Koutouan, C.; Clerc, V. Le; Baltenweck, R.; Claudel, P.; Halter, D.; Hugueney, P.; Hamama, L.; Suel, A.; Huet, S.; Merlet, M.H.B.; et al. Link between Carrot Leaf Secondary Metabolites and Resistance to *Alternaria Daudi*. *Sci. Rep.* 2018, 8, 13746.
24. Marie, L.R.; Claude K.; Jean, J.H.; Valerie, L.C.; Latifa, H.; Emmanuel G.; Mathilde, B.; Role of Phenylpropanoids and flavonoids in plant resistance to pests and diseases. *Molecules.* 2022,27,8371.
25. Upasani, S.M.; Kotkar, H.M.; Mendki, P.S.; Maheshwari, V.L. Partial Characterization and Insecticidal Properties of Ricinus Communis L Foliage Flavonoids. *Pest Manag. Sci.* 2003, 59, 1349–1354.
26. Tissier, A.; Ziegler, J.; Vogt, T. Specialized Plant Metabolites: Diversity and Biosynthesis. In *Ecological Biochemistry: Environmental and Interspecies Interactions*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2015; pp. 14–37.
27. Haraguchi, H.; Yoshida, N.; Ishikawa, H.; Tamura, Y.; Mizutani, K.; Kinoshita, T. Protection of Mitochondrial Functions against Oxidative Stresses by Isoflavans from *Glycyrrhiza Glabra*. *J. Pharm. Pharmacol.* 2010, 52, 219–223.
28. Stoepler, T.M.; Wolf, T.K. North American Grapevine Yellows Disease: Current Knowledge and Management Recommendations for Wine Growers; Virginia Cooperative Extension: Blacksburg, United States, Publication AREC-48P; 2013.
29. Bollina, V.; Kumaraswamy, G.K.; Kushalappa, A.C.; Choo, T.M.; Dion, Y.; Rioux, S.; Faubert, D.; Hamzehzarghani, H. Mass Spectrometry-Based Metabolomics Application to Identify Quantitative Resistance-Related Metabolites in Barley against Fusarium Head Blight. *Mol. Plant Pathol.* 2010, 11, 769–782.
30. Rees, A.; Dodd, G.F.; Spencer, J.P.E. The Effects of Flavonoids on Cardiovascular Health: A Review of Human Intervention Trials and Implications for Cerebrovascular Function. *Nutrients* 2018, 10, 1852.
31. Samanta, A.; Das, G.; Das, S. Roles of Flavonoids in Plants. *Carbon* 2011, 100, 12–35.
32. Song, Z.; Chen, W.; Du, X.; Zhang, H.; Lin, L.; Xu, H. Chemical Constituents of *Picea neoveiticii*. *Phytochemistry* 2011, 72, 490–494.
33. Del Rio, J.A.; Gonzalez, A.; Fuster, M.D.; Botia, J.M.; Gomez, P.; Frias, V.; Ortuño, A. Tylose Formation and Changes in Phenolic Compounds of Grape. *Phytopathol. Mediterr.* 2001, 40, 394–399.
34. Bruneton, J. *Pharmacognosie: Phytochimie et Plantes Médicinales*; Tec & doc Cachan: Paris, France, 1999; ISBN 2-7430-0315-4.
35. Kröner, A.; Marnet, N.; Andrivon, D.; Val, F. Nicotiflorin, Rutin and Chlorogenic Acid: Phenylpropanoids Involved Differently in Quantitative Resistance of Potato Tubers to Biotrophic and Necrotrophic Pathogens. *Plant Physiol. Biochem.* 2012, 57, 23–31.
36. Tak, J.H.; Isman, M.B. Enhanced Cuticular Penetration as the Mechanism for Synergy of Insecticidal Constituents of Rosemary Essential Oil in *Trichoplusia Ni*. *Sci. Rep.* 2015, 5, 12690.
37. Lorenc-Kukuła, K.; Jafra, S.; Oszmiański, J.; Szopa, J. Ectopic Expression of Anthocyanin 5-O-Glucosyltransferase in Potato Tu- ber Causes Increased Resistance to Bacteria. *J. Agric. Food Chem.* 2005, 53, 272–281.
38. Naoumkina, M.A.; Zhao, Q.; Gallego-Giraldo, L.; Dai, X.; Zhao, P.X.; Dixon, R.A. Genome-Wide Analysis of Phenylpropanoid Defense Pathways. *Mol. Plant Pathol.* 2010, 11, 829–846.
39. Dudareva, N.; Negre, F.; Nagegowda, D.A.; Orlova, I. Plant Volatiles: Recent Advances and Future Perspectives. *CRC. Crit. Rev. Plant Sci.* 2006, 25, 417–440.
40. Hijaz, F.M.; Manthey, J.A.; Folimonova, S.Y.; Davis, C.L.; Jones, S.E.; Reyes-De-Corcuera, J.I. An HPLC-MS Characterization of the Changes in Sweet Orange Leaf Metabolite Profile Following Infection by the Bacterial Pathogen *Candidatus Liberibacter Asiaticus*. *PLoS ONE* 2013, 8, e79485.
41. Bednarek, P. Chemical Warfare or Modulators of Defense Responses—The Function of Secondary

- Metabolites in Plant Immun-ity. *Curr. Opin. Plant Biol.* 2012, 15, 407–414.
42. Yogendra, K.N.; Kushalappa, A.C.; Sarmiento, F.; Rodriguez, E.; Mosquera, T. Metabolomics Deciphers Quantitative Resistance Mechanisms in Diploid Potato Clones against Late Blight. *Funct. Plant Biol.* 2015, 42, 284–298.
  43. Szatmári, Á.; Zvara, Á.; Móricz, Á.M.; Besenyi, E.; Szabó, E.; Ott, P.G.; Puskás, L.G.; Bozsó, Z. Pattern Triggered Immunity (PTI) in Tobacco: Isolation of Activated Genes Suggests Role of the Phenylpropanoid Pathway in Inhibition of Bacterial Pathogens. *PLoS ONE* 2014, 9, e102869.
  44. Yang, W.; Xu, X.; Li, Y.; Wang, Y.; Li, M.; Wang, Y.; Ding, X.; Chu, Z. Rutin-Mediated Priming of Plant Resistance to Three Bacterial Pathogens Initiating the Early SA Signal Pathway. *PLoS ONE* 2016, 11, e0146910.
  45. Tak, J.H.; Isman, M.B. Enhanced Cuticular Penetration as the Mechanism for Synergy of Insecticidal Constituents of Rosemary Essential Oil in *Trichoplusia Ni.* *Sci. Rep.* 2015, 5, 12690.
  46. Yao, Q.; Peng, Z.; Tong, H.; Yang, F.; Xing, G.; Wang, L.; Zheng, J.; Zhang, Y.; Su, Q. Tomato Plant Flavonoids Increase Whitefly Resistance and Reduce Spread of Tomato Yellow Leaf Curl Virus. *J. Econ. Entomol.* 2019, 112, 2790–2796.
  47. Zabala, G.; Zou, J.; Tuteja, J.; Gonzalez, D.O.; Clough, S.J.; Vodkin, L.O. Transcriptome Changes in the Phenylpropanoid Pathway of Glycine Max in Response to *Pseudomonas Syringae* Infection. *BMC Plant Biol.* 2006, 6, 26.
  48. Jeandet, P.; Hébrard, C.; Deville, M.A.; Cordelier, S.; Dorey, S.; Aziz, A.; Crouzet, J. Deciphering the Role of Phytoalexins in Plant-Microorganism Interactions and Human Health. *Molecules* 2014, 19,
  49. Margaria, P.; Ferrandino, A.; Caciagli, P.; Kedrina, O.; Schubert, A.; Palmano, S. Metabolic and Transcript Analysis of the Flavonoid Pathway in Diseased and Recovered Nebbiolo and Barbera Grapevines (*Vitis Vinifera* L.) Following Infection by *Flavescence Dorée* Phytoplasma. *Plant Cell Environ.* 2014, 37, 2183–2200.
  50. Zagrean-Tuza, C.; Mot, A.C.; Chmiel, T.; Bende, A.; Turcu, I. Sugar Matters: Sugar Moieties as Reactivity-Tuning Factors in Quercetin: O-Glycosides. *Food Funct.* 2020, 11, 5293–5307.
  51. Leiss, K.A.; Maltese, F.; Choi, Y.H.; Verpoorte, R.; Klinkhamer, P.G.L. Identification of Chlorogenic Acid as a Resistance Factor for Thrips in Chrysanthemum. *Plant Physiol.* 2009, 150, 1567–1575.
  52. Wang, Y.C.; Qian, W.J.; Li, N.N.; Hao, X.Y.; Wang, L.; Xiao, B.; Wang, X.C.; Yang, Y.J. Metabolic Changes of Caffeine in Tea Plant (*Camellia Sinensis* (L.) O. Kuntze) as Defense Response to *Colletotrichum fructicola*. *J. Agric. Food Chem.* 2016, 64, 6685–6693.
  53. Tugizimana, F.; Djami-Tchatchou, A.T.; Steenkamp, P.A.; Piater, L.A.; Dubery, I.A. Metabolomic Analysis of Defense-Related Reprogramming in Sorghum Bicolor in Response to *Colletotrichum sublineolum* Infection Reveals a Functional Metabolic Web of Phenylpropanoid and Flavonoid Pathways. *Front. Plant Sci.* 2019, 9, 1840.
  54. Treutter, D. Significance of Flavonoids in Plant Resistance: A Review. *Environ. Chem. Lett.* 2006, 4, 147–157.
  55. Ardila, H.D.; Martínez, S.T.; Higuera, B.L. Levels of Constitutive Flavonoid Biosynthetic Enzymes in Carnation (*Dianthus Caryophyllus* L.) Cultivars with Differential Response to *Fusarium Oxysporum* f. Sp. *Dianthi.* *Acta Physiol. Plant.* 2013, 35, 1233–1245.
  56. Tan, B.A.; Daim, L.D.J.; Ithnin, N.; Ooi, T.E.K.; Md-Noh, N.; Mohamed, M.; Mohd-Yusof, H.; Appleton, D.R.; Kulaveerasingam, H. Expression of Phenylpropanoid and Flavonoid Pathway Genes in Oil Palm Roots during Infection by *Ganoderma Boninense*. *Plant Gene* 2016, 7, 11–20.
  57. Tattini, M.; Gravano, E.; Pinelli, P.; Mulinacci, N.; Romani, A. Flavonoids Accumulate in Leaves and Glandular Trichomes of *Phillyrea Latifolia* Exposed to Excess Solar Radiation. *New Phytol.* 2000, 148, 69–77.
  58. Panche, A.N.; Diwan, A.D.; Chandra, S.R. Flavonoids: An Overview. *J. Nutr. Sci.* 2016, 5, e47.
  59. Ramarosan ML, Koutouan C, Helesbeux JJ, Le Clerc V, Hamama L, Geoffriau E, Briard M. Role of phenylpropanoids and flavonoids in plant resistance to pests and diseases. *Molecules.* 2022 Nov 30;27(23):8371.